

Orthogonal Dualities and Asymptotics of Dynamic Stochastic Higher Spin Vertex Models, using the Drinfeld Twister

Jeffrey Kuan and Zhengye Zhou

Accessibility Statement:

A WCAG2.1AA compliant version of this PDF will be available at the first author's professional website.

This PDF was created in consultation with the British Dyslexia Association's Style Guide 2023; however, arXiv administrators do not allow the dyslexia-friendly color contrast.

For any issues accessing the information in this document, please email the first author.

Abstract

We introduce a new, algebraic method to construct duality functions for integrable dynamic models. This method will be implemented on dynamic stochastic higher spin vertex models, where we prove the duality functions between the dynamic stochastic higher spin vertex models and non dynamic stochastic higher spin vertex models are the ${}_3\varphi_2$ functions. A degeneration of these duality functions is dual q -Krawtchouk polynomials, which agree with orthogonal polynomial dualities between ASEP and dynamic ASEP of Groenevelt–Wagenaar [19].

The method involves using the universal twister of $U_q(sl_2)$, viewed as a quasi-triangular, quasi- $*$ -Hopf algebra. The algebraic method is presented very generally and is expected to produce duality functions for other dynamic integrable models.

As an application of the duality, we prove that the asymptotic fluctuations of the dynamic stochastic six vertex model with step initial conditions are governed by the Tracy–Widom distribution.

Contents

1	Introduction	3
2	Background	6
2.1	Orthogonal Polynomial Background	6
2.2	Probabilistic Background	8
2.2.1	Definition of duality	8
2.2.2	Definition of model	10
2.3	Algebraic Background	12
2.3.1	Quantum groups as quasi-triangular quasi-Hopf *-algebras	12
3	Main Results	16
3.1	Probabilistic Results	16
3.1.1	Markov Duality	17
3.1.2	Asymptotics	21
3.2	Algebraic Results	22
4	Proofs	27
4.1	Proof of Theorem 2 (assuming Theorem 1)	27
4.2	A general result using orthogonality	27
4.2.1	Matching notation to the dynamic stochastic six vertex model	31
4.2.2	A “trick” for rewriting the duality function	32
4.2.3	Proving the Bounds	36
4.3	Proof of Theorem 3	39
4.3.1	Yang-Baxter-Equation	39
4.3.2	Reducing from \mathcal{B} to B	40

4.3.3	Transposition	41
4.4	Proof of Proposition 1	42
4.4.1	Transposition formula	42
4.4.2	The Ground State Transformation	43
4.4.3	Inversion formula	44
4.5	Proof of Theorem 4 and Corollary 3	45
4.6	Proof of Proposition 2	47
4.6.1	Calculating limits of $B_{m,l}(u, z, \lambda)$	48
4.6.2	Calculating The Ground State Transformation	52
4.6.3	Proof of Corollary 1	53
4.7	Proof of Proposition 3	55
4.7.1	Matching of Notation	56
4.7.2	Proof in the General Case:	57
5	Orthogonal Polynomial Duality	58
5.1	The dual quantum group	59
5.2	Proof of Orthogonal Polynomial Duality	60
6	Appendix	64
6.1	Examples of Matching	64
6.1.1	The spin 1/2 case	64
6.1.2	The spin 1 case	66
6.2	Computer Simulations	68

1 Introduction

Markov duality is a widely used tool in the theory of Markov processes, wherein an analysis of a Markov process is reduced to an analysis of a “simpler” Markov process. However, despite its ubiquity, discovering Markov duality is a “black art”[20]. Over the last 30 years, starting with

the seminal papers [34, 35], researchers have developed an algebraic machinery to discover Markov duality in interacting particle systems.

In more recent years, there have been new developments on so-called “dynamic” models. Dynamic stochastic vertex models and dynamic ASEP were introduced by Borodin [6], using the Felder–Varchenko elliptic quantum group. His result produced an contour integral formula for expectations of certain observables which resembled a duality result. Later work of [7] proved that the observable is indeed a duality function between dynamic ASEP and usual ASEP, using direct calculations involving q -Hermite polynomials. Later work of Aggarwal [1] applied fusion to produce dynamic stochastic higher spin vertex models with explicit formulas for the weights, and also produced contour integral formulas for expectations. An unusual feature of these dynamic models is that the one point marginals have scaling exponent $1/4$ and converge to a non-trivial random variable. (See also the SPDE work of [13] where the scaling limit occurs).

In this paper, the main probabilistic result is a duality for the dynamic stochastic higher spin vertex models, and Tracy–Widom fluctuations for the dynamic stochastic six vertex model. The duality functions are themselves vertex weights, which was an idea communicated to the authors by Michael Wheeler [36, 37] and William Mead [33].

The proof utilizes algebraic machinery. In particular, the new ingredients are (1) the universal Drinfeld twister of the quantum group $U_q(sl_2)$, in contrast to the use of the Felder–Varchenko elliptic quantum group; and (2) calculations are done using the universal R -matrix rather than applying fusion. As far as we know, this is the first probabilistic application of the previous two tools. The algebraic methods are expected to apply to other Drinfeld–Jimbo quantum groups.

We also make a remark on orthogonal polynomial duality, where the

duality functions are polynomials which are orthogonal with respect to the reversible measures of the process. Because of the orthogonality, there are **a priori** applications in probability, such as to Boltzmann-Gibbs principles [3, 4] or correlation formulas [11, 30]. In this sense, orthogonal polynomial dualities are in improvement on “triangular” dualities, for which the probabilistic applications are not immediately obvious. A recent paper [16] developed a new algebraic method for constructing orthogonal polynomial dualities in interacting particle systems, generalizing the previous methods of [10, 11, 18, 38]. Because the method here involves Michael Wheeler’s idea that duality functions are themselves vertex weights, new algebraic insights for orthogonal polynomial dualities will be needed. These are discussed in section 5.

Shortly after version 1 of this paper appeared on the arXiv, a preprint [19] of Groenevelt and Wagenaar constructed a dynamic ASEP(q, j) with orthogonal polynomial duality (with respect to the reversible measures). In version 2 of this paper, we show that their duality function is a degeneration of ours. We then use the duality function to prove Tracy–Widom fluctuations. To our knowledge, this is the first time that orthogonal polynomials have been used for asymptotic fluctuations.

Acknowledgements. The authors would like to thank (in no particular order) Michael Wheeler, William Mead, Wolter Groenevelt, Alexei Borodin, Amol Aggarwal, Ivan Corwin, Carel Wagenaar, Konstantin Matetski, and Leonid Petrov for helpful discussions. The discussions with Wolter Groenevelt and William Mead were held at the workshop “Recent Developments in Stochastic Duality” in Eindhoven, with support from EURANDOM and the NWO grant 613.001.753 “Duality for interacting particle systems.”

The first author would also like to thank his friends on “CATS (Cult of Accessibility and Taylor Swift)”, a Swiftie Discord Server where we

discuss Taylor Swift and accessibility (acknowledging that this PDF is not accessible). Personal thanks goes to @Vega for setting up the server. The first author was also supported by National Science Foundation grant DMS-2000331.

2 Background

2.1 Orthogonal Polynomial Background

When n is a nonnegative integer, the q -Pochhammer symbol is defined by:

$$(a; q)_n = \prod_{j=0}^{n-1} (1 - q^j a) \quad (a; q)_{-n} = \prod_{j=1}^n \frac{1}{1 - q^{-j} a}.$$

From the q -Pochhammer symbols, we can define the hypergeometric series

$${}_p\varphi_r \left(\begin{matrix} a_1, a_2, \dots, a_p \\ b_1, b_2, \dots, b_r \end{matrix} \middle| q, z \right) = \sum_{k=0}^{\infty} \frac{z^k}{(q; q)_k} \prod_{j=1}^p (a_j; q)_k \prod_{j=1}^r (b_j; q)_k^{-1}.$$

For any integer n , define the q -deformed integers and factorial by

$$[n] = \frac{q^n - q^{-n}}{q - q^{-1}}, \quad [n]! = \frac{(q^2; q^2)_n}{(q^{-1} - q)^n} q^{-n(n+1)/2}.$$

We extend this definition such that for any $a \in \mathbb{R}$ and integer n ,

$$\frac{[a+n]!}{[a]!} = \frac{(q^{2(a+1)}; q^2)_n}{(q^{-1} - q)^n} q^{-na - n(n+1)/2}.$$

Now we introduce the Racah-Wilson polynomials [2], which is given by the q -hypergeometric function:

$$W_n(x; \alpha, \beta, \gamma, M|q) = {}_4\varphi_3 \left(\begin{matrix} q^{-n}, q^{n+1}\alpha\beta, q^{-x}, q^{x-M}\gamma \\ q\alpha, q\beta\gamma, q^{-M} \end{matrix} \middle| q, q \right). \quad (1)$$

It satisfies the following orthogonal relation:

$$\sum_{x=0}^N W_n(x; \alpha, \beta, \gamma, M|q) W_m(x; \alpha, \beta, \gamma, M|q) w(x) = \delta_{m,n} h_n, \quad (2)$$

where

$$h_n = \frac{(q; q)_n (1 - \alpha\beta q) (\beta q; q)_n (\alpha\gamma^{-1} q; q)_n (\alpha\beta q^{M+2}; q)_n (q^{-M}\gamma)^n}{(\alpha\beta q; q)_n (1 - \alpha\beta q^{2n+1}) (\alpha q; q)_n (\beta\gamma q; q)_n (q^{-M}; q)_n} \\ \times \frac{(q^{-M+1}\gamma; q)_\infty (\alpha^{-1}\beta^{-1}q^{-M-1}; q)_\infty (\alpha^{-1}\gamma; q)_\infty (\beta^{-1}; q)_\infty}{(\alpha^{-1}q^{-M}\gamma; q)_\infty (\beta^{-1}q^{-M}; q)_\infty (\gamma q; q)_\infty (\alpha^{-1}\beta^{-1}q^{-1}; q)_\infty}, \\ w(x) = \frac{(q^{-M}\gamma; q)_x (1 - q^{-M+2x}\gamma) (\alpha q; q)_x (\beta\gamma q; q)_x (q^{-M}; q)}{(q; q)_x (1 - q^{-M}\gamma) (\alpha^{-1}q^{-M}\gamma; q)_x (\beta^{-1}q^{-M}; q)_x (\gamma q; q)_x (\alpha\beta q)^x}.$$

Next, we define the $q - 6j$ symbol [22]

$$\left\{ \begin{matrix} a & b & e \\ d & c & f \end{matrix} \right\}_q \\ = (-1)^{2c+2d+2e} \sqrt{[2e+1][2f+1]} \Delta(a, b, e) \Delta(a, c, f) \Delta(c, e, d) \Delta(d, b, f) \\ \times \sum_w \left(\frac{(-1)^{-w} [a+b+c+d+1-w]!}{[c+d-e-w]! [b+d-f-w]! [a+c-f-w]! [a+b-e-w]!} \right. \\ \left. \times \frac{1}{[w]! [w+e+f-b-c]! [w-a-d+e+f]!} \right), \quad (3)$$

where the sum is taken only over w with non-negative arguments in square brackets inside the summation and

$$\Delta(j_1, j_2, j_3) = \sqrt{\frac{[-j_1 + j_2 + j_3]! [j_1 - j_2 + j_3]! [j_1 + j_2 - j_3]!}{[j_1 + j_2 + j_3 + 1]!}}.$$

The $q - 6j$ symbol can be expressed through the Racah-Wilson polyno-

mial, by setting

$$\begin{aligned} n &= a + b - e, & x &= c + d - e, & M &= a + b + c + d + 1, \\ \alpha &= q^{2(-a-d+e+f)}, & \beta &= q^{2(-a-b-c+d+1)}, & \gamma &= q^{2(a+e+f-d+1)} \end{aligned}$$

so that

$$\left\{ \begin{array}{ccc} a & b & e \\ d & c & f \end{array} \right\}_q = \sqrt{\frac{w(x)}{h_n}} W_n(x; \alpha, \beta, \gamma, M | q^2). \quad (4)$$

The dual q -Krawtchouk polynomial $K_n(x; c, N | q)$ is defined as

$$K_n(x; c, N | q) = {}_3\phi_2 \left(\begin{array}{c} q^{-n}, q^{-x}, -cq^{x-N} \\ q^{-N}, 0 \end{array}; q, q \right), \quad (5)$$

It satisfies the following orthogonal relation:

$$\begin{aligned} & \sum_{x=0}^N \frac{(-cq^{-N}, q^{-N}; q)_x}{(q, -cq; q)_x} \frac{(1 + cq^{2x-N})}{(1 + cq^{-N})} (-c)^{-x} q^{x(2N-x)} K_m(x; c, N | q) K_n(x; c, N | q) \\ &= (-c^{-1}; q)_N \frac{(q; q)_n}{(q^{-N}; q)_n} (cq^{-N})^n \delta_{mn}, \quad c > 0. \end{aligned} \quad (6)$$

In this paper, it will be helpful to use a rescaled version of the q -Krawtchouk polynomial, defined as

$$k(n, x; \rho; N, q) = c_k(n, \rho, N) K_n(x; q^{2\rho}, N; q^2) \quad (7)$$

with $c_k(n; \rho; N) = (-1)^n q^{-n\rho} q^{\frac{1}{2}n(N-1)}$. The greek letter *rho* is intentionally chosen to remind the reader of density.

2.2 Probabilistic Background

2.2.1 Definition of duality

Definition 1. *Two Markov processes $\xi(t)$ and $\eta(t)$ on states spaces \mathfrak{X} and \mathfrak{Y} , respectively, are dual with respect to the function $D(\xi, \eta)$ on $\mathfrak{X} \times \mathfrak{Y}$*

if

$$\mathbf{E}_\xi[D(\xi(t), \eta)] = \mathbf{E}_\eta[D(\xi, \eta(t))] \quad \text{for all } t \geq 0.$$

If, in addition, $\mathfrak{X} = \mathfrak{Y}$ and $\xi(t) = \eta(t)$ then $\xi(t)$ is self-dual with respect to the function $D(\xi, \eta)$.

If \mathfrak{X} and \mathfrak{Y} are discrete, an equivalent definition of duality is the intertwining relation

$$L_\xi D = D L_\eta^T,$$

where L_ξ is the generator of $\xi(t)$ viewed as a matrix with rows and columns indexed by \mathfrak{X} , L_η is the generator of $\eta(t)$ with rows and columns indexed by \mathfrak{Y} , and D is the duality function viewed as a matrix with rows indexed by \mathfrak{X} and columns indexed by \mathfrak{Y} . The superscript T indicates matrix transpose. Note that this is using the convention in probability, that a stochastic matrix has rows summing to 1.

Observe that all Markov processes are dual to each other with respect to constant functions, so part of the difficulty of duality is finding duality functions which have useful applications.

2.2.2 Definition of model

Recall from Borodin's paper [6] the weights for the dynamic stochastic six vertex model:

$$\begin{aligned}
\text{weight}(\odot) &= a_0^{\text{stoch}}(q, \lambda, w) = 1, & \text{weight}_{\lambda, w}(+) &= d_1^{\text{stoch}}(q, \lambda, w) = 1, \\
\text{weight}(|) &= a_1^{\text{stoch}}(q, \lambda, w) = \frac{1 - q^{\frac{1}{2}}w}{1 - q^{-\frac{1}{2}}w} \cdot \frac{q^{-1} - e^{2\pi i\lambda}}{1 - e^{2\pi i\lambda}}, \\
\text{weight}(\lrcorner) &= b_0^{\text{stoch}}(q, \lambda, w) = \frac{1 - q^{-1}}{1 - q^{-\frac{1}{2}}w} \cdot \frac{q^{\frac{1}{2}}w - e^{2\pi i\lambda}}{1 - e^{2\pi i\lambda}}, \\
\text{weight}(\ulcorner) &= c_1^{\text{stoch}}(q, \lambda, w) = \frac{\left(q^{\frac{1}{2}} - q^{-\frac{1}{2}}\right)w}{1 - q^{-\frac{1}{2}}w} \cdot \frac{\left(q^{\frac{1}{2}}w\right)^{-1} - e^{2\pi i\lambda}}{1 - e^{2\pi i\lambda}}, \\
\text{weight}(-) &= d_0^{\text{stoch}}(q, \lambda, w) = \frac{q^{-1} - q^{-\frac{1}{2}}w}{1 - q^{-\frac{1}{2}}w} \cdot \frac{q - e^{2\pi i\lambda}}{1 - e^{2\pi i\lambda}}.
\end{aligned} \tag{8}$$

Here, λ is the dynamic parameter and w is the spectral parameter. Briefly note that

$$\frac{d_0}{a_1} = \frac{q^{-1} - q^{-1/2}w}{1 - q^{1/2}w} \frac{q - e^{2\pi i\lambda}}{q^{-1} - e^{2\pi i\lambda}} = q^{-1} \cdot \frac{q - e^{2\pi i\lambda}}{q^{-1} - e^{2\pi i\lambda}} \in (q^{-1}, q)$$

In Aggarwal's paper [1], the fused weights are defined as follows. Define

$$\begin{aligned}
&\psi_{\Lambda, J, u, \lambda}(i_1, j_1; i_2, j_2) \\
&= \mathbf{1}_{i_1+j_1=i_2+j_2} q^{(j_2-i_1)J} \left(\frac{u}{s}\right)^{j_1} \frac{(q^{i_1-j_2+1}; q)_{j_2} (q^{j_2-J}; q)_{j_1} (suq^J; q)_{i_1-j_2} (s^2q^{i_1-j_2}; q)_{j_1}}{(su; q)_{i_1+j_1} (q; q)_{j_2} (q^{j_2-J}; q)_{j_1-j_2}} \\
&\times \frac{(us^{-1}\kappa^{-1}q^{-i_1}; q)_{j_2} (q^{1-i_1-J}u^{-1}s^{-1}\kappa^{-1}; q)_{j_1} (q\kappa^{-1}; q)_{j_1} (q^{j_2-2i_1+1}s^{-2}\kappa^{-1}; q)_{i_1-j_2}}{(q^{1-j_2}\kappa^{-1}; q)_{j_1} (q^{j_2-i_1-J+1}s^{-2}\kappa^{-1}; q)_{j_1} (q^{j_2-2i_1-J}s^{-2}\kappa^{-1}; q)_{j_2} (q^{2j_2-2i_1-J+1}s^{-2}\kappa^{-1}; q)_{i_1-j_2}} \\
&\times {}_{10}W_9(q^{-j_2}\kappa^{-1}; q^{-j_1}, q^{-j_2}, q^{j_1-J}\kappa^{-1}, q^{1-i_1}s^{-2}\kappa^{-1}, su^{-1}q^{i_1-j_2+1}, usq^{i_1-j_2+J}, q^{-i_1}\kappa^{-1}; q, q)
\end{aligned} \tag{9}$$

with

$$q = e^{-4\pi i \eta}; \quad s = e^{2\pi i \eta \Lambda}; \quad \varkappa = e^{2\pi i \lambda}; \quad \tilde{\varkappa} = q^{J-2j_1} \varkappa; \quad \kappa = \tilde{\varkappa}^{-1},$$

and where

$${}_{r+1}W_r(a_1; a_4, a_5, \dots, a_{r+1}; q, z) = \sum_{k=0}^{\infty} \frac{z^k (a_1; q)_k}{(q; q)_k} \frac{1 - a_1 q^{2k}}{1 - a_1} \prod_{j=4}^{r+1} \frac{(a_j; q)_k}{(qa_1/a_j; q)_k} \quad (10)$$

denotes the very-well poised basic hypergeometric series. These weights sum to 1 in the sense that

$$\sum_{i_2, j_2} \psi(i_1, j_1; i_2, j_2) = 1.$$

In (9), J is the horizontal spin and Λ is the vertical spin. As before λ is the dynamic parameter, η is the asymmetry parameter and u is the spectral parameter.

From these weights, a discrete–time Markov chain can be constructed as a vertex model (see [1] for a detailed definition). In this paper, we consider the model on finite sites with closed boundaries (i.e. no horizontal incoming arrows through the left boundary and no horizontal outgoing arrows through the right boundary), or on the infinite line.

Now we briefly describe the model defined on $[1, N] \times \mathbf{Z}^+$ (See Figure 1). For each vertex $(x, y) \in [1, N] \times \mathbf{Z}^+$, there is an arrow configuration, which is a quadruple

$$(i_1, j_1; i_2, j_2) = (i_1, j_1; i_2, j_2)_{(x,y)} = (i_1(x, y), j_1(x, y); i_2(x, y), j_2(x, y))$$

of non-negative integers, where j_1 denotes the number of horizontal incoming arrows at (x, y) ; i_1 denotes the number of vertical incoming arrows; j_2 denotes the number of horizontal outgoing arrows; and i_2

denotes the number of vertical outgoing arrows. At any vertex, the total number of incoming arrows is equal to the total number of outgoing arrows (i.e. $i_1 + j_1 = i_2 + j_2$). Let J_k be the horizontal spin, and I_k be the vertical spin for strip k , which means $0 \leq j_1(\cdot, y), j_2(\cdot, y) \leq 2J_y$, and $0 \leq i_1(x, \cdot), i_2(x, \cdot) \leq 2I_x$. Each vertex (x, y) is associated with a dynamical parameter $\lambda(x, y)$ according to the following rules:

$$\begin{aligned}\lambda(x+1, y) &= \lambda(x, y) - 4\eta(i_1(x+1, y) - I_{x+1}), \\ \lambda(x, y+1) &= \lambda(x, y) + 4\eta(j_1(x, y) - J_y).\end{aligned}$$

Given the empty left boundary (i.e. $j_1(1, \cdot) = 0$) and arbitrary bottom boundary (i.e. $i_1(\cdot, 1)$), we now update the configuration from left to right and from bottom to top with transition probability at site (x, y) defined as

$$\mathcal{P}_{x,y}(i_2, j_2 | i_1, j_1) = \psi_{I_x, J_y, u_x, \lambda(x,y)}(i_1, j_1; i_2, j_2). \quad (11)$$

Similarly to the stochastic six vertex model [8], there is an equivalent way to define the model as an interacting particle system: consider the \mathcal{P} -distributed random configuration and cut it by horizontal lines $y = t + \frac{1}{2}$, $t = 0, 1, 2, \dots$, then $\eta_x(t) = i_1(x, t + \frac{1}{2})$ is a Markov chain. Since the horizontal arrows are to the right, which means the particles only jump to the right, we say that the model has asymmetry to the right.

2.3 Algebraic Background

2.3.1 Quantum groups as quasi-triangular quasi-Hopf *-algebras

The quantum group $U_q(sl_2)$ is a unital algebra generated by E_{\pm} and q^H with relations:

$$[H, E_{\pm}] = \pm 2E_{\pm}, \quad [E_+, E_-] = \frac{q^H - q^{-H}}{q - q^{-1}}.$$






Figure 1: This shows one-time update of the dynamic stochastic higher spin vertex model. The update occurs sequentially from left to right. The dashed lines indicate that the update has not yet occurred, and each solid line shows the number of arrows.

The co-unit is $\epsilon(q^H) = 1, \epsilon(E_{\pm}) = 1$ The choice of co-product in this paper will be

$$\Delta(H) = H \otimes \text{id} + \text{id} \otimes H, \quad \Delta(E_{\pm}) = E_{\pm} \otimes q^{H/2} + q^{-H/2} \otimes E_{\pm}.$$

The above definitions define a bialgebra structure on $U_q(sl_2)$. There is also an antipode which defines a Hopf algebra structure, but we will not (explicitly) use the antipode.

Recall Drinfeld's universal R -matrix:

$$R_{12}^D = q^{\frac{1}{2}H \otimes H} \sum_{i=0}^{\infty} (q - q^{-1})^i \frac{q^{-\frac{i(i+1)}{2}}}{[i]!} q^{\frac{i}{2}H} E_+^i \otimes q^{-\frac{i}{2}H} E_-^i.$$

This R -matrix satisfies the property that

$$R^D \Delta(X) = \Delta'(X) R^D$$

for any $X \in U_q(sl_2)$, where $\Delta' = P \circ \Delta$ is the opposite comultiplication defined by $P(X \otimes Y) = Y \otimes X$. This property makes $U_q(sl_2)$ an *almost-cocommutative* Hopf algebra. The R -matrix additionally satisfies

$$R_{12}^D R_{21}^D = 1, \quad (\epsilon \otimes \epsilon) R_{12}^D = 1 \quad R_{12}^D (\Delta \otimes \text{id})(R^D) = R_{23}^D (\text{id} \otimes \Delta)(R^D),$$

which makes it a *coboundary* Hopf algebra. Additionally,

$$(\Delta \otimes \text{id})(R^D) = R_{13}^D R_{23}^D, \quad (\text{id} \otimes \Delta)(R^D) = R_{13}^D R_{12}^D,$$

making it a *quasi-triangular* Hopf algebra. In any quasi-triangular Hopf algebra, the R -matrix satisfies the quantum Yang-Baxter equation [23]

$$R_{12}^D R_{13}^D R_{23}^D = R_{23}^D R_{13}^D R_{12}^D.$$

In fact, there is a spectral-dependent version of the R -matrix, denoted

$R^D(z)$, satisfying the Yang–Baxter Equation

$$R_{12}^D(u)R_{13}^D(uv)R_{23}^D(v) = R_{23}^D(v)R_{13}^D(uv)R_{12}^D(u). \quad (12)$$

The R –matrix can be constructed from the quantized affine Lie algebra $U_q(\widehat{sl_2})$, although that construction is not needed for this paper.

There is a generalization of Hopf algebras to quasi–Hopf algebras, which can be constructed with a twisting procedure. Suppose there is a twister satisfying the following shifted cocycle condition [5]:

$$F_{12}(\lambda)(\Delta \otimes \text{id})F(\lambda) = F_{23}(\lambda - 2\eta H_1)(\text{id} \otimes \Delta)F(\lambda); \quad (13)$$

then the twisted R –matrix

$$R(z, \lambda) = F_{21}(\lambda) (R^D(z))^{-1} F_{12}^{-1}(\lambda) \quad (14)$$

satisfies the dynamical Yang–Baxter Equation [15]

$$\begin{aligned} R_{12}(u, \lambda - 2\eta H_3)R_{13}(uv, \lambda)R_{23}(v, \lambda - 2\eta H_1) \\ = R_{23}(v, \lambda)R_{13}(uv, \lambda - 2\eta H_2)R_{12}(u, \lambda), \end{aligned} \quad (15)$$

where η is defined in a way such that $q = e^{2\pi i\eta}$.

Now set

$$\Delta_\lambda(X) = F(\lambda)\Delta(X)F^{-1}(\lambda),$$

which is called the twisted co-product. Then,

$$R(z, \lambda)\Delta_\lambda(X) = \Delta'_\lambda(X)R(z, \lambda)$$

for any $X \in U_q(sl_2)$. Additionally,

$$(\Delta_\lambda \otimes \text{id})(R(z, \lambda)) = \Phi_{312}R_{13}(z, \lambda)\Phi_{132}^{-1}R_{23}(z, \lambda)\Phi_{123},$$

$$(\text{id} \otimes \Delta_\lambda)(R(z, \lambda)) = \Phi_{231}^{-1}R_{13}(z, \lambda)\Phi_{213}R_{12}(z, \lambda), \Phi_{123}^{-1}$$

where $\Phi = (F_{23}(\lambda)(\text{id} \otimes \Delta)F(\lambda))(F_{12}(\lambda)(\Delta \otimes \text{id})F(\lambda))^{-1}$. These properties makes $U_q(\mathfrak{sl}_2)$ a quasi-Hopf algebra. For more details about quasi-Hopf algebra, we refer to [21].

There is also a $*$ -Hopf algebra structure on $U_q(\mathfrak{sl}_2)$ for nonzero real values of q . The involution is given by the explicit formula

$$E_+^* = E_-, \quad E_-^* = E_+, \quad (q^H)^* = q^H.$$

The $*$ -algebra structure can also be defined using the antipode, in addition to this explicit formula.

Finally, recall that for $J \in \{1/2, 1, 3/2, 2, 5/2, \dots\}$, the $(2J+1)$ -dimensional representations are given by

$$\begin{aligned} \rho^{(J)}(H)|J+m\rangle &= 2m|J+m\rangle & -J \leq m \leq J \\ \rho^{(J)}(E_{\pm})|J+m\rangle &= \sqrt{[J \mp m][J \pm m + 1]}|J+m \pm 1\rangle, \end{aligned}$$

Denote this representation by V^{2J} . We interpret $J+m$ to be the number of particles and $J-m$ to be the number of holes at each site.

3 Main Results

3.1 Probabilistic Results

The main theorems of this paper are purely probabilistic results, which can be stated without algebra. The first is a Markov duality result, and the second is an application to Tracy–Widom asymptotics. Below, the weights for the model are defined using different notations from those described in section 2.2.2. More specifically, define the weights $S(i_1, j_1, i_2, j_2)$ to be equal to $\psi(i_1, j_1, i_2, j_2)$ with q replaced by q^2 and u replaced by $\frac{1}{zq^2}$.

3.1.1 Markov Duality

Let J_i be the vertical spin at site i , μ and ξ be the vertex models with transition matrices given by (26) and (27), respectively. Then we have the following duality results (see Corollary 3).

Theorem 1. *The dynamic stochastic higher spin vertex model μ , with asymmetry to the left, is dual to a (non–dynamic) stochastic higher spin vertex model ξ with asymmetry to the right, with respect to the duality functions:*

$$\begin{aligned} \mathcal{D}_c(\mu, \xi) &= \prod_{i=1}^N \left(e^{-2\pi i \lambda} q^{2(-c+N_{[1,i-1]}(\mu+\xi-2J)+\mu_i-2J_i)}; q^2 \right)_{2J_i-\mu_i} \\ &\times {}_3\varphi_2 \left(\begin{matrix} q^{-2(2J_i-\mu_i)}, q^{-2\xi_i}, e^{-2\pi i \lambda} q^{2(2N_{[1,i-1]}(\mu-J)+\mu_i-2J_i)} \\ q^{-4J_i}, e^{-2\pi i \lambda} q^{2(-c+N_{[1,i-1]}(\mu+\xi-2J)+\mu_i-2J_i)} \end{matrix} \middle| q^2, q^2 \right) \\ &\times q^{-4J_i N_{[1,i-1]}(\xi) - 2\xi_i N_{[1,i]}(\mu)} \end{aligned}$$

where c is a constant and $N_{[1,i]}(\eta) = \sum_{k=1}^i \eta_k$.

Taking c to $\pm\infty$ so that $q^c \rightarrow 0$ yields an orthogonal duality:

Corollary 1. *The dynamic stochastic higher spin vertex model μ , with asymmetry to the left, is dual to a (non–dynamic) stochastic higher spin vertex model ξ with asymmetry to the right, with respect to the duality functions:*

$$\begin{aligned} \mathcal{D}_{ort}(\mu, \xi) &= \prod_{i=1}^N {}_3\varphi_2 \left(\begin{matrix} q^{2(2J_i-\mu_i)}, q^{2\xi_i}, e^{-2\pi i \lambda} q^{2(2N_{[1,i-1]}(\mu-J)+\mu_i)} \\ q^{4J_i}, 0 \end{matrix} \middle| q^{-2}, q^{-2} \right) \\ &\times q^{-4J_i N_{[1,i]}(\xi) - 2\xi_i N_{[1,i-1]}(\mu)} \quad (16) \end{aligned}$$

Moreover, when $-e^{-2\pi i\lambda} > 0$, there are orthogonal relations

$$\sum_{\xi} \mathcal{D}_{ort}(\mu, \xi) \mathcal{D}_{ort}(\mu', \xi) w(\xi; q, J) = \frac{\delta_{\mu, \mu'}}{W(\mu; q, J, \lambda)}, \quad (17)$$

$$\sum_{\mu} \mathcal{D}_{ort}(\mu, \xi) \mathcal{D}_{ort}(\mu, \xi') W(\mu; q, J, \lambda) = \frac{\delta_{\xi, \xi'}}{w(\xi; q, J)}, \quad (18)$$

with

$$w(\xi; q, J) = q^{\frac{1}{2} \sum_{i=1}^N (2J_i \xi_i - 2\xi_i N_{[1, i]}(2J))} \prod_{i=1}^N q^{-\xi_i (\xi_i - 2J_i)} \left[\begin{matrix} 2J_i \\ \xi_i \end{matrix} \right]_{q^{-2}}$$

and

$$W(\mu; q, J, \lambda) = \prod_{i=1}^N \mathcal{W} \left(2J_i - \mu_i; q^{-1}, 2J_i, \log_{q^{-2}} \left(-e^{-2\pi i\lambda} q^{4N_{[1, i-1]}(\mu - J)} \right) \right),$$

$$\mathcal{W}(x; q, N, \rho) = \frac{1 + q^{4x+2\rho-2N}}{1 + q^{2\rho-2N}} \frac{(-q^{2\rho-2N}; q^2)_x}{(-q^{2\rho+2}; q^2)_x} \frac{q^{-x(2\rho+1+x-2N)}}{(-q^{-2\rho}; q^2)_N} \left[\begin{matrix} N \\ x \end{matrix} \right]_{q^2}. \quad (19)$$

Remark 1. $\mathcal{D}_{ort}(\mu, \xi)$ is equal to K_L in Theorem 3.14 of [19], where K_L is proven to be an orthogonal duality between dyanmic ASEP and ASEP, the orthogonal relation is given by Corollary 3.15 in [19]. The measure w and W are reversible measures for ASEP and dynamic ASEP, respectively. If we let $J_i = J$ for any i , then up to a constant,

$$\lim_{J \rightarrow \infty} \mathcal{D}_{ort}(\mu, \xi) = \prod_{i=1}^N q^{2\xi_i N_{[1, i]}(\mu)},$$

which is a self duality function for q -TAZRP (totally asymmetric zero range process). Note that this degeneration confirms the correct directions of the asymmetry.

In the limit to the non–dynamic model as $\lambda = -\mathbf{i}\infty$ in $\mathcal{T}^{stoch}(z, \lambda)$, we obtain a new duality function:

Corollary 2. *The stochastic higher spin vertex model, with asymmetry to the left, is dual to a stochastic higher spin vertex model with asymmetry to the right, with respect to the duality function*

$$\begin{aligned} & \mathcal{D}_{new}(\mu, \xi) \\ &= \prod_{i=1}^N \frac{(q^{2(C_0+N_{[1,i-1]}(\mu+\xi-2J)+\mu_i-2J_i+1)}; q^2)_{2J_i-\mu_i}}{(q^{-2(C_0+N_{[1,i-1]}(\mu+\xi-2J)+\xi_i)}; q^2)_{2J_i-\mu_i}} \\ & \times {}_2\varphi_1 \left(\begin{matrix} q^{-2(2J_i-\mu_i)}, q^{-2(2J_i-\xi_i)} \\ q^{-4J_i} \end{matrix} \middle| q^2, q^{-2(C_0+N_{[1,i]}(\mu+\xi-2J))} \right) q^{-4J_i N_{[1,i-1]}(\xi) - 2\xi_i N_{[1,i]}(\mu)} \end{aligned}$$

where C_0 is a fixed constant.

Remark 2. Take $C_0 \rightarrow \infty$ in \mathcal{D}_{new} we get a triangular duality

$$\mathcal{D}_{tr}(\mu, \xi) = \prod_{i=1}^N 1_{\{\mu_i \geq \xi_i\}} \frac{[\mu_i]![2J_i - \xi_i]!}{[\mu_i - \xi_i]!} q^{-4J_i N_{[1,i-1]}(\xi) + \mu_i \xi_i + 2\mu_i N_{[1,i-1]}(\xi) - 2J_i \xi_i},$$

which actually is equal to the duality in [26]. To see this apply the “charge–parity symmetry” to ξ in Theorem 2.5(a).

A simple way to check the duality result is to look at “two–site” dualities or degenerate cases. In the next examples, we use that for the “two–site” dualities the transition matrices are 4×4 matrices, and duality matrices are calculated up to a constant in the total number of particles. Below, $\mathcal{T}^{stoch}(z, \lambda)$ is the transition matrix for process η and $\mathcal{T}_{rev}^{stoch}(z, -\mathbf{i}\infty)$ is the

transition matrix for the non dynamic process ξ .

$$\mathcal{T}_{\text{rev}}^{\text{stoch}}(z, -\mathbf{i}\infty) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \frac{q^2-1}{(zq^2-1)} & \frac{q^2(z-1)}{zq^2-1} & 0 \\ 0 & \frac{z-1}{zq^2-1} & \frac{z(q^2-1)}{zq^2-1} & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathcal{T}^{\text{stoch}}(z, \lambda) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \frac{(q^2-1)(1-e^{-2\pi i\lambda}z)}{(zq^2-1)(1-e^{-2\pi i\lambda})} & \frac{(q^2-e^{-2\pi i\lambda})(z-1)}{(zq^2-1)(1-e^{-2\pi i\lambda})} & 0 \\ 0 & \frac{(z-1)(1-e^{-2\pi i\lambda}q^2)}{(zq^2-1)(1-e^{-2\pi i\lambda})} & \frac{(q^2-1)(z-e^{-2\pi i\lambda})}{(zq^2-1)(1-e^{-2\pi i\lambda})} & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Comparing these matrices with Borodin's weights in (8), the notation matches with $q^{\frac{1}{2}}w$ replaced by z^{-1} and q being replaced with q^2 (see the 6 for more details).

Example 1. The matrix \mathcal{D}_c equals

$$\begin{bmatrix} (e^{-2\pi i\lambda}q^{-2c-2}-1)(e^{-2\pi i\lambda}q^{-2c-4}-1) & -(e^{-2\pi i\lambda}(e^{-2\pi i\lambda}-q^{2c+2})(q^{2c}-1))q^{-4c-6} & -(e^{-2\pi i\lambda}(e^{-2\pi i\lambda}-q^{2c+2})(q^{2c}-1))q^{-4c-6} & (e^{-4\pi i\lambda}(q^{2c}-q^2)(q^{2c}-1))q^{-4c-8} \\ 1-e^{-2\pi i\lambda}q^{-2c-2} & -(e^{-2\pi i\lambda}q^{-2c-2}-1)q^{-2} & (e^{-2\pi i\lambda}(q^{2c}-1))q^{-2c-4} & (e^{-2\pi i\lambda}(q^{2c}-1))q^{-2c-6} \\ 1-e^{-2\pi i\lambda}q^{-2c-2} & (e^{-2\pi i\lambda}(q^{2c+2}-1))q^{-2c-4} & -(q^{-2c}e^{-2\pi i\lambda}-1)q^{-4} & (e^{-2\pi i\lambda}(q^{2c}-1))q^{-2c-6} \\ 1 & q^{-4} & q^{-4} & q^{-8} \end{bmatrix}$$

and satisfies $\mathcal{T}^{\text{stoch}}(z, \lambda)\mathcal{D}_c = \mathcal{D}_c(\mathcal{T}_{\text{rev}}^{\text{stoch}}(z, -\mathbf{i}\infty))^*$.

Example 2. The matrix \mathcal{D}_{new} equals

$$\mathcal{D}_{\text{new}} = \begin{bmatrix} ((q^{2C_0}-q^2)(q^{2C_0}-q^4))q^{-6} & (q^{2C_0}(q^{2C_0}-q^2))q^{-6} & (q^{2C_0}(q^{2C_0}-q^2))q^{-6} & q^{4C_0-6} \\ 1-q^{2C_0-2} & -(q^{2C_0}-q^2)q^{-4} & -q^{2C_0-4} & -q^{2C_0-6} \\ 1-q^{2C_0-2} & -q^{2C_0-4} & -(q^{2C_0}-1)q^{-4} & -q^{2C_0-6} \\ 1 & q^{-4} & q^{-4} & q^{-8} \end{bmatrix}$$

and satisfies $\mathcal{T}^{\text{stoch}}(z, -\mathbf{i}\infty)\mathcal{D}_{\text{new}} = \mathcal{D}_{\text{new}}(\mathcal{T}_{\text{rev}}^{\text{stoch}}(z, -\mathbf{i}\infty))^*$.

Example 3. The matrix \mathcal{D}_{tr} equals

$$\mathcal{D}_{\text{tr}} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ q^{-1/2} & 1 & 0 & 0 \\ q^{-1/2} & 0 & q^{-2} & 0 \\ q & q^{3/2} & q^{3/2} & q^4 \end{bmatrix}$$

and satisfies $\mathcal{T}^{\text{stoch}}(z, -\mathbf{i}\infty)\mathcal{D}_{tr} = \mathcal{D}_{tr}(\mathcal{T}_{\text{rev}}^{\text{stoch}}(z, -\mathbf{i}\infty))^*$.

3.1.2 Asymptotics

As an application of the duality, we can prove that the large–time fluctuations of the dynamic stochastic six–vertex model are governed by the Tracy–Widom distribution:

Theorem 2. *Let $\mu(t)$ denote the dynamic stochastic six vertex model with asymmetry to the right, and started with step initial conditions (i.e. all lattice sites to the right of the origin contains an up–arrow). Suppose $q < 1, z < 1$ and that $\nu \in (z, z^{-1})$.*

Let L be a large parameter, and let the spatial variable y and the time variable t depend on L in such a way that $y = \lfloor \nu L \rfloor$ and $t = L$. Let the dynamic parameter λ depend on L in such a way that λ is purely imaginary and $\liminf_{L \rightarrow \infty} \lambda/(\mathbf{i}L) \leq 0$. Then for any real value of s ,

$$\lim_{L \rightarrow \infty} \mathbf{P} \left(\frac{N_y(\mu(t)) - m_\nu L}{\sigma_\nu L^{1/3}} \leq s \right) \rightarrow F_2(s)$$

where $F_2(s)$ is cumulative distribution function of the Tracy–Widom distribution, and m_L, σ_L equal

$$m_\nu := \frac{(\sqrt{\nu} - \sqrt{z})^2}{1 - z}, \quad \sigma_\nu := \frac{z^{1/2} \nu^{-1/6}}{1 - z} ((1 - \sqrt{\nu z})(\sqrt{\nu z^{-1}} - 1))^{2/3}$$

Remark 3. Somewhat surprisingly, the asymptotic outputs m_L, σ_L and $F_2(s)$ do not depend on the dynamic parameter λ . A similar phenomenon occurs for the usual stochastic six vertex model [8], where the parameters do not depend on the asymmetry parameter q . Computer simulations in the appendix also support the lack of dependence on λ . Following the proof of the theorem (section 4.1), additional intuition will be provided, based on the form of the duality function.

¹For example, this holds if $\lambda/\mathbf{i} \leq 0$ for all values of L .

Remark 4. In more standard probability notation, the usual convention is that asymmetry parameter q and spectral parameter z take values in $(0, 1)$. Since the quantum group $U_q(sl_2)$ is invariant under the replacements $q \leftrightarrow q^{-1}$, in an algebraic setting the notational convention is different.

3.2 Algebraic Results

The proof of this probabilistic result uses algebraic methods. These methods are very general, therefore we summarize them here. Note that the main concept of this proof, that “duality functions can be constructed from vertex models”, was communicated to us by Michael Wheeler and William Mead [36, 37, 33].

Consider the following setup. There is a matrix $R(z, \lambda)$, acting on a tensor power $V \otimes W$, which depends on a quantization parameter q , a spectral parameter z and the dynamical parameter λ and satisfies the dynamical Yang–Baxter Equation (15). More generally, consider the tensor product $V_0 \otimes V_1 \otimes \cdots \otimes V_N$, with V_0 called the “auxiliary space” and $V_1 \otimes \cdots \otimes V_N$ called “the bulk space.” Let $2J_i + 1$ be the dimension of V_i .

Let P be the permutation map on two sites, defined by $P : V \otimes W \rightarrow W \otimes V$ with $P(v \otimes w) = w \otimes v$. Let the $\check{\cdot}$ notation refer to

$$\check{R}_{i,i+1} = P_{i,i+1} R_{i,i+1} = R_{i+1,i} P_{i,i+1}.$$

The transfer matrix $\mathcal{T}(\lambda)$ is defined by

$$\mathcal{T}(z, \lambda) = \check{R}_{N-1,N} (z, \lambda - 2\eta N_{[1,N-2]}(H)) \check{R}_{N-2,N-1} (z, \lambda - 2\eta N_{[1,N-3]}(H)) \cdots \check{R}_{1,2} (z, \lambda),$$

where $N_{[a,b]}(H) = H_a + H_{a+1} \cdots + H_b$. Define

$$\mathcal{B}(u, z, \lambda) = R_{0,N}(u, \lambda - 2\eta N_{[1,N-1]}(H)) R_{0,N-1}(uz, \lambda - 2\eta N_{[1,N-2]}(H)) \cdots R_{0,1}(uz^{N-1}, \lambda)$$

and for each $0 \leq m, l \leq \dim V_0$

$$\langle \mu | B_{ml}(u, z, \lambda) | \xi \rangle = \langle m, \mu | \mathcal{B}(u, z, \lambda) | l, \xi \rangle.$$

Theorem 3. Define $\check{\mathfrak{R}}$ so that

$$\check{\mathfrak{R}}(z, \lambda) = (F_{12}(\lambda)(\Pi^{-1})^{\otimes 2}(F_{21}(\lambda))^* (R(z, \lambda)P)^* (F_{21}(\lambda)^*)^{-1} \Pi^{\otimes 2} F_{12}^{-1}(\lambda))^*.$$

Then there is the intertwining relationship

$$\check{R}_{i,i+1}(z, \lambda - 2\eta N_{[1,i-1]}(H)) B_{m,l}(u, z, \lambda) = B_{m,l}(u, z, \lambda) (\check{\mathfrak{R}}_{i,i+1}(z, \lambda(l)))^*, \quad (20)$$

where $\lambda(l) = \lambda - 2\eta N_{[1,i-1]}(H) - 2\eta H_0(l)$ with $H_0(l)$ denoting the constant $2(l - J_0)$.

If a few more properties hold, then there is an intertwining relationship between stochastic matrices. First, suppose that $R(z, \lambda)$ satisfies the transposition formula

$$(F_{21}(\lambda)^*)^{-1} \Pi^{\otimes 2} F_{12}(\lambda)^{-1} P R(z, \lambda) = P R(z, \lambda)^* (F_{21}(\lambda)^*)^{-1} \Pi^{\otimes 2} F_{12}(\lambda)^{-1}, \quad (21)$$

where Π is a permutation matrix. Second, suppose there is a ground state symmetry

$$G_{i,i+1}^{-1}(\lambda) \check{R}_{i,i+1}(z, \lambda) G_{i,i+1}(\lambda) = \check{S}_{i,i+1}(z, \lambda) \quad (22)$$

such that $S_{i,i+1}(z, \lambda)$ is a stochastic matrix, meaning that each row sums to 1 and all entries are non-negative, for some values of z, λ . Further, we

assume there is a diagonal matrix C such that

$$\Pi G(-\mathbf{i}\infty)\Pi = CPG(-\mathbf{i}\infty)P. \quad (23)$$

Last, suppose that this same $S(z, \lambda)$ satisfies the inversion formula

$$\Pi^{\otimes 2}S(z, \lambda)\Pi^{\otimes 2} = \mathfrak{S}(z^{-1}, \lambda), \quad (24)$$

where $\mathfrak{S}(z, \lambda)$ is defined by substituting q with q^{-1} in $S(z, \lambda)$.

Note that equations (22) and (24) can be combined into

$$(\check{R}(z, \lambda))^* = G^{-1}(\lambda)\Pi^{\otimes 2}(\check{\mathfrak{S}}(z^{-1}, \lambda))^* \Pi^{\otimes 2}G(\lambda). \quad (25)$$

Define the stochastic transfer matrix by

$$\begin{aligned} \mathcal{T}^{\text{stoch}}(z, \lambda) = \check{S}_{N-1, N}(z, \lambda - 2\eta N_{[1, N-2]}(H)) \check{S}_{N-2, N-1}(z, \lambda - 2\eta N_{[1, N-3]}(H)) \\ \cdots \check{S}_{1, 2}(z, \lambda), \end{aligned} \quad (26)$$

and

$$\begin{aligned} \mathcal{T}_{\text{rev}}^{\text{stoch}}(z, -\mathbf{i}\infty) \\ = (\mathfrak{S}P)_{1, 2}(z^{-1}, -\mathbf{i}\infty) \cdots (\mathfrak{S}P)_{N-2, N-1}(z^{-1}, -\mathbf{i}\infty) (\mathfrak{S}P)_{N-1, N}(z^{-1}, -\mathbf{i}\infty). \end{aligned} \quad (27)$$

Remark 5. Notice that in the definition of $\mathcal{T}^{\text{stoch}}(z, \lambda)$, each \check{S} has row sums to 1, thus $\mathcal{T}^{\text{stoch}}(z, \lambda)$ could be viewed as a transfer matrix for the dynamic stochastic vertex model with drift to the left, with the dynamic parameter of site N being equal to $\lambda + 2\eta N_{[1, N]}(H)$.

Also Notice that

$$\begin{aligned} \lim_{H_0(l) \rightarrow \infty} (\check{\mathfrak{R}}(z, \lambda(l)))^* &= (\Pi^{-1})^{\otimes 2} (R(z, -\mathbf{i}\infty)P)^* \Pi^{\otimes 2} \\ &= (\Pi^{-1})^{\otimes 2} \left((R^D(z))^{-1} P \right)^* \Pi^{\otimes 2}, \end{aligned}$$

where in the last equality we used the fact that $PR^D(z)PR^D(z^{-1}) = \text{Id}$ (see (11) in [31]). This means it reduces to the non-dynamic case in this limit.

In the theorem below, we will take the quantity $H_0(l)$ to ∞ , as in such limit, the twister terms in $\check{\mathfrak{R}}(z, \lambda)$ become identities.

Theorem 4. *With all of the above assumptions (specifically equations (15), (21)–(24), there is the “two-site” duality relation*

$$\begin{aligned} &\check{S}_{i,i+1}(z, \lambda - 2\eta N_{[1,i-1]}(H)) \\ &\quad \times G_{i,i+1}^{-1}(\lambda - 2\eta N_{[1,i-1]}(H)) \left(\lim_{H_0(l) \rightarrow \infty} B_{\mathbf{m},1}(u, z, \lambda) \right) G_{i,i+1}^{-1}(-\mathbf{i}\infty) \\ &= G_{i,i+1}^{-1}(\lambda - 2\eta N_{[1,i-1]}(H)) \left(\lim_{H_0(l) \rightarrow \infty} B_{\mathbf{m},1}(u, z, \lambda) \right) G_{i,i+1}^{-1}(-\mathbf{i}\infty) \\ &\quad \times (\mathfrak{S}(z^{-1}, -\mathbf{i}\infty)P)_{i,i+1}^*. \end{aligned}$$

Corollary 3. *There is the duality relation*

$$\mathcal{T}^{\text{stoch}}(z, \lambda)\mathcal{D} = \mathcal{D}(\mathcal{T}_{\text{rev}}^{\text{stoch}}(z, -\mathbf{i}\infty))^*,$$

where

$$\mathcal{D} = \text{const} \cdot \mathcal{G}^{-1}(\lambda) \left(\lim_{H_0(l) \rightarrow \infty} B_{\mathbf{m},1}(u, z, \lambda) \right) \mathcal{G}^{-1}(-\mathbf{i}\infty)$$

with

$$\langle \mu | \mathcal{G}(\lambda) | \mu \rangle = \prod_{i=1}^N \frac{q^{2J_i N_{[1,i-1]}(\mu) + J_i \mu_i}}{\sqrt{[\mu_i]! [2J_i - \mu_i]!} (e^{-2\pi i \lambda} q^{4(1+N_{[1,i]}(\mu-J))}; q^2)_{2J_i - \mu_i}}.$$

On the right-hand-side of the duality relation, the $-i\infty$ means that it is a non-dynamic stochastic vertex model, and the subscript refers to space reversal. The term “const” is a normalization constant so that the duality function is nontrivial in the limit.

To relate these algebraic ideas to the dynamic stochastic higher-spin vertex models, it needs to be shown that this specific model fits in the general framework. The next proposition states exactly that:

Proposition 1. *The dynamic stochastic higher-spin vertex model satisfies the equations (15)–(24).*

The next step is to computationally verify that the duality functional \mathcal{D} has the expression stated in Theorem 1.

Proposition 2. *The duality functional \mathcal{D} has the expression stated in Theorem 1.*

The final step is to verify that the fusion procedure of Aggarwal [1] produces the same stochastic models as the use of the universal R -matrix and twister in his paper.

Proposition 3. *The stochastic weights (9) from [1] are the same weights as $S(z, \lambda)$, up to a change of variable.*

An explicit example of the matching of the weights in the spin 1 case can be found in the Appendix, section 6.

4 Proofs

4.1 Proof of Theorem 2 (assuming Theorem 1)

The second probabilistic result, Theorem 2, follows from Theorem 1. This will be proved first. Before continuing with the proof, we outline the idea of the proof. The first step is to bound the difference between the duality function for the dynamic model and the duality function for the non–dynamic model. The second step is to use this bound (on the duality functions) to then bound the difference between the height function for the dynamic model and the height function for the non–dynamic model. It is this step that requires orthogonality of the duality functions. Finally, using the previously known result that the (rescaled) height function for the non–dynamic model converges to the Tracy–Widom distribution, this proves convergence that the (rescaled) height function of the dynamic model also converges to the Tracy–Widom distribution. The second step above holds more generally for models with orthogonal polynomial duality. It is based on the following proposition, which we state in its own subsection because of its generality. Furthermore, the proposition will inform us of the necessary bounds in the context of the dynamic stochastic six vertex model.

4.2 A general result using orthogonality

Consider the following setup and notation. Let $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in \mathbb{N}}, \mathbf{P})$ be a filtered probability space. Let X_t^λ be a stochastic process with values in a countable state space \mathcal{S} , so in other words X_t is a \mathcal{S} –valued random variable on $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in \mathbb{N}}, \mathbf{P})$. Let $Q_\lambda(t)$ denote the probability measure on \mathcal{S} which is the pushforward of P under X_t^λ . Let $Q_\lambda(t, x)$ denote the weight of $x \in \mathcal{S}$ under $Q_\lambda(t)$.

Suppose that we have a family of measures $\{W_\lambda\}_{\lambda \in \mathbf{R}_{\geq 0}}$ on the state space \mathcal{S} , and let \mathcal{H}_λ denote the Hilbert space

$$\mathcal{H}_\lambda = L^2(\mathcal{S}, W_\lambda).$$

Let $\langle f, g \rangle_\lambda$ denote the usual inner product with respect to W_λ , i.e.

$$\langle f, g \rangle_\lambda = \sum_{x \in \mathcal{S}} f(x)g(x)W_\lambda(x)$$

Suppose that \mathcal{H}_λ has an orthonormal basis with respect to W_λ , denoted by $\{D_\lambda^S\}$ where S indexes the basis.

Let $h(L)$ be an element of \mathcal{H}_λ (for all values of λ) which depends on a large parameter L . Let $a_\lambda^S(L)$ be defined by

$$h(L) = \sum_S a_\lambda^S(L) D_\lambda^S,$$

or equivalently

$$a_\lambda^S(L) = \langle h(L), D_\lambda^S \rangle_\lambda = \sum_{x \in \mathcal{S}} h(L)[x] D_\lambda^S(x) W_\lambda(x).$$

Now suppose that t depends on L as well, and for simplicity set $t = L$. The proposition is the following, which we state in large generality to include a large variety of processes:

Proposition 4. *Suppose that we have the upper bound for all S and L :*

$$\left(\sum_{y \in \mathcal{S}} h(L)[y] \cdot |D_\lambda^S(y)W_\lambda(y) - D_0^S(y)W_0(y)| \right) \leq C_1 L^2 \quad (28)$$

and

$$\sum_{y \in \mathcal{S}} h(L)[y] D_\lambda^S(y) W_\lambda(y) \leq C_2 L^2. \quad (29)$$

Further suppose that we have the upper bounds for all S :

$$\sum_{x \in S} |D_0^S[x] Q_0(L, x)| \leq M_1(L, S) \quad (30)$$

$$\left(\sum_{x \in S} D_\lambda^S[x] (Q_\lambda(L, x) - Q_0(L, x)) \right) \leq M_3(L, S) \quad (31)$$

where for $j = 1, 3$, the terms $M_j(L, S)$ satisfy

$$\sum_S L^2 M_j(L, S) < \infty, \quad \lim_{L \rightarrow \infty} L^2 M_j(L, S) = 0$$

and additionally

$$\lim_{L \rightarrow \infty} \sum_S \left(\sum_{x \in S} |D_\lambda^S[x] - D_0^S[x]| \cdot |Q_0(L, x)| \right) \left(\sum_{y \in S} |h(L)[y] D_\lambda^S(y) W_\lambda(y)| \right) = 0 \quad (32)$$

Then

$$\lim_{L \rightarrow \infty} \sum_{x \in S} |h(L)[x] Q_\lambda(L, x) - h(L)[x] Q_0(L, x)| = 0.$$

Proof. Note that, by the definition of $a_\lambda^S(L)$,

$$\begin{aligned} \sum_{x \in S} |h(L)[x] Q_\lambda(L, x) - h(L)[x] Q_0(L, x)| \\ = \sum_{x \in S} \sum_S |a_\lambda^S(L) D_\lambda^S[x] Q_\lambda(L, x) - a_0^S(L) D_0^S[x] Q_0(L, x)| \end{aligned}$$

Recalling the elementary inequality

$$|b_1 b_2 b_3 - c_1 c_2 c_3| = |(b_1 - c_1) c_2 c_3 + b_1 (b_2 - c_2) c_3 + b_1 b_2 (b_3 - c_3)|,$$

we have the upper bound

$$\sum_{x \in \mathcal{S}} \sum_S \left(|(a_\lambda^S(L) - a_0^S(L)) D_0^S[x] Q_0(L, x)| \right. \\ \left. + |a_\lambda^S(L) (D_\lambda^S[x] - D_0^S[x]) Q_0(L, x)| + |a_\lambda^S(L) D_\lambda^S[x] (Q_\lambda(L, x) - Q_0(L, x))| \right).$$

Next, we will express everything in terms of the functions D (note that this can be done because the functions D are orthogonal). Again using the definition of $a_\lambda^S(L)$: the first term is equal to

$$\sum_{x \in \mathcal{S}} \sum_S \sum_{y \in \mathcal{S}} |h(L)[y] (D_\lambda^S(y) W_\lambda(y) - D_0^S(y) W_0(y)) D_0^S[x] Q_0(L, x)|,$$

while the second term is equal to

$$\sum_{x \in \mathcal{S}} \sum_S \sum_{y \in \mathcal{S}} |h(L)[y] D_\lambda^S(y) W_\lambda(y) (D_\lambda^S[x] - D_0^S[x]) Q_0(L, x)|$$

and the third term is equal to

$$\sum_{x \in \mathcal{S}} \sum_S \sum_{y \in \mathcal{S}} |h(L)[y] D_\lambda^S(y) W_\lambda(y) D_\lambda^S[x] (Q_\lambda(L, x) - Q_0(L, x))|.$$

The first term can be re-written as

$$\sum_S \left(\sum_{x \in \mathcal{S}} |D_0^S[x] Q_0(L, x)| \right) \left(\sum_{y \in \mathcal{S}} |h(L)[y]| \cdot |D_\lambda^S(y) W_\lambda(y) - D_0^S(y) W_0(y)| \right)$$

and the second term can be re-written as

$$\sum_S \left(\sum_{x \in \mathcal{S}} |D_\lambda^S[x] - D_0^S[x]| \cdot |Q_0(L, x)| \right) \left(\sum_{y \in \mathcal{S}} |h(L)[y] D_\lambda^S(y) W_\lambda(y)| \right)$$

and the third term can be re-written as

$$\sum_S \left(\sum_{x \in \mathcal{S}} |D_\lambda^S[x]| \cdot (Q_\lambda(L, x) - Q_0(L, x)) \right) \left(\sum_{y \in \mathcal{S}} |h(L)[y] D_\lambda^S(y) W_\lambda(y)| \right).$$

The lemma now follows from the assumed bounds and the dominated convergence theorem.

4.2.1 Matching notation to the dynamic stochastic six vertex model

Proposition 4 holds for general stochastic processes, including the dynamic stochastic six vertex model as stated in Theorem 2. First, let us match the notation of the proposition to the notation of the dynamic stochastic six vertex model. The countable state space \mathcal{S} is simply $\mathcal{S} = \{0, 1\}^{\mathbf{Z}_{>0}}$, which is the set of all possible particle configurations on the positive integers. The entire sample space Ω can be defined as $\{0, 1\}^{\mathbf{Z}_{>0} \times \mathbf{Z}_{>0}}$, which is the set of all possible outcomes of Bernoulli trials at all individual lattice site. Each X_t^λ is measurable with respect to the σ -algebra \mathcal{F}_t^λ , where \mathcal{F}_t^λ is the cylindrical σ -algebra consisting of events of the form

$$(\{a_1, \dots, a_t\} \times \{0, 1\}^{\mathbf{Z}_{>0}})^{\mathbf{Z}_{>0}}$$

where $a_1, \dots, a_t \in \{0, 1\}$. Then $Q_\lambda(t)$ is a probability measure on $\mathcal{S} = \{0, 1\}^{\mathbf{Z}_{>0}}$, and $Q_\lambda(t, x)$ (for $x \in \mathcal{S}$) is the probability that the dynamic stochastic six vertex model (with parameter λ) is in configuration x and time t . The functions D_λ^S , where $S \subseteq \mathbf{Z}_+$ (here we are writing the state space as occupation variables), are the duality functions in equation (16)

$$D_\lambda^S[\mu] = \mathcal{D}_{ort}(\mu, S).$$

To maintain consistency with notation, we will use $D_{-\infty}$ instead of D_0 .

Recall that the duality function is stated in Corollary 1 as

$$\begin{aligned} & \mathcal{D}_{ort}(\mu, \xi) \\ &= \prod_{i=1}^N {}_3\varphi_2 \left(\begin{matrix} q^{2(2J_i - \mu_i)}, q^{2\xi_i}, e^{-2\pi i \lambda} q^{4N_{[1, i-1]}(\mu - J)} \\ q^{4J_i}, 0 \end{matrix} ; q^{-2}, q^{-2} \right) q^{-4J_i N_{[1, i]}(\xi) - 2\xi_i N_{[1, i-1]}(\mu)}, \end{aligned}$$

However, this function is stated where the dynamic stochastic six vertex model has asymmetry to the left, whereas Theorem 2 is stated with asymmetry to the right to match the notation of [8]. For the remainder of this section, we temporarily translate the duality function into the notation of [8]. Reversing the direction of the lattice, the duality function is now

$$\begin{aligned} \mathfrak{D}_M := \prod_{i=0}^M {}_3\varphi_2 \left(\begin{matrix} q^{2(2J_i - \mu_i)}, q^{2\xi_i}, e^{-2\pi i \lambda} q^{4N_{[i+1, M]}(\mu - J)} \\ q^{4J_i}, 0 \end{matrix} ; q^{-2}, q^{-2} \right) \\ \times q^{-4J_i N_{[i, M]}(\xi) - 2\xi_i N_{[i+1, M]}(\mu)}, \quad (33) \end{aligned}$$

where M is a non-negative integer.

4.2.2 A “trick” for rewriting the duality function

Next, we will specialize certain values for J_i . In this case, we will use a “trick,” which is to take $J_0 \rightarrow \infty$ so that the lattice site acts as an “absorbing” site (cf. [25]). The trick is more specific to totally asymmetric models, and will simplify many calculations. In general, recall that duality for asymmetric models with open boundary conditions ([27], [24], [29]) are different than their counterparts on the infinite line, even in the bulk. We will state the re-written duality function as a lemma

Lemma 1. *Fix $J_i = 1/2$ for $i \geq 1$. Suppose that J_0 and M both depend*

on a large parameter K in such a way that $J_0 = M = K$. Then

$$\begin{aligned} \lim_{K \rightarrow \infty} q^{-4K} e^{2\pi i \lambda \xi_0} \left((q^{2\xi_0}; q^{-2})_{\xi_0} \frac{q^{-2\xi_0}}{(q^{-2}; q^{-2})_{\xi_0}} \right)^{-1} \mathfrak{D}_K(\tau_K \mu, \tau_K \xi) \\ = 1_{\{\mu_y=1 \ \forall y \geq 1\}} e^{-2\pi i \lambda |\xi|} \prod_{y \in S} \prod_{z \geq y+1} q^{-21_{\mu_z=0}} \end{aligned}$$

where τ_K is the shift operator

$$(\tau_K \mu)_y = \mu(y + K).$$

Proof.

Now, assume that $J_i = 1/2$ for all $i \geq 1$. Then, using the identities

$$\begin{aligned} N_{[i, M]}(\mu) &= |\mu| - N_{[0, i-1]}(\mu), \\ 4N_{[i+1, M]}(\mu - J) &= 2 \sum_{j=i+1}^M (1_{\{\mu_j=1\}} - 1_{\{\mu_j=0\}}) := 2N_{[i+1, M]}^0(\mu) \end{aligned}$$

the duality function is thus

$$\begin{aligned} D_0 \prod_{i=1}^M {}_3\varphi_2 \left(\begin{matrix} q^{2(1-\mu_i)}, q^{2\xi_i}, e^{-2\pi i \lambda} q^{2N_{[i+1, M]}^0(\mu)} \\ q^2, 0 \end{matrix} ; q^{-2}, q^{-2} \right) \\ \times q^{-2(|\xi| - N_{[0, i-1]}(\xi)) - 2\xi_i(|\mu| - N_{[0, i]}(\mu))}, \end{aligned}$$

where D_0 is term corresponding to $i = 0$ in the summand. The q -hypergeometric series terminates after two terms, so we obtain

$$\begin{aligned} D_0 C(\xi) q^{-2|\xi||\mu|} \prod_{i=1}^M q^{2\xi_i N_{[0, i]}(\mu)} \\ \times \left(1 - \frac{(1 - q^{2(1-\mu_i)})(1 - q^{2\xi_i})(1 - e^{-2\pi i \lambda} q^{2N_{[i+1, M]}^0(\mu)})}{1 - q^2} \frac{q^{-2}}{1 - q^{-2}} \right) \end{aligned}$$

where $C_M(\xi)$ is the constant

$$\prod_{i=1}^M q^{-2(|\xi| - N_{[0, i-1]}(\xi))}.$$

If $\xi_i = 0$ then the term in the product is 1. So let $S = \{y_1, \dots, y_m\} \subseteq \{1, \dots, M\}$ denote the points where $\xi_y = 1$. Then the duality function simplifies to

$$D_0 C_M(\xi) q^{-2|\xi||\mu|} \times \prod_{i=1}^m \left(1_{\{\mu_{y_i}=0\}} q^{2N_{[0, y_i]}(\mu)} e^{-2\pi i \lambda} q^{2N_{[y_i+1, M]}^0(\mu)} + 1_{\{\mu_{y_i}=1\}} q^{2N_{[0, y_i]}(\mu)} \right)$$

This duality function then extends to the half-infinite line by taking $M \rightarrow \infty$. Before doing so, we specialize to the initial conditions of step initial conditions for the stochastic six vertex model, where particles are densely packed to the right of the origin. Therefore, focusing on the terms depending on M , which are

$$C'_M(\xi, \mu) = \prod_{i=1}^M q^{-2(|\xi| - N_{[0, i-1]}(\xi))} \prod_{\{y \in S: \mu_y=0\}} \prod_{z=y+1}^M q^{21_{\{\mu_z=1\}} - 21_{\{\mu_z=0\}}},$$

Note that for large enough values of the index i , we have $|\xi| - N_{[0, i-1]}(\xi) = 0$. So for large enough M , we must have

$$C'_{M+1}(\xi, \mu) = C'_M(\xi, \mu) q^{2|\{y \in S: \mu_y=0\}|}.$$

Thus, the leading order term occurs when $\mu_y = 0$ for every y such that $\xi_y = 1$. Also note the summation of $2N_{[0, y]}(\mu)$ and $2 \sum_{z>y} 1_{\mu_z=1}$ turns into the constant $2K$.

Now, returning to the “trick,” take $J_0 \rightarrow \infty$, so that the “one-site” duality

function there is

$${}_3\varphi_2 \left(\begin{matrix} q^{2(2J_0-\mu_0)}, q^{2\xi_0}, e^{-2\pi i\lambda} q^{4N_{[1,M]}(\mu-J)} \\ q^{4J_0}, 0 \end{matrix}; q^{-2}, q^{-2} \right) q^{-4J_0 N_{[0,M]}(\xi) - 2\xi_0 N_{[1,M]}(\mu)},$$

which equals

$$\lim_{J_0 \rightarrow \infty} q^{-4J_0 N_{[0,M]}(\mu) - 2\xi_0 N_{[1,M]}(\mu)} \times \sum_{n=0}^{\infty} \frac{(e^{-2\pi i\lambda} q^{4N_{[1,M]}(\mu-1/2)}; q^{-2})_n (q^{4J_0-2\mu_0}; q^{-2})_n (q^{2\xi_0}; q^{-2})_n}{(q^{4J_0}; q^{-2})_n} \frac{q^{-2n}}{(q^{-2}; q^{-2})_n}.$$

In fact, for the step initial conditions where $\mu_0 = 0$, then the dependence on J_0 disappears in the summand and we obtain

$$\lim_{J_0 \rightarrow \infty} q^{-4J_0 N_{[0,M]}(\mu) - 2\xi_0 N_{[1,M]}(\mu)} \times \sum_{n=0}^{\infty} (e^{-2\pi i\lambda} q^{4N_{[1,M]}(\mu-1/2)}; q^{-2})_n (q^{2\xi_0}; q^{-2})_n \frac{q^{-2n}}{(q^{-2}; q^{-2})_n}.$$

From here, we take M and J_0 both to infinity. Due to the term $(q^{2\xi_0}; q^{-2})_n$, the series terminates at $n = \xi_0$. Since ξ is the dual process with only finitely many particles, this means that the leading order term is simply

$$q^{-4J_0 N_{[0,M]}(\mu)} q^{-2\xi_0 N_{[1,M]}(\mu)} q^{4\xi_0 N_{[1,M]}(\mu-1/2)}.$$

Note that for μ in the state space, the value of the height function $N_{[0,M]}(\mu)$ will be proportional to M . More specifically, for any fixed μ and large enough M

$$N_{[0,M]}(\mu) = M - \tilde{N}(\mu)$$

where $\tilde{N}(\mu)$ denotes the place where the densely packed region in μ

begins, i.e.

$$\min\{y : \mu_z = 1 \text{ for all } z \geq y\}.$$

Therefore, the leading order term is

$$q^{-4J_0(M-\tilde{N}(\mu))} q^{-2\xi_0\tilde{N}(\mu)}.$$

This is then supported on the μ where $\tilde{N}(\mu) = M$, which is the state obtained where jumps always occur. Finally, accounting for the shift operator results in the indicator function.

4.2.3 Proving the Bounds

To apply proposition 4, we need to prove five separate inequalities. The latter three (the ones bounded by M_1, M_2, M_3) only depend on the duality function D and the measures Q , and not on h or W . We focus on those three inequalities first.

Proving the bound in M_1

Recall that we want an upper bound of the form

$$\sum_{x \in \mathcal{S}} |D_0^S[x] Q_0(L, x)| \leq M_1(L, S).$$

The sum is supported on one element, and $Q_0(L, x)$ is exponentially decaying while $D_0^S[x]$ is bounded, so this bound is trivial.

Proving the bound in M_3

Recall that we want an upper bound of the form

$$\left(\sum_{x \in \mathcal{S}} D_\lambda^S[x] (Q_\lambda(L, x) - Q_0(L, x)) \right) \leq M_3(L, S)$$

This is true for the same reasons as the previous bound.

Proving the bound in M_2

We now move on to the inequality

$$\sum_S \left(\sum_{x \in \mathcal{S}} |D_\lambda^S[x] - D_0^S[x]| \cdot |Q_0(L, x)| \right) \left(\sum_{z \in \mathcal{S}} |h(L)[z] D_\lambda^S(z) W_\lambda(z)| \right) \rightarrow 0$$

as $L \rightarrow \infty$. The term in the first parentheses is exponentially decaying in L while the term in the second parentheses is bounded. The summation is over a set which is growing polynomially in L , thus the bound holds.

Proving the bounds in C_1, C_2

Recall that we want to prove

$$\left(\sum_{y \in \mathcal{S}} h(L)[y] \cdot |D_\lambda^S(y) W_\lambda(y) - D_0^S(y) W_0(y)| \right) \leq C_1 L^2$$

and

$$\sum_{y \in \mathcal{S}} h(L)[y] D_\lambda^S(y) W_\lambda(y) \leq C_2 L^2.$$

If the second bound holds for all values of λ then taking its difference at $\lambda = 0$ proves the first bound. This bound holds because the sum only is nonzero for one term, and $h(L)[y]$ is linear in L and the W term is bounded.

Finally, we explain the Tracy–Widom convergence. Thus far, all we have done is compare the height function of the dynamic stochastic six vertex model to the usual stochastic six vertex model, and have not considered any particular scaling. The expression

$$\sum_{x \in \mathcal{S}} h(L)[x] Q_\lambda(L, x)$$

simply equals

$$\mathbf{E} [h(L)[X_L^\lambda]],$$

so the lemma states that the random variable

$$h(L)[X_L^\lambda] - h(L)[X_L^0]$$

converges to 0 in mean. But since $h(L)[X_L^0]$ converges in distribution to Tracy–Widom distribution, then so does $h(L)[X_L^\lambda]$.

Remark 6. If the limit of the duality function were computed without the “trick” then the duality would be a product of linear combinations of the duality functions of [35] and [25] (see also [32] and [28]). These two duality functions are essentially equivalent to each other up to $q \leftrightarrow q^{-1}$; furthermore note that the dynamic parameter essentially interpolates between the two values q and q^{-1} . Thus, the duality function provides a heuristic for why the dynamic parameter is irrelevant in the asymptotics: if the dynamic parameter were scaled large enough to see asymptotics other than Tracy–Widom, then it would be the asymptotic regime where q is replaced by q^{-1} ; but in this regime, there is no liquid region with nontrivial asymptotics.

Remark 7. The duality function of [35] can be modified (as in [9]) to give the q –moments of the height function. Thus, asymptotically, the duality function here can be related to these q –moments of the height function.

Remark 8. In section 7 of [1], it is proven that the dynamic (J, γ) –PEP has different fluctuations - namely, the height function rescaled by $T^{1/4}$ converges to a non–deterministic limit, rather than converging to a deterministic limit with linear rescaling. A natural question is how to “reconcile” the two different scaling regimes. First, note that the dynamic (J, γ) –PEP was constructed as a degeneration of the dynamic q –Hahn Boson. While the usual q –Hahn Boson does satisfy a duality [16], [25], the duality is not known to have an algebraic origin. Rather, the algebraically–constructed duality for the q –Boson is manually checked to hold for the q –Hahn Bo-

son. Therefore, the algebraic intuition that the quantum group is invariant under $q \leftrightarrow q^{-1}$ does not necessarily apply to the dynamic q -Hahn Boson. This intuition does hold for the “dynamic” q -Boson; but note that the dynamic q -Boson itself does not depend on the dynamic parameter anyway (example 6.4 of [1]; see also [19] and [27]). This intuition would seem to indicate that the duality for the q -Hahn Boson can *not* be proven from quantum groups.

Now move on to the first probabilistic result, Theorem 1, which follows from Theorem 3, Theorem 4, Corollary 3, Proposition 1, Proposition 2 and 3. Each of these will be proved in one of the subsections below.

4.3 Proof of Theorem 3

4.3.1 Yang-Baxter-Equation

From the dynamical Yang–Baxter equation and replacing the lattice sites $\{1, 2, 3\}$ with $\{0, i, i + 1\}$, we get

$$\begin{aligned} R_{0,i}(u, \lambda - 2\eta H_{i+1})R_{0,i+1}(uv, \lambda)R_{i,i+1}(v, \lambda - 2\eta H_0) \\ = R_{i,i+1}(v, \lambda)R_{0,i+1}(uv, \lambda - 2\eta H_i)R_{0,i}(u, \lambda). \end{aligned}$$

Now letting $P_{i,j}$ be the permutation operator that permutes site i and j , and since P^2 is the identity,

$$\begin{aligned} P_{i,i+1}R_{0,i}(u, \lambda - 2\eta H_{i+1})P_{i,i+1}P_{i,i+1}R_{0,i+1}(uv, \lambda)P_{i,i+1}P_{i,i+1}R_{i,i+1}(v, \lambda - 2\eta H_0)P_{i,i+1} \\ = P_{i,i+1}R_{i,i+1}(v, \lambda)R_{0,i+1}(uv, \lambda - 2\eta H_i)R_{0,i}(u, \lambda)P_{i,i+1}. \end{aligned}$$

Now using that

$$P_{j,k}R_{i,j}(z, \lambda)P_{j,k} = R_{i,k}(z, \lambda), \quad P_{j,k}R_{i,j}(z, \lambda - 2\eta H_k)P_{j,k} = R_{i,k}(z, \lambda - 2\eta H_j),$$

we get

$$\begin{aligned} R_{0,i+1}(u, \lambda - 2\eta H_i) R_{0,i}(uv, \lambda) R_{i+1,i}(v, \lambda - 2\eta H_0) \\ = P_{i,i+1} R_{i,i+1}(v, \lambda) R_{0,i+1}(uv, \lambda - 2\eta H_i) R_{0,i}(u, \lambda) P_{i,i+1}. \end{aligned}$$

Plugging in

$$\check{R}_{i,i+1}(z, \lambda) := P_{i,i+1} R_{i,i+1}(z, \lambda) = R_{i+1,i}(z, \lambda) P_{i,i+1},$$

we obtain

$$\begin{aligned} R_{0,i+1}(u, \lambda - 2\eta H_i) R_{0,i}(uv, \lambda) \check{R}_{i,i+1}(v, \lambda - 2\eta H_0) \\ = \check{R}_{i,i+1}(v, \lambda) R_{0,i+1}(uv, \lambda - 2\eta H_i) R_{0,i}(u, \lambda). \end{aligned}$$

Thus, for any $i = 1, \dots, N - 1$,

$$\mathcal{B}(u, z, \lambda) \check{R}_{i,i+1}(q, \lambda - 2\eta N_{[1,i-1]}(H) - 2\eta H_0) = \check{R}_{i,i+1}(q, \lambda - 2\eta N_{[1,i-1]}(H)) \mathcal{B}(u, z, \lambda).$$

As a note, the H_0 on the left-hand-side explains why a dynamic ASEP can be dual to a usual ASEP; namely, when $H_0 = \infty$ then the value of λ is irrelevant.

4.3.2 Reducing from \mathcal{B} to B

We continue from the last section, now defining, for each $m, l \leq \dim V_0$

$$\langle \mu | B_{m,l}(u, z, \lambda) | \xi \rangle = \langle m, \mu | \mathcal{B}(u, z, \lambda) | l, \xi \rangle$$

we have that

$$\begin{aligned} \langle m, \mu | \check{R}_{i,i+1}(z, \lambda - 2\eta N_{[1,i-1]}(H)) \mathcal{B}(u, z, \lambda) | l, \xi \rangle = \\ \langle m, \mu | \mathcal{B}(u, z, \lambda) (\check{R}_{i,i+1}(z, \lambda - 2\eta N_{[1,i-1]}(H) - 2\eta H_0)) | l, \xi \rangle \end{aligned}$$

with $m + N(\mu) = l + N(\xi)$, where $N(\mu)$ is the total number of particles in μ . This implies

$$\begin{aligned} & \sum_{\nu} \langle \mu | \check{R}_{i,i+1}(z, \lambda - 2\eta N_{[1,i-1]}(H)) | \nu \rangle \langle m, \nu | \mathcal{B}(u, z, \lambda) | l, \xi \rangle \\ &= \sum_{\xi} \langle m, \mu | \mathcal{B}(u, z, \lambda) | l, \xi \rangle \langle l, \xi | \check{R}_{i,i+1}(z, \lambda - 2\eta N_{[1,i-1]}(H) - 2\eta H_0) | l, \xi \rangle, \end{aligned}$$

which can be stated as

$$\begin{aligned} & \check{R}_{i,i+1}(z, \lambda - 2\eta N_{[1,i-1]}(H)) B_{m,l}(u, z, \lambda) \\ &= B_{m,l}(u, z, \lambda) \check{R}_{i,i+1}(z, \lambda - 2\eta N_{[1,i-1]}(H) - 2\eta H_0(l)) \end{aligned}$$

for all values of m and l . Here, $H_0(l)$ is the constant $2(l - J_0)$.

4.3.3 Transposition

Recall that $R^D(z)$ satisfies

$$\Pi^{\otimes 2} R^D(z) = (R^D(z))^* \Pi^{\otimes 2},$$

where Π is a permutation matrix. More specifically,

$$\langle \beta | \Pi | \alpha \rangle = 1_{\{\beta=2J-\alpha\}}.$$

Now, by (14) and the fact that P intertwines with $\Pi^{\otimes 2}$, we get

$$(F_{21}(\lambda)^*)^{-1} \Pi^{\otimes 2} F_{12}(\lambda)^{-1} \check{R}(z, \lambda) = P R(z, \lambda)^* (F_{21}(\lambda)^*)^{-1} \Pi^{\otimes 2} F_{12}(\lambda)^{-1}.$$

Notice that by definition,

$$\check{\mathfrak{R}}^*(z, \lambda) = \check{R}(z, \lambda).$$

For an index of numbers I , let $\Pi_I = \prod_{i \in I} \Pi_i$. Therefore we can write

$$\check{R}_{i,i+1}(z, \lambda - 2\eta N_{[1,i-1]}(H)) B_{m,l}(u, z, \lambda) = B_{m,l}(u, z, \lambda) \check{\mathfrak{R}}^*(z, \lambda(l)),$$

where $\lambda(l) = \lambda - 2\eta N_{[1,i-1]}(H) - 2\eta H_0(l)$.

4.4 Proof of Proposition 1

In this section, we show that the dynamic R matrix satisfies the transposition formula (21), there exists a ground state symmetry (22), and the stochastic S matrix satisfies the inversion formula (24).

4.4.1 Transposition formula

First of all, there is a transposition formula for $R^D(z)$ [31]

$$\Pi^{\otimes 2} \Lambda^{\otimes 2} R^D(z) = R^D(z)^* \Pi^{\otimes 2} \Lambda^{\otimes 2}.$$

Applying the permutation operator P and plugging in (14) yields (21). Since we used a different representation from [31], we calculate Λ here.

Proposition 5. $\Lambda = \text{id}$ and $\Pi^{\otimes 2} R^D(z) = R^D(z)^* \Pi^{\otimes 2}$.

Proof.

As shown in [31], Λ does not depend on z , thus we take $z = 0$ and explicitly solve for Λ .

Recall

$$R_{12}^D(0) = q^{\frac{1}{2}H \otimes H} \sum_{i=0}^{\infty} (q - q^{-1})^i \frac{q^{-\frac{i(i+1)}{2}}}{[i]!} q^{\frac{i}{2}H} E_+^i \otimes q^{-\frac{i}{2}H} E_-^i.$$

Then

$$(R_{12}^D(0))^* = \sum_{i=0}^{\infty} (q - q^{-1})^i \frac{q^{-\frac{i(i+1)}{2}}}{[i]!} E_-^i q^{\frac{i}{2}H} \otimes E_+^i q^{-\frac{i}{2}H} q^{\frac{1}{2}H \otimes H}.$$

When $i = \eta_1 - \xi_1 = \xi_2 - \eta_2 \geq 0$

$$\begin{aligned} \frac{\langle \eta_1, \eta_2 | \Lambda^{\otimes 2} | \eta_1, \eta_2 \rangle}{\langle \xi_1, \xi_2 | \Lambda^{\otimes 2} | \xi_1, \xi_2 \rangle} &= \frac{\langle \eta_1, \eta_2 | R^D(0) | \xi_1, \xi_2 \rangle}{\langle 2J_1 - \eta_1, 2J_2 - \eta_2 | R^D(0)^* | 2J_1 - \xi_1, 2J_2 - \xi_2 \rangle} \\ &= q^{2(\eta_1 - J_1)(\eta_2 - J_2) - 2(J_1 - \xi_1)(J_2 - \xi_2) + i(\eta_1 - \eta_2 - \xi_2 + \xi_1 - 2J_1 + 2J_2)} \\ &= q^{(\eta_1 - 2J_1 + \xi_1)(\eta_1 + \eta_2 - \xi_1 - \xi_2)} = 1. \end{aligned}$$

Thus $\langle \beta | \Lambda | \alpha \rangle = 1_{\{\alpha = \beta\}}$.

4.4.2 The Ground State Transformation

From the form of the universal R -matrix in subsection 2.3.1, it is immediate that

$$\check{R}_{i,j}(z, \lambda) |0, 0\rangle = |0, 0\rangle. \quad (34)$$

Taking $N = 2$ in $B_{m,l}$, we obtain

$$\begin{aligned} \check{R}_{i,i+1}(z, \lambda) B_{m,l}(u, z, \lambda) |0, 0\rangle &= B_{m,l}(u, z, \lambda) \check{R}_{i,i+1}(z, \lambda - 2\eta h^0(l)) |0, 0\rangle \\ &= B_{ml}^1(u, z, \lambda) |0, 0\rangle. \end{aligned}$$

In words, $B_{m,l}(u, z\lambda) |0, 0\rangle$ is an eigenvector of $\check{R}_{i,i+1}(z, \lambda)$ of eigenvalue 1.

Consider the value

$$\langle \xi | G_{i,i+1}(\lambda) | \xi \rangle = \langle \xi | B_{m,l}(u, z\lambda) |0, 0\rangle, \quad (35)$$

where for weight conservation we require $m + \xi_1 + \xi_2 = l$. Now note that

$$\begin{aligned}
& \sum_{\mu} \langle \xi_{i+1}, \xi_i | \check{S}_{i,i+1}(z, \lambda) | \mu_i, \mu_{i+1} \rangle \\
&= \sum_{\mu} \frac{\langle \mu_i, \mu_{i+1} | G_{i,i+1} | \mu_i, \mu_{i+1} \rangle}{\langle \xi_i, \xi_{i+1} | G_{i,i+1} | \xi_i, \xi_{i+1} \rangle} \langle \xi_{i+1}, \xi_i | \check{R}_{i,i+1}(z, \lambda) | \mu_i, \mu_{i+1} \rangle \\
&= \frac{\langle \xi_{i+1}, \xi_i | \check{R}_{i,i+1}(z, \lambda) B_{m,l}(u, z, \lambda) | 0, 0 \rangle}{\langle \xi_{i+1}, \xi_i | B_{m,l}(u, z, \lambda) | 0, 0 \rangle} \\
&= 1,
\end{aligned}$$

demonstrating that the ground state transformation exists.

4.4.3 Inversion formula

We see that

$$\begin{aligned}
b_0^{\text{stoch}}(q^{-1}, \lambda, w^{-1}) &= \frac{1 - q}{1 - q^{1/2}w^{-1}} \cdot \frac{q^{-1/2}w^{-1} - e^{2\pi i\lambda}}{1 - e^{2\pi i\lambda}} \\
&= \frac{(1 - q)wq^{-1/2}}{wq^{-1/2} - 1} \cdot \frac{q^{-1/2}w^{-1} - e^{2\pi i\lambda}}{1 - e^{2\pi i\lambda}} \\
&= c_1^{\text{stoch}}(q, \lambda, w).
\end{aligned}$$

and also

$$\begin{aligned}
a_1^{\text{stoch}}(q^{-1}, \lambda, w^{-1}) &= \frac{1 - q^{-1/2}w^{-1}}{1 - q^{1/2}w^{-1}} \frac{q - e^{2\pi i\lambda}}{1 - e^{2\pi i\lambda}} \\
&= \frac{wq^{-1/2}(1 - q^{-1/2}w^{-1})}{wq^{-1/2} - 1} \frac{q - e^{2\pi i\lambda}}{1 - e^{2\pi i\lambda}} \\
&= d_0^{\text{stoch}}(q, \lambda, w).
\end{aligned}$$

Thus, this proves that Π must exist in the spin 1/2 case. For the general case, the standard fusion argument applies (e.g. section 4.3 of [1] or section 3.2 of [25]).

4.5 Proof of Theorem 4 and Corollary 3

Theorem 4 is obtained by plugging (23) and (25) in (20) and taking $H_0(l) \rightarrow \infty$. To extend to multiple sites, we first extend $G_{i,i+1}$ to multiple sites.

Proposition 6. *Define*

$$\langle \mu | \mathcal{G}(\lambda) | \mu \rangle = \prod_{i=1}^N \frac{q^{2J_i N_{[1,i-1]}(\mu) + J_i \mu_i}}{\sqrt{[\mu_i]! [2J_i - \mu_i]!} (e^{-2\pi i \lambda} q^{4N_{[1,i]}(\mu - J) + 2}; q^2)_{2J_i - \mu_i}}.$$

Then

$$\mathcal{T}^{\text{stoch}}(z, \lambda) = \mathcal{G}^{-1}(\lambda) \mathcal{T}(z, \lambda) \mathcal{G}(\lambda).$$

Proof. Similar to the two site case in section 4.4.2, $B_{m,l}(\lambda) | \vec{0} \rangle$ is an eigenvector of $\mathcal{T}(z, \lambda)$, so let

$$\begin{aligned} \langle \mu | \mathcal{G}(\lambda) | \mu \rangle &= \lim_{J_0 \rightarrow \infty} \langle \mu | B_{2J_0 - N(\mu), 2J_0}(\lambda) | \vec{0} \rangle \\ &= C(N(\mu)) \prod_{i=1}^N \frac{q^{2J_i N_{[1,i-1]}(\mu) + J_i \mu_i}}{\sqrt{[\mu_i]! [2J_i - \mu_i]!} (e^{-2\pi i \lambda} q^{4N_{[1,i]}(\mu - J) + 2}; q^2)_{2J_i - \mu_i}}. \end{aligned}$$

Then conjugating $\mathcal{T}(z, \lambda)$ by \mathcal{G} makes it stochastic, and we will show it is equal to $\mathcal{T}^{\text{stoch}}$ by induction.

When $N = 2$,

$$\langle \mu | \mathcal{G}(\lambda) | \mu \rangle = \langle \mu | G_{1,2}(\lambda) | \mu \rangle.$$

Now we assume there are N sites, so define

$$\mathcal{T}_N^{\text{stoch}}(z, \lambda) = \mathcal{G}_N^{-1}(\lambda) \mathcal{T}_N(z, \lambda) \mathcal{G}_N(\lambda),$$

so it suffices to show that

$$\check{S}_{N,N+1}(z, \lambda - 2\eta N_{[1,N-1]}(H)) \mathcal{T}_N^{\text{stoch}}(z, \lambda) = \mathcal{G}_{N+1}^{-1} \check{R}_{N,N+1}(z, \lambda - 2\eta N_{[1,N-1]}(H)) \mathcal{T}_N(z, \lambda) \mathcal{G}_{N+1},$$

Recalling that $\check{R}_{i,i+1} : V_1 \otimes V_{i+1} \rightarrow V_{i+1} \otimes V_1$, we obtain

$$\begin{aligned} & \frac{\langle \xi | G_{N,N+1}(\lambda - 2\eta N_{[1,N-1]}(H)) \check{R}_{N-1,N} | \mu \rangle}{\langle \xi | \check{R}_{N-1,N} G_{N,N+1}(\lambda - 2\eta N_{[1,N-1]}(H)) | \mu \rangle} \\ &= \frac{\langle \xi_1, \xi_{N+1} | G(\lambda - 2\eta N_{[1,N-1]}(H)) |_{V_1 \otimes V_{N+1}} | \xi_1, \xi_{N+1} \rangle}{\langle \mu_N, \mu_{N+1} | G(\lambda - 2\eta N_{[1,N-1]}(H)) |_{V_i \otimes V_{N+1}} | \mu_N, \mu_{N+1} \rangle} \\ &= \frac{\sqrt{[2J_N - \mu_N]! [\mu_N]!} (e^{-2\pi i \lambda} q^{4N_{[1,N]}(\mu-J)+2}; q^2)_{2J_N - \mu_N} q^{(J_1+2J_{N+1})\xi_1 - (J_N+2J_{N+1})\mu_N}}{\sqrt{[2J_1 - \xi_1]! [\xi_1]!} (e^{-2\pi i \lambda} q^{4N_{[1,N]}(\xi-J)+2}; q^2)_{2J_1 - \xi_1}}. \end{aligned}$$

and

$$\begin{aligned} & \frac{\langle \xi | G_{N,N+1}(\lambda - 2\eta N_{[1,N-1]}(H)) \mathcal{T}_N(z, \lambda) \mathcal{G}_N(\lambda) | \mu \rangle}{\langle \xi | \mathcal{T}_N(z, \lambda) \mathcal{G}_{N+1}(\lambda) | \mu \rangle} \\ &= \frac{\langle \xi | G_{N,N+1}(\lambda - 2\eta N_{[1,N-1]}(H)) \mathcal{T}_N(z, \lambda) | \mu \rangle \langle \mu | G_{N,N+1}(\lambda - 2\eta N_{[1,N-1]}(H)) \mathcal{G}_N(\lambda) | \mu \rangle}{\langle \xi | \mathcal{T}_N(z, \lambda) G_{N,N+1}(\lambda - 2\eta N_{[1,N-1]}(H)) | \mu \rangle \langle \mu | \mathcal{G}_{N+1}(\lambda) | \mu \rangle} \\ &= \frac{1}{\sqrt{[2J_1 - \xi_1]! [\xi_1]!}} q^{-2J_{N+1}N_{[1,N]}(\xi) + (J_1+2J_{N+1})\xi_1}. \end{aligned}$$

Similarly,

$$\begin{aligned} \frac{\langle \xi | \check{R}_{N,N+1} \mathcal{G}_N^{-1} | \mu \rangle}{\langle \xi | \mathcal{G}_N^{-1} \check{R}_{N,N+1} | \mu \rangle} &= \frac{\sqrt{[2J_1 - \mu_1]! [\mu_1]!} (e^{-2\pi i \lambda} q^{4N_{[1,N]}(\mu-J)+2}; q^2)_{2J_1 - \mu_1}}{\sqrt{[2J_{N+1} - \xi_{N+1}]! [\xi_{N+1}]!} (e^{-2\pi i \lambda} q^{4N_{[2,N+1]}(\xi-J)+2}; q^2)_{2J_{N+1} - \xi_{N+1}}} \\ & \quad q^{2(J_{N+1}-J_1)N_{[2,N]}(\xi) + J_{N+1}\xi_{N+1} - J_1\mu_1}. \end{aligned}$$

Thus,

$$\begin{aligned}
& \frac{\langle \xi | G_{N,N+1}^{-1}(\lambda - 2\eta N_{[1,N-1]}(H)) \check{R}_{N,N+1} \mathcal{G}_N^{-1} | \mu \rangle}{\langle \xi | \mathcal{G}_{N+1}^{-1} \check{R}_{N,N+1} | \mu \rangle} \\
&= \frac{\langle \xi | \check{R}_{N,N+1} \mathcal{G}_N^{-1} | \mu \rangle \langle \xi | G_{N,N+1}^{-1}(\lambda - 2\eta N_{[1,N-1]}(H)) \mathcal{G}_N^{-1} | \xi \rangle}{\langle \xi | \mathcal{G}_N^{-1} \check{R}_{N,N+1} | \mu \rangle \langle \xi | \mathcal{G}_{N+1}^{-1} | \xi \rangle} \\
&= \sqrt{[2J_1 - \mu_1]! [\mu_1]!} q^{2J_{N+1} N_{[2,N]}(\mu) - J_1 \mu_1}.
\end{aligned}$$

Combining the above equations, we obtain the desired equality. Q.E.D.

To finish the proof of Corollary 3, we need to extend from a single time update to multiple time updates. These are standard arguments, which we repeat below. First, to interpret the intertwining of transfer matrices as a particle system satisfying duality, we apply the same arguments as in section 4.4 of [25]. Also, note that the dynamic parameter changes in each time step. Since updating each time step is equivalent to fusing the horizontal spin and the dynamic parameter changes in fusion (section 4.3 of [1], our duality result extends to multiple time steps.

4.6 Proof of Proposition 2

Let $\sigma'_i = \mu_i - J_i$ and $\sigma_i = \xi_i - J_i$ ($-J_i \leq \sigma_i, \sigma'_i \leq J_i$). By an automorphism between the current paper and the setups in [5] we see that the action of $R(\infty, \lambda)$ on $V^{2J_1} \otimes V^{2J_2}$ has the following entries

$$\begin{aligned}
& \langle J_1 + \sigma'_1 | \langle J_2 + \sigma'_2 | R^{2J_1, 2J_2}(\infty, \lambda) | J_1 + \sigma_1 \rangle | J_2 + \sigma_2 \rangle \\
&= (-1)^{\sigma'_2 - \sigma_2} q^{C_{j(\lambda)} + C_{j(\lambda) + \sigma_1 + \sigma_2} - C_{j(\lambda) + \sigma_1} - C_{j(\lambda) + \sigma'_2}} \\
&\times \frac{\mathcal{N}_\psi^{(J_2)}(q^{2j(\lambda)+1}, \sigma'_2) \mathcal{N}_\psi^{(J_1)}(q^{2j(\lambda)+1} q^{2\sigma'_2}, \sigma'_1)}{\mathcal{N}_\psi^{(J_2)}(q^{2j(\lambda)+1} q^{2\sigma_1}, \sigma_2) \mathcal{N}_\psi^{(J_1)}(q^{2j(\lambda)+1}, \sigma_1)} \left\{ \begin{matrix} J_1 & j(\lambda) + \sigma_1 + \sigma_2 & j(\lambda) + \sigma'_2 \\ J_2 & j(\lambda) & j(\lambda) + \sigma_1 \end{matrix} \right\}_q,
\end{aligned} \tag{36}$$

where $q^{2j(\lambda)+1} := e^{-\pi i \lambda}$, $C_n = n(n+1)$, the last symbol represents the 6-j coefficient (3) (also see eq. (2.54) in [14]) and

$$\begin{aligned} & \mathcal{N}_\psi^{(J_1)} \left(q^{2j(\lambda)+1}, \sigma_1 \right) \\ &= (-1)^{-2J_1 + \frac{\sigma_1}{2}} \frac{\sqrt{[J_1 + 2j(\lambda) + \sigma_1 + 1]!}}{(q^{4j(\lambda)+4}; q^2)_{J_1 + \sigma_1}} \frac{(q - q^{-1})^{J_1} q^{(2j(\lambda)+1)J_1} q^{J_1 \sigma_1}}{\sqrt{[2j(\lambda) + 1 + 2\sigma_1] [-J_1 + 2j(\lambda) + \sigma_1]!}}. \end{aligned}$$

For simplicity, denote

$$f_{\mathcal{N}} = \frac{\mathcal{N}_\psi^{(J_2)} \left(q^{2j(\lambda)+1}, \sigma'_2 \right) \mathcal{N}_\psi^{(J_1)} \left(q^{2j(\lambda)+1} q^{2\sigma'_2}, \sigma'_1 \right)}{\mathcal{N}_\psi^{(J_2)} \left(q^{2j(\lambda)+1} q^{2\sigma_1}, \sigma_2 \right) \mathcal{N}_\psi^{(J_1)} \left(q^{2j(\lambda)+1}, \sigma_1 \right)}$$

and

$$f_{6j} = (-1)^{\sigma'_2 - \sigma_2} q^{C_{j(\lambda)} + C_{j(\lambda) + \sigma_1 + \sigma_2} - C_{j(\lambda) + \sigma_1} - C_{j(\lambda) + \sigma'_2}} \left\{ \begin{array}{ccc} J_1 & j(\lambda) + \sigma_1 + \sigma_2 & j(\lambda) + \sigma'_2 \\ J_2 & j(\lambda) & j(\lambda) + \sigma_1 \end{array} \right\}_q.$$

4.6.1 Calculating limits of $B_{m,l}(u, z, \lambda)$

In this section, we calculate several limits of $B_{m,l}(u, z, \lambda)$. For simplicity, we only show an example when $u = \infty$, other values can be calculated in the same way.

Recall the fact that as $N \rightarrow \infty$

$$\frac{[\alpha \pm N]!}{[\beta \pm N]!} \sim (\mp)^{\alpha - \beta} \frac{q^{\mp \frac{1}{2}(\alpha - \beta)(\alpha + \beta + 1)}}{(q - q^{-1})^{\alpha - \beta}} q^{-(\alpha - \beta)N}.$$

We make use of this fact repeatedly to obtain the following limits, which are stated in the next proposition.

Proposition 7. *Suppose $\dim V_i = 2J_i + 1$ when $i \geq 0$, $c = 2J_0 - l$ is a fixed positive integer. Let $J = (J_1, \dots, J_N)$, then up to a constant in $N(\xi)$*

and $N(\mu)$, we have the limits:

$$\begin{aligned}
& \lim_{J_0 \rightarrow \infty} \langle \mu | B_{2J_0-c+N(\xi-\mu), 2J_0-c}(\infty, \lambda) | \xi \rangle \\
&= \prod_{i=1}^N \frac{(q^{2(2j(\lambda)-c+N_{[1,i-1]}(\mu+\xi-2J)+\mu_i-2J_i+1)}; q^2)_{2J_i-\mu_i}}{\left(q^{4(j(\lambda)+1+N_{[1,i]}(\mu-J))}; q^2 \right)_{2J_i-\mu_i} \sqrt{[\mu_i]![2J_i-\xi_i]![2J_i-\mu_i]![\xi_i]!}} \\
& \quad 3\varphi_2 \left(\begin{matrix} q^{-2(2J_i-\mu_i)}, q^{-2\xi_i}, q^{2(2j(\lambda)+2N_{[1,i-1]}(\mu-J)+\mu_i-2J_i+1)} \\ q^{-4J_i}, q^{2(2j(\lambda)-c+N_{[1,i-1]}(\mu+\xi-2J)+\mu_i-2J_i+1)} \end{matrix}, q^2, q^2 \right) \\
& \quad q^{2J_i N_{[1,i-1]}(\mu-\xi)+J_i(\mu_i+\xi_i)-2\xi_i N_{[1,i]}(\mu)},
\end{aligned}$$

$$\begin{aligned}
& \lim_{\substack{J_0 \rightarrow \infty \\ j(\lambda) \rightarrow \infty}} \langle \mu | B_{2J_0-c+N(\xi-\mu), 2J_0-c}(\infty, \lambda) | \xi \rangle \\
&= \prod_{i=1}^N 1_{\{\mu_i \geq \xi_i\}} \frac{1}{[\mu_i - \xi_i]!} \sqrt{\frac{[\mu_i]![2J_i - \xi_i]!}{[2J_i - \mu_i]![\xi_i]!}} \\
& \quad \times q^{2J_i N_{[1,i-1]}(\mu-\xi)+J_i(\mu_i-\xi_i)+\mu_i \xi_i+2\mu_i N_{[1,i-1]}(\xi)},
\end{aligned}$$

$$\begin{aligned}
& \lim_{\substack{j(\lambda) \rightarrow \infty \\ 2j(\lambda)-c \rightarrow C_0}} \lim_{J_0 \rightarrow \infty} \langle \mu | B_{2J_0-c+N(\xi-\mu), 2J_0-c}(\infty, \lambda) | \xi \rangle \\
&= \prod_{i=1}^N \frac{(q^{2(C_0+N_{[1,i-1]}(\mu+\xi-2J)+\mu_i-2J_i+1)}; q^2)_{2J_i-\mu_i}}{\left(q^{4(j(\lambda)+1+N_{[1,i]}(\mu-J))}; q^2 \right)_{2J_i-\mu_i} \sqrt{[\mu_i]![2J_i-\xi_i]![2J_i-\mu_i]![\xi_i]!}} \\
& \quad 3\varphi_2 \left(\begin{matrix} q^{-2(2J_i-\mu_i)}, q^{-2\xi_i}, 0 \\ q^{-4J_i}, q^{2(C_0+N_{[1,i-1]}(\mu+\xi-2J)+\mu_i-2J_i+1)} \end{matrix}, q^2, q^2 \right) \\
& \quad q^{2J_i N_{[1,i-1]}(\mu-\xi)+J_i(\mu_i+\xi_i)-2\xi_i N_{[1,i]}(\mu)},
\end{aligned}$$

$$\begin{aligned}
& \lim_{c \rightarrow \infty} \lim_{J_0 \rightarrow \infty} \langle \mu | B_{2J_0-c+N(\xi-\mu), 2J_0-c}(\infty, \lambda) | \xi \rangle \\
&= \prod_{i=1}^N \frac{[2J_i]! q^{2J_i N_{[1,i-1]}(\mu+\xi)+J_i(\mu_i+\xi_i)}}{\sqrt{[\mu_i]![2J_i-\xi_i]![2J_i-\mu_i]![\xi_i]!} \left(q^{4(j(\lambda)+1+N_{[1,i]}(\mu-J))}; q^2 \right)_{2J_i-\mu_i}}.
\end{aligned}$$

Proof.

We consider the limit of (36) as $J_1 \rightarrow \infty$ with $J_1 - \sigma_1 = c$, where c is a fixed integer such that $-J_i \leq \sigma_i, \sigma'_i \leq J_i$. Denote $c' = J_1 - \sigma'_1 = c + \sigma'_2 - \sigma_2$. Then

$$\lim_{\substack{J_1 \rightarrow \infty \\ J_1 - \sigma_1 = c}} f_{\mathcal{N}} = \frac{(-1)^{1/2\sigma_2 + 1/2\sigma'_2 - J_2}}{(q - q^{-1})^{1/2\sigma_2 + 1/2\sigma'_2 - J_2} (q^{4j(\lambda) + 4 + 4\sigma'_2}; q^2)_{J_2 - \sigma'_2}} q^{J_2 - \sigma_2/4 - (3\sigma'_2)/4 + J_2\sigma'_2} \\ q^{2J_2j(\lambda) - j(\lambda)\sigma_2 - j(\lambda)\sigma'_2 + (c\sigma_2)/2 + (c\sigma'_2)/2 - (\sigma_2\sigma'_2)/2 - \sigma_2^2/4 - \sigma_2'^2/4} \\ \frac{\sqrt{[-c + 2j(\lambda)]! [\sigma'_2 + J_2 + 2j(\lambda) + 1]!}}{\sqrt{[2j(\lambda) + 1 + 2\sigma'_2] [2\sigma'_2 + 2j(\lambda) - c']! [-J_2 + \sigma'_2 + 2j(\lambda)]!}},$$

and

$$\lim_{\substack{J_1 \rightarrow \infty \\ J_1 - \sigma_1 = c}} f_{6j} = (-1)^{4j(\lambda) + \sigma'_2 - \sigma_2} q^{\sigma_2 - J_2^2/2 - J_2/2 + J_2\sigma_2 - \sigma'_2/2 + 2J_1\sigma_2} \\ q^{c(-J_2 - 2\sigma_2) + j(\lambda)(\sigma_2 - \sigma'_2) + \sigma_2^2 - (\sigma'_2)^2/2} \sqrt{[2j(\lambda) + 2\sigma'_2 + 1] \Delta(J_2, j(\lambda), j(\lambda) + \sigma'_2)} \\ \times \sqrt{[J_2 + \sigma_2]! [J_2 - \sigma_2]! [2j(\lambda) - c]! [c]! [c']! [2j(\lambda) + 2\sigma'_2 - c']!} \\ \sum_w \left(\frac{(-1)^{-w} q^{w(J_2 + 2j(\lambda) + \sigma'_2 + 1)}}{[c - w]! [J_2 - \sigma'_2 - w]! [J_2 + \sigma_2 - w]! [w]! [w + \sigma'_2 - \sigma_2]! [w + 2j(\lambda) + \sigma'_2 - J_2 - c]!} \right)$$

Thus

$$\lim_{\substack{J_1 \rightarrow \infty \\ J_1 - \sigma_1 = c}} \langle J_1 + \sigma'_1 | \langle J_2 + \sigma'_2 | R^{2J_1, 2J_2}(\infty, \lambda) | J_1 + \sigma_1 \rangle | J_2 + \sigma_2 \rangle \\ = C_1(J_1, J_2, \sigma_2, \sigma'_2) \mathcal{F}(J_2, j(\lambda), c, \sigma_2, \sigma'_2),$$

where

$$\mathcal{F}(J_2, j(\lambda), c, \sigma_2, \sigma'_2) = \frac{[-c + 2j(\lambda)]!}{(q^{4j(\lambda)+4+4\sigma'_2}; q^2)_{J_2-\sigma'_2} [2j(\lambda) + \sigma'_2 - J_2 - c]! [\sigma'_2 - \sigma_2]!}$$

$$\sqrt{\frac{[J_2 + \sigma'_2]! [J_2 - \sigma_2]! [c']!}{[J_2 - \sigma'_2]! [J_2 + \sigma_2]! [c]!}} 3\varphi_2 \left(\begin{matrix} q^{-2c}, q^{-2(J_2-\sigma'_2)}, q^{-2(J_2+\sigma_2)} \\ q^{2(\sigma'_2-\sigma_2+1)}, q^{2(2j(\lambda)+\sigma'_2-J_2-c+1)} \end{matrix} \middle| q^2, q^{2(J_2+2j(\lambda)+\sigma'_2+2)} \right)$$

$$q^{j(\lambda)(2J_2-2\sigma'_2)+c(-J_2-3\sigma_2/2+\sigma'_2/2)-\sigma_2\sigma'_2/2+3\sigma_2^2/4-3\sigma_2'^2/4+J_2\sigma'_2},$$

and

$$C_1(J_1, J_2, \sigma_2, \sigma'_2) = \frac{(-1)^{1/2\sigma_2+1/2\sigma'_2-J_2}}{(q - q^{-1})^{1/2\sigma_2+1/2\sigma'_2-J_2}} q^{j_2/2+(3\sigma_2)/4-(5\sigma'_2)/4+2J_1\sigma_2-J_2^2/2}.$$

Note that $C_1(J_1, J_2, \sigma_2, \sigma'_2)$ does not depend on $c, \sigma_1, \sigma'_1, \lambda$ and gives a constant in $B_{m,l}(u, z, \lambda)$ thus can be discarded in duality function.

Next, we convert back to particle–hole notation and also apply various identities (e.g.(1.5.2),(3.2.4),(3.2.5) in [17]) to simplify the results. For example, up to a constant,

$$\lim_{J_0 \rightarrow \infty} \langle \mu | B_{2J_0-c+N(\xi-\mu), 2J_0-c}(\infty, \lambda) | \xi \rangle$$

$$= \prod_{i=1}^N \mathcal{F}(J_i, j(\lambda) + N_{[1,i-1]}(\mu - J), c + N_{[1,i-1]}(\mu - \xi), \xi_i - J_i, \mu_i - J_i)$$

$$= \prod_{i=1}^N \frac{[2j(\lambda) - c + N_{[1,i-1]}(\mu + \xi - 2J)]! [\mu_i - \xi_i]!^{-1}}{(q^{4(j(\lambda)+1+N_{[1,i]}(\mu-J))}; q^2)_{2J_i-\mu_i} [2j(\lambda) - c + N_{[1,i-1]}(\mu + \xi - 2J) + \mu_i - 2J_i]!}$$

$$\sqrt{\frac{[\mu_i]! [2J_i - \xi_i]! [c + N_{[1,i]}(\mu - \xi)]!^{-1}}{[2J_i - \mu_i]! [\xi_i]! [c + N_{[1,i-1]}(\mu - \xi)]!}} q^{2J_i N_{[1,i-1]}(\mu) + J_i(\mu_i - \xi_i) - \xi_i N_{[1,i-1]}(\mu)}$$

$$\begin{aligned}
& 3\varphi_2 \left(\begin{matrix} q^{-2(c+N_{[1,i-1]}(\mu-\xi))}, q^{-2(2J_i-\mu_i)}, q^{-2\xi_i} \\ q^{2(\mu_i-\xi_i+1)}, q^{2(2j(\lambda)-c+N_{[1,i-1]}(\mu+\xi-2J)+\mu_i-2J_i+1)} \end{matrix}, q^2, q^{2(2(j(\lambda)+N_{[1,i-1]}(\mu-J))+\mu_i+2)} \right) \\
&= \prod_{i=1}^N \frac{[2j(\lambda) - c + N_{[1,i-1]}(\mu + \xi - 2J)]!}{\left(q^{4(j(\lambda)+1+N_{[1,i]}(\mu-J))}; q^2 \right)_{2J_i-\mu_i} [2j(\lambda) - c + N_{[1,i-1]}(\mu + \xi - 2J) + \mu_i - 2J_i]!} \\
&\frac{1}{\sqrt{[\mu_i]![2J_i - \xi_i]![2J_i - \mu_i]![\xi_i]!}} q^{2J_i N_{[1,i-1]}(\mu) + J_i(\mu_i + \xi_i) - \xi_i N_{[1,i]}(\mu)} \\
& 3\varphi_2 \left(\begin{matrix} q^{-2(2J_i-\mu_i)}, q^{-2\xi_i}, q^{2(2j(\lambda)+2N_{[1,i-1]}(\mu-J)+\mu_i-2J_i+1)} \\ q^{-4J_i}, q^{2(2j(\lambda)-c+N_{[1,i-1]}(\mu+\xi-2J)+\mu_i-2J_i+1)} \end{matrix}, q^2, q^2 \right) \\
&= \prod_{i=1}^N \frac{\left(q^{2(2j(\lambda)-c+N_{[1,i-1]}(\mu+\xi-2J)+\mu_i-2J_i+1)}; q^2 \right)_{2J_i-\mu_i}}{\left(q^{4(j(\lambda)+1+N_{[1,i]}(\mu-J))}; q^2 \right)_{2J_i-\mu_i} \sqrt{[\mu_i]![2J_i - \xi_i]![2J_i - \mu_i]![\xi_i]!}} \\
& 3\varphi_2 \left(\begin{matrix} q^{-2(2J_i-\mu_i)}, q^{-2\xi_i}, q^{2(2j(\lambda)+2N_{[1,i-1]}(\mu-J)+\mu_i-2J_i+1)} \\ q^{-4J_i}, q^{2(2j(\lambda)-c+N_{[1,i-1]}(\mu+\xi-2J)+\mu_i-2J_i+1)} \end{matrix}, q^2, q^2 \right) \\
& q^{2J_i N_{[1,i-1]}(\mu-\xi) + J_i(\mu_i + \xi_i) - 2\xi_i N_{[1,i]}(\mu)}.
\end{aligned}$$

The other limits can be derived in the same way.

Q.E.D.

4.6.2 Calculating The Ground State Transformation

We perform the calculation for (35). Note that G is not unique: we solve for one such that (23) is satisfied.

Proposition 8. Suppose $\dim V_i = 2J_i + 1$ where $i \geq 0$. Define up to a map $G : V_1 \otimes V_2 \rightarrow V_1 \otimes V_2$

$$\begin{aligned} & \langle \xi_1, \xi_2 | G(\lambda) | \xi_1, \xi_2 \rangle \\ &= \frac{([2J_1 - \xi_1]![2J_2 - \xi_2]![\xi_2]![\xi_1]!)^{-1/2}}{\left(q^{4(j(\lambda)+1+\xi_1-J_1)}; q^2\right)_{2J_1-\xi_1} \left(q^{4(j(\lambda)+1+\xi_1+\xi_2-J_1-J_2)}; q^2\right)_{2J_2-\xi_2}} q^{(J_1+2J_2)\xi_1+J_2\xi_2}. \end{aligned} \quad (37)$$

Then (23) is satisfied with the constant $C(J_1, J_2, \xi_1 + \xi_2) = q^{2(J_1+J_2)(J_1+J_2-\xi_1-\xi_2)}$.

Proof. It is clear that the ground state G does not depend on u , so we can set $u = \infty$. Take $l = 2J_0$ and also $J_0 \rightarrow \infty$ to find that

$$\begin{aligned} & \langle \xi_1, \xi_2 | G(\lambda) | \xi_1, \xi_2 \rangle \\ &= \lim_{J_0 \rightarrow \infty} \langle \xi | B_{2J_0-\xi_1-\xi_2, 2J_0}(\lambda) | 0, 0 \rangle \\ &= \frac{([2J_1 - \xi_1]![2J_2 - \xi_2]![\xi_2]![\xi_1]!)^{-1/2}}{\left(q^{4(j(\lambda)+1+\xi_1-J_1)}; q^2\right)_{2J_1-\xi_1} \left(q^{4(j(\lambda)+1+\xi_1+\xi_2-J_1-J_2)}; q^2\right)_{2J_2-\xi_2}} q^{(J_1+2J_2)\xi_1+J_2\xi_2}. \end{aligned}$$

Next, we have

$$\frac{\langle \xi_1, \xi_2 | \Pi G(-\mathbf{i}\infty) \Pi | \xi_1, \xi_2 \rangle}{\langle \xi_1, \xi_2 | P G(-\mathbf{i}\infty) P | \xi_1, \xi_2 \rangle} = q^{2(J_1+J_2)(J_1+J_2-\xi_1-\xi_2)}.$$

Thus we have $\langle \mu_1, \mu_2 | C | \xi_1, \xi_2 \rangle = \delta_{\mu_i=\xi_i} q^{2(J_1+J_2)(J_1+J_2-\xi_1-\xi_2)}$ in (23).

Q.E.D.

4.6.3 Proof of Corollary 1

This section provides explicit calculations for Corollary 1.

Recall that [17]

$${}_3\varphi_2 \left[\begin{matrix} q^{-n}, a, b \\ d, e \end{matrix} ; q, \frac{deq^n}{ab} \right] = \frac{(e/a; q)_n}{(e; q)_n} {}_3\phi_2 \left[\begin{matrix} q^{-n}, a, d/b \\ d, aq^{1-n}/e \end{matrix} ; q, q \right]$$

and the inversion formula

$$\begin{aligned} & {}_r\phi_s \left[\begin{matrix} a_1, \dots, a_r \\ b_1, \dots, b_s \end{matrix} ; q, z \right] \\ &= \sum_{n=0}^{\infty} \frac{(a_1^{-1}, \dots, a_r^{-1}; q^{-1})_n}{(q^{-1}, b_1^{-1}, \dots, b_s^{-1}; q^{-1})_n} \left(\frac{a_1 \cdots a_r z}{b_1 \cdots b_s q} \right)^n. \end{aligned}$$

Combining the above two identities we get

$${}_3\phi_2 \left[\begin{matrix} q^n, 1/a, 1/b \\ 1/d, 1/e \end{matrix} ; q^{-1}, q^{-1} \right] = \frac{(e/a; q)_n}{(e; q)_n} {}_3\phi_2 \left[\begin{matrix} q^{-n}, a, d/b \\ d, aq^{1-n}/e \end{matrix} ; q, q \right].$$

Taking e to ∞ yields

$${}_3\phi_2 \left[\begin{matrix} q^n, 1/a, 1/b \\ 1/d, 0 \end{matrix} ; q^{-1}, q^{-1} \right] = a^{-n} {}_3\phi_2 \left[\begin{matrix} q^{-n}, a, d/b \\ d, 0 \end{matrix} ; q, q \right]. \quad (38)$$

Recall that

$$\begin{aligned} & \lim_{c \rightarrow -\infty} \mathcal{D}_c(\mu, \xi) \\ &= \prod_{i=1}^N {}_3\phi_2 \left(\begin{matrix} q^{-2(2J_i - \mu_i)}, q^{-2\xi_i}, e^{-2\pi i \lambda} q^{2(2N_{[1, i-1]}(\mu - J) + \mu_i - 2J_i)} \\ q^{-4J_i}, 0 \end{matrix} ; q^2, q^2 \right) \\ & q^{-4J_i N_{[1, i-1]}(\xi) - 2\xi_i N_{[1, i]}(\mu)}. \end{aligned}$$

Applying (38) by taking $a = q^{-2(2J_i - \mu_i)}$, $n = \xi_i$, $d = q^{-4J_i}$ and $b^{-1} =$

$e^{-2\pi i\lambda} q^{2(2N_{[1,i-1]}(\mu-J)+\mu_i)}$, we obtain that

$$\begin{aligned} \lim_{c \rightarrow -\infty} \mathcal{D}_c(\mu, \xi) &= \prod_{i=1}^N {}_3\varphi_2 \left(\begin{matrix} q^{2(2J_i-\mu_i)}, q^{2\xi_i}, e^{-2\pi i\lambda} q^{2(2N_{[1,i-1]}(\mu-J)+\mu_i)} \\ q^{4J_i}, 0 \end{matrix} \middle| q^{-2}, q^{-2} \right) \\ &\quad q^{-2\xi_i(2J_i-\mu_i)-4J_iN_{[1,i-1]}(\xi)-2\xi_iN_{[1,i]}(\mu)} \\ &= \prod_{i=1}^N {}_3\varphi_2 \left(\begin{matrix} q^{2(2J_i-\mu_i)}, q^{2\xi_i}, e^{-2\pi i\lambda} q^{2(2N_{[1,i-1]}(\mu-J)+\mu_i)} \\ q^{4J_i}, 0 \end{matrix} \middle| q^{-2}, q^{-2} \right) \\ &\quad q^{-4J_iN_{[1,i]}(\xi)-2\xi_iN_{[1,i-1]}(\mu)}, \end{aligned}$$

which is equal to $\mathcal{D}_{ort}(\mu, \xi)$ up to a constant.

Q.E.D.

4.7 Proof of Proposition 3

In [1], the author computes explicit formulas for the stochastic weights of the dynamic stochastic higher spin vertex model by using fusion. Essentially, he proves a recursive relation for the weights and then solves the relation explicitly. Here, we prove that the same recursive relation is satisfied by our weights. Because the (difficult) task of solving the weights was already done, a proof of the recursive relation suffices to prove Proposition 3. While “spiritually” the fusion of Aggarwal is the same as the algebraic fusion that arises from the universal R -matrix and twister, for reasons of completeness a proof still needs to be provided.

The first step in matching is to match notation across different authors, which is best done with an example. For simplicity, we only consider the case when spectral $u = 0$ in (9) and only match the hypergeometric term. We will prove that for general u , the weights also match.

4.7.1 Matching of Notation

We match the notation with an example. First, note that

$$\begin{aligned}
& \lim_{u \rightarrow 0} \psi(i_1, j_1; i_2, j_2) \\
&= \mathbf{1}_{i_1+j_1=i_2+j_2} \left(\frac{q^{j_1/2-i_1j_1-2i_2j_1-i_1^2+i_2^2+j_1^2/2}}{s^{2j_1+2j_2}\kappa^{j_2}} \right) \frac{(q^{i_1-j_2+1}; q)_{j_2} (q^{j_2-J}; q)_{j_1} (s^2q^{i_1-j_2}; q)_{j_1}}{(q; q)_{j_2} (q^{j_2-J}; q)_{j_1-j_2}} \\
&\times \frac{(q\kappa^{-1}; q)_{j_1} (q^{j_2-2i_1+1}s^{-2}\kappa^{-1}; q)_{i_1-j_2}}{(q^{1-j_2}\kappa^{-1}; q)_{j_1} (q^{j_2-i_1-J+1}s^{-2}\kappa^{-1}; q)_{j_1} (q^{j_2-2i_1-J}s^{-2}\kappa^{-1}; q)_{j_2} (q^{2j_2-2i_1-J+1}s^{-2}\kappa^{-1}; q)_{i_1-j_2}} \\
&\times {}_8W_7(q^{-j_2}\kappa^{-1}; q^{-j_1}, q^{-j_2}, q^{j_1-J}\kappa^{-1}, q^{1-i_1}s^{-2}\kappa^{-1}, q^{-i_1}\kappa^{-1}; q, s^2\kappa q^{1+2i_1-j_2+J})
\end{aligned}$$

Now set $q^{2j(\lambda)+1} = e^{-2\pi i\lambda}$ and apply (2.5.1) and (3.2.1) in [17] to get

$$\begin{aligned}
& {}_8W_7(q^{-j_2}\kappa^{-1}; q^{-j_1}, q^{-j_2}, q^{j_1-J}\kappa^{-1}, q^{1-i_1}s^{-2}\kappa^{-1}, q^{-i_1}\kappa^{-1}; q, s^2\kappa q^{1+2i_1-j_2+J}) \\
&= \frac{(q^{J-i_1+i_2-3j_1-2j(\lambda)}, q^{i_2+2j(\lambda)+2})_{i_1-\Lambda-J+2j_1+2j(\lambda)} (q^{-j_1}, q^{J+\Lambda-i_2-j_1-2j(\lambda)+1})_{J+\Lambda-i_1-2j_1-2j(\lambda)}}{(q^{J-i_1+i_2-2j_1+1}, q^{i_2-j_1+1})_{i_1-\Lambda-J+2j_1+2j(\lambda)} (q^{J-2j_1-2j(\lambda)}, q^{\Lambda-i_2+1})_{J+\Lambda-i_1-2j_1-2j(\lambda)}} \\
&{}^4\varphi_3 \left(\begin{matrix} q^{J+\Lambda-i_1-2j_1-2j(\lambda)}, q^{J-j_1-2j(\lambda)}, q^{J+\Lambda-i_1-i_2-j_1-2j(\lambda)}, q^{\Lambda-i_2-2j(\lambda)} \\ q^{J+\Lambda-i_1-i_2-2j_1-4j(\lambda)-1}, q^{J+\Lambda-i_2-j_1-2j(\lambda)+1}, q^{J+\Lambda-i_1-j_1-2j(\lambda)+1} \end{matrix} \middle| q, q \right).
\end{aligned}$$

Now setting $q = Q^2$,

$$\begin{aligned}
& {}^4\varphi_3 \left(\begin{matrix} q^{J+\Lambda-i_1-2j_1-2j(\lambda)}, q^{J-j_1-2j(\lambda)}, q^{J+\Lambda-i_1-i_2-j_1-2j(\lambda)}, q^{\Lambda-i_2-2j(\lambda)} \\ q^{J+\Lambda-i_1-i_2-2j_1-4j(\lambda)-1}, q^{J+\Lambda-i_2-j_1-2j(\lambda)+1}, q^{J+\Lambda-i_1-j_1-2j(\lambda)+1} \end{matrix} \middle| q, q \right) \\
&= {}^4\varphi_3 \left(\begin{matrix} Q^{2(J+\Lambda-i_1-2j_1-2j(\lambda))}, Q^{2(J-j_1-2j(\lambda))}, Q^{2(J+\Lambda-i_1-i_2-j_1-2j(\lambda))}, Q^{2(\Lambda-i_2-2j(\lambda))} \\ Q^{2(J+\Lambda-i_1-i_2-2j_1-4j(\lambda)-1)}, Q^{2(J+\Lambda-i_2-j_1-2j(\lambda)+1)}, Q^{2(J+\Lambda-i_1-j_1-2j(\lambda)+1)} \end{matrix} \middle| Q^2, Q^2 \right)
\end{aligned}$$

On the other hand, applying a symmetry of $6j$ -symbol (see [22]) and

the equivalent definition through Racah–Wilson polynomial (4), we have

$$\begin{aligned}
& \langle i_1, j_1 | S^{\Lambda, J}(\infty, \lambda) | i_2, j_2 \rangle \sim \langle i_1, j_1 | R^{\Lambda, J}(\infty, \lambda) | i_2, j_2 \rangle \\
& \sim \left\{ \begin{array}{ccc} \Lambda/2 & j(\lambda) + j_1 + i_1 - J/2 - \Lambda/2 & j(\lambda) + j_1 - J/2 \\ J/2 & j(\lambda) & j(\lambda) + i_2 - \Lambda/2 \end{array} \right\}_Q \\
& \sim \left\{ \begin{array}{ccc} i_2/2 - \Lambda/2 + j_2/2 + j(\lambda) & i_1 - \Lambda/2 - J/2 + j_2 + j(\lambda) & J/2 + i_2/2 - j_2/2 \\ i_2/2 - J/2 + j_2/2 + j(\lambda) & j(\lambda) & \Lambda/2 - i_2/2 + j_2/2 \end{array} \right\}_Q \\
& \sim {}_4\varphi_3 \left(\begin{array}{c} Q^{2(J+\Lambda-i_1-2j_1-2j(\lambda))}, Q^{2(J-j_1-2j(\lambda))}, Q^{2(J+\Lambda-i_1-i_2-j_1-2j(\lambda))}, Q^{2(\Lambda-i_2-2j(\lambda))} \\ Q^{2(J+\Lambda-i_1-i_2-2j_1-4j(\lambda)-1)}, Q^{2(J+\Lambda-i_2-j_1-2j(\lambda)+1)}, Q^{2(J+\Lambda-i_1-j_1-2j(\lambda)+1)} \end{array} \middle| Q^2, Q^2 \right) \\
& \sim \lim_{u \rightarrow 0} \psi(i_1, j_1; i_2, j_2)
\end{aligned}$$

where $A \sim B$ means the hypergeometric term ${}_4\varphi_3$ are equal in both A and B , although the constants in front of the hypergeometric term may be different. Thus the matching of the notation is demonstrated.

4.7.2 Proof in the General Case:

Recall the dynamical Yang-Baxter Equation is:

$$R_{12}(u, \lambda - 2\eta H_3) R_{13}(uv, \lambda) R_{23}(v, \lambda - 2\eta H_1) = R_{23}(v, \lambda) R_{13}(uv, \lambda - 2\eta H_2) R_{12}(u, \lambda), \quad (39)$$

For each positive integer, let V^{2J} denote the $(2J + 1)$ -dimensional representation of $U_q(\mathfrak{sl}_2)$. Recall that there is a projection

$$V^1 \otimes V^{J-1} \rightarrow \mathcal{P}(V^1 \otimes V^{J-1}) \cong V^J,$$

where \mathcal{P} is the projection $\frac{1}{q-q^{-1}} \check{R}^{J-1,1}(q^2, \lambda)$. One can verify by direct calculation that $\mathcal{P}^2 = \mathcal{P}$ using the explicit formulas for the R -matrix. Let $\iota : V^J \rightarrow V^{J-1} \otimes V^1$ denote the inclusion.

Below, use the same notations as [1]. By the dynamic Yang–Baxter

Equation, where R_{13} is W_I and R_{23} is \widehat{W}_{J-1} , we see that

$$\begin{aligned} \langle j_2, i_2 | W_J | j_1, i_1 \rangle &= \sum_{k=0}^1 \sum_{l=0}^1 \sum_{m=0}^1 \langle j_2, i_2 | \mathcal{P} \otimes \text{id} | m, l, i_2 \rangle \langle m, l, i_2 | W_I | k, l, \delta \rangle \\ &\quad \times \langle k, l, \delta | \widehat{W}_{J-1} | k, j_1 - k, i_1 \rangle \langle k, j_1 - k, i_1 | \iota \otimes \text{id} | j_1, i_1 \rangle \end{aligned}$$

where $l + \delta = j_1 - k + i_1$. Note that this term is only nonzero if $j_2 = m + l$, so the summation only has four terms. In the above equation, the dependence on the parameters is dropped.

Having set $u = q^{2(J-1)}$ in (39) so that $\mathcal{P} = R_{12}$, the two terms are now

$$W_1(i_2, m; \delta, k | q^{2(J-1)}z, \lambda), \quad \widehat{W}_{J-1}(i_1, j_1 - k; \delta, l | z, \lambda - 2\eta H_1(k)).$$

We write δ in terms of i_2, m, k by

$$\delta = j_1 + i_1 - k - l = j_2 + i_2 - k - l = i_2 - k + m.$$

We can see that this matches with the four terms in the recurrence relation in [1];

$$\begin{aligned} &\widehat{W}_J(i_1, j_1; i_2, j_2 | v, \lambda) \\ &= \widehat{W}_{J-1}(i_1, j_1; i_2, j_2 | v, \lambda - 2\eta) W_1(i_2, 0; i_2, 0 | v + 2\eta(J-1), \lambda) \\ &\quad + \widehat{W}_{J-1}(i_1, j_1 - 1; i_2 - 1, j_2 | v, \lambda + 2\eta) W_1(i_2 - 1, 1; i_2, 0 | v + 2\eta(J-1), \lambda) \\ &\quad + \widehat{W}_{J-1}(i_1, j_1; i_2 + 1, j_2 - 1 | v, \lambda - 2\eta) W_1(i_2 + 1, 0; i_2, 1 | v + 2\eta(J-1), \lambda) \\ &\quad + \widehat{W}_{J-1}(i_1, j_1 - 1; i_2, j_2 - 1 | v, \lambda + 2\eta) W_1(i_2, 1; i_2, 1 | v + 2\eta(J-1), \lambda) \end{aligned}$$

where $e^v = z$. Therefore the weights match and Proposition 3 is proved.

Q.E.D.

5 Orthogonal Polynomial Duality

This section will discuss how the algebraic approach in this paper can

be used to prove orthogonal polynomial duality. We provide a new proof of a previous orthogonal polynomial duality in the nondynamic case[16]. It is not clear how to recover the dynamic dualities of [19].

5.1 The dual quantum group

In order to produce *orthogonal polynomial duality*, we will use the Hopf $*$ -algebra on the dual quantum group. This is because the duality functionals are constructed from elements of the dual quantum group, and the $*$ -structure leads to orthogonality in the duality function. The definitions here are taken from [23] as well.

Recall that for a Hopf algebra A over a field \mathbf{k} , the dual Hopf algebra A^* is the Hopf subalgebra of $\text{Hom}_{\mathbf{k}}(A, \mathbf{k})$ consisting of the matrix elements of finite-dimensional A -modules. The right and left regular representation of A on A^* are given by

$$\begin{aligned}\mathcal{R}(a) &: f \mapsto \langle \text{id} \otimes a, \Delta(f) \rangle \\ \mathcal{L}(a) &: f \mapsto \langle S^{-1}(a) \otimes \text{id}, \Delta(f) \rangle.\end{aligned}$$

In the case of $U_q(\mathfrak{g})$, where q is a nonzero real number, \mathbf{k} is the field of complex numbers, and \mathfrak{g} is a finite-dimensional semisimple Lie algebra associated to a Lie group G , the dual Hopf algebra is denoted $\mathbf{C}[G]_q$ and called “the quantized algebra of functions.” The finite-dimensional modules of $U_q(\mathfrak{g})$ will be denoted by $L(\Lambda)$, where $\Lambda \in P_+$, where P_+ is the set of all dominant weights. A linear basis of $\mathbf{C}[G]_q$ is given by a quantized version of the Peter–Weyl theorem, which states that the action of $\mathcal{L} \otimes \mathcal{R}$ defines an isomorphism

$$\mathbf{C}[G]_q \cong \bigoplus_{\Lambda \in P_+} L(\Lambda)^* \otimes L(\Lambda)$$

where $c_{l,v}^{L(\Lambda)}$ is mapped to $l \otimes v$.

The $*$ -structure on $\mathbb{C}[G]_q$ can be defined on the basis elements $c_{l,v}^{L(\Lambda)}$. Fix an orthonormal basis $\{v_\lambda^{(i)}\}$ where λ ranges over the set of weights of $L(\Lambda)$, and each $v_\lambda^{(i)} \in L(\Lambda)_\lambda$, where i takes values in $1, 2, \dots, \dim L(\Lambda)_\lambda$. Let $\{l_{-\lambda}^{(i)}\}$ denote the dual basis in $L(\Lambda)^*$, and let $c_{-\mu,i,\lambda,j}^\Lambda$ denote the matrix element (with respect to these bases) corresponding to the pair $(l_{-\mu}^{(i)}, v_\lambda^{(j)})$. Then

$$(c_{-\mu,i,\lambda,j}^\Lambda)^* = q^{2(\mu-\lambda,\rho)} c_{\mu,i,-\lambda,j}^{w_0\Lambda}$$

where ρ is half the sum of the positive roots and w_0 is the element of the Weyl group of maximal length.

5.2 Proof of Orthogonal Polynomial Duality

In [16], the authors proved an orthogonal polynomial duality for a so-called “multi-species partial exclusion process”. Here we demonstrate how the method of the dual Hopf algebra can prove the same result.

Recall that the Hopf $*$ -algebra structure is given by

$$(c_{-\mu,i,\lambda,j}^\Lambda)^* = q^{2(\mu-\lambda,\rho)} c_{\mu,i,-\lambda,j}^{w_0\Lambda}$$

We will also use Proposition 2.1.4 of Chapter 3 of [23], which states:

Let $\Lambda, \Lambda' \in P_+$ and $l \in L(\Lambda)_{-\lambda}^*$, $l' \in L(\Lambda')_{-\lambda'}$. Suppose that $v' \in L(\Lambda')_\gamma$ and that $v_\Lambda \in L(\Lambda)_\Lambda$ is a highest weight vector. Then there exist $a_\nu \in \mathbb{C}$, $l_\nu \in ((U_q \mathfrak{b}_+)l)_{-\lambda+\mu}$, and $l'_\nu \in ((U_q \mathfrak{b}_-)l')_{-\lambda'-\nu}$ such that

$$c_{l',v'}^{\Lambda'} c_{l,v_\Lambda}^\Lambda = q^{2((\Lambda,\gamma)-(\lambda,\lambda'))} c_{l,v_\Lambda}^\Lambda c_{l',v'}^{\Lambda'} + q^{2(\Lambda,\gamma)} \sum_{\nu \in Q_+} a_\nu c_{l_\nu,v_\Lambda}^\Lambda c_{l'_\nu,v'}^{\Lambda'}$$

where Q_+ is the lattice of positive roots of \mathfrak{g} .

In [23], the authors choose certain weights for l', l, v' such that the sum on the right-hand-side is empty. For our purposes, we will take

l' to be a lowest-weight vector, which means that l'_ν can only equal 0, resulting in the sum being empty. In order to relate this relation to the $*$ -structure, we will also take v' to be a lowest weight vector and l to be a highest weight vector. Then for these choices, $-\gamma = \lambda' = \Lambda'$ and $\lambda = -\Lambda$ so that

$$c_{-\Lambda', -\Lambda'}^{\Lambda'} c_{\Lambda, \Lambda}^{\Lambda} = c_{\Lambda, \Lambda}^{\Lambda} c_{-\Lambda', -\Lambda'}^{\Lambda'}.$$

Furthermore, for sl_2 the weight lattice is one dimensional and the Weyl group is trivial, so the $*$ -algebra structure implies

$$(c_{-\Lambda, i, -\Lambda, j}^{\Lambda})^* = c_{\Lambda, i, \Lambda, j}^{\Lambda}.$$

and setting $\Lambda' = -\Lambda$ then

$$[(c_{-\Lambda, i, -\Lambda, j}^{\Lambda}), c_{\Lambda, i, \Lambda, j}^{\Lambda}] = 0.$$

Recall from linear algebra that the exponential of a skew-Hermitian matrix is unitary. However, in the non-commutative setting, however, we will need the q -exponential

$$e_q(z) = \sum_{n=0}^{\infty} \frac{z^n}{(q; q)_n} = \frac{1}{(z; q)_{\infty}}, \quad |z| < 1, \quad \text{and} \quad \mathcal{E}_q(z) = \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}} z^n}{(q; q)_n} = (-z; q)_{\infty}$$

If $XY = qYX$ then

$$\mathcal{E}_q(X + Y) = \mathcal{E}_q(X)\mathcal{E}_q(Y), \quad e_q(Y + X) = e_q(Y)e_q(X)$$

The two q -exponentials are inverses in the sense that $e_q(Z)\mathcal{E}_q(-Z) = 1$ for any Z .

Lemma 2. *Now let X be an arbitrary element such that $X^*X = Q^{-1}XX^*$. Then*

$$U(X) = \mathcal{E}_Q(X^*)e_Q(-X)\mathcal{E}_Q(-X)e_Q(X^*)$$

is unitary in the sense that $U(X)U(X^*) = 1$.

Proof. Note that in the case $q = 1$, this becomes the statement that $\exp(2(X^* - X))$ is unitary. Indeed,

$$U(X)U(X)^* = \mathcal{E}_Q(X^*)e_Q(-X)\mathcal{E}_Q(-X)e_Q(X^*)e_Q(X)\mathcal{E}_Q(-X^*)e_Q(-X^*)\mathcal{E}_Q(X).$$

Once we show that $e_Q(X)$ and $e_Q(X^*)$ commute, then the inversion formula implies this equals 1. To prove the commutation, use the q -Baker–Campbell–Hausdorff formula:

$$\begin{aligned} e_Q(\lambda X)Y\mathcal{E}_Q(-\lambda X) &= \sum_{n=0}^{\infty} \frac{\lambda^n}{(q; q)_n} [X, Y]'_n \\ [X, Y]'_0 &= Y, \quad [X, Y]'_{n+1} = X[X, Y]'_n - q^n [X, Y]'_n X, \quad n = 0, 1, 2, \dots \end{aligned}$$

So if $Q = q^{A(\Lambda, \Lambda)}$ then

$$e_Q(\lambda X)X^*\mathcal{E}_Q(-\lambda X) = X^*$$

because

$$[X, X^*]'_1 = XX^* - QX^*X = 0.$$

Therefore

$$e_Q(\lambda X)e_Q(X^*)\mathcal{E}_Q(-\lambda X) = e_Q(X^*),$$

which implies that

$$e_Q(\lambda X)e_Q(-X^*) = e_Q(-X^*)e_Q(\lambda X).$$

Q.E.D.

Recall that $X^*X = XX^*$ for $X = c_{\Lambda, i, \Lambda, j}^{\Lambda}$. The unitary symmetry is then

$$\exp(X - X^*) = \sum_{k_1=0}^{\infty} \sum_{k_2=0}^{\infty} \frac{X^{k_1} (-1)^{k_2} (X^*)^{k_2}}{k_1! k_2!}.$$

We now clarify the relationship between $c_{\pm\Lambda, i, \pm\Lambda, j}^{\Lambda}$ and the $B_{\mathbf{ml}}$ previously

calculated. Recall that

$$\langle \mu | B_{m,l}(u, z, \lambda) | \xi \rangle = \langle m, \mu | \mathcal{B}(u, z, \lambda) | l, \xi \rangle.$$

An equivalent definition is that

$$B_{ml} = \langle m | a_1 \dots a_N | l \rangle$$

where a_1, \dots, a_N are quantum group elements uniquely defined by ξ and μ . Thus, B_{ml} defines a linear map on $U_q(sl_2)$, which is a matrix element of a finite dimensional module (in particular, the (l, m) -entry of the spin J representation).

More precisely, for $0 \leq l, m \leq 2J$ and $\Lambda = J$

$$B_{ml} = c_{m-J, l-J}^J.$$

So setting $J = \Lambda$ and $m = l = 0$, we get that

$$B_{00} = c_{-\Lambda, -\Lambda}^\Lambda.$$

If $J = \Lambda$ and $m = l = 2J$ then

$$B_{2J, 2J} = c_{\Lambda, \Lambda}^\Lambda$$

Then each summand in the unitary symmetry is then

$$\frac{(-1)^{k_2}}{k_1! k_2!} (c_{\Lambda\Lambda}^\Lambda)_1^{k_1} (c_{-\Lambda, -\Lambda}^\Lambda)_2^{k_2},$$

where

$$\begin{aligned} & (c_{\Lambda\Lambda}^\Lambda)^{k_1} (c_{-\Lambda, -\Lambda}^\Lambda)^{k_2} \\ &= B_{00}(-i\infty) \cdots B_{00}(-i\infty) B_{00}(-i\infty) \\ & \quad \times B_{2J, 2J}(-i\infty) \cdots B_{2J, 2J}(-i\infty) B_{2J, 2J}(-i\infty). \end{aligned}$$

From here, we can compute an orthogonal polynomial duality for the (non–dynamic) stochastic higher spin vertex model. However, we note that the calculation would be same calculation as the inner–product method [11]. The inner–product method does not generalize as well as an algebraic method (see the discussion in [16], for example), so therefore we do not repeat the argument. The most natural algebraic approach to orthogonal polynomial duality would involve the twisted algebra of functions (see section 7 of chapter 3 of [23], which is best suited for a future paper.

Additionally, orthogonal polynomial dualities are usually taken to be orthogonal with respect to the reversible measures of a process. The stochastic vertex model, being totally asymmetric, is not reversible. However, in the non–dynamic stochastic vertex model, the duality functions are orthogonal with respect to the $ASEP(q, j)$ [12]. A recent preprint [19] constructed a dynamic version of $ASEP(q, j)$ with duality. Since the $ASEP(q, j)$ is not integrable, different methods would be needed. This is another reason orthogonality (from an algebraic perspective) is best suited for a future paper.

6 Appendix

6.1 Examples of Matching

6.1.1 The spin 1/2 case

$$G = \text{diag}(1, q^2 - e^{-2\pi i \lambda}, q(1 - e^{-2\pi i \lambda}), 1)$$

$$R = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \frac{q(z-1)}{zq^2-1} & \frac{(q^2-1)(z-e^{-2\pi i \lambda})}{(zq^2-1)(1-e^{-2\pi i \lambda})} & 0 \\ 0 & \frac{(q^2-1)(1-e^{-2\pi i \lambda}z)}{(zq^2-1)(1-e^{-2\pi i \lambda})} & \frac{(z-1)(e^{-4\pi i \lambda}q^2 - e^{-2\pi i \lambda}q^4 - e^{-2\pi i \lambda} + q^2)}{q(zq^2-1)(1-e^{-2\pi i \lambda})^2} & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$S = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \frac{(z-1)(1-e^{-2\pi i\lambda}q^2)}{(zq^2-1)(1-e^{-2\pi i\lambda})} & \frac{(q^2-1)(z-e^{-2\pi i\lambda})}{(zq^2-1)(1-e^{-2\pi i\lambda})} & 0 \\ 0 & \frac{(q^2-1)(1-e^{-2\pi i\lambda}z)}{(zq^2-1)(1-e^{-2\pi i\lambda})} & \frac{(q^2-e^{-2\pi i\lambda})(z-1)}{(zq^2-1)(1-e^{-2\pi i\lambda})} & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

To compare S to (8), let $q^{\frac{1}{2}}w := z^{-1}$, and $q \rightarrow q^2$ so that

$$S = \begin{bmatrix} \text{weight}(\odot) & 0 & 0 & 0 \\ 0 & \text{weight}(\text{--}) & \text{weight}(\lrcorner) & 0 \\ 0 & \text{weight}(\ulcorner) & \text{weight}(|) & 0 \\ 0 & 0 & 0 & \text{weight}(+) \end{bmatrix}$$

6.1.2 The spin 1 case

$$\psi(0, 0; 0, 0) = 1$$

$$\psi(0, 1; 0, 1) = \frac{(qe^{-2\pi i\lambda} - 1)(qu - 1)}{(-e^{-2\pi i\lambda} + q)(q - u)}$$

$$\psi(0, 1; 1, 0) = -\frac{(ue^{-2\pi i\lambda} - 1)(q^2 - 1)}{(-e^{-2\pi i\lambda} + q)(q - u)}$$

$$\psi(0, 2; 0, 2) = \frac{(uq^2 - 1)(qe^{-2\pi i\lambda} - 1)(q^2e^{-2\pi i\lambda} - 1)(qu - 1)}{q^2(-e^{-2\pi i\lambda} + q)(e^{-2\pi i\lambda} - 1)(q - u)(u - 1)}$$

$$\psi(0, 2; 1, 1) = \frac{(q^2e^{-2\pi i\lambda} - 1)(que^{-2\pi i\lambda} - 1)(qu - 1)(q - 1)(q + 1)^2}{q^2(q - u)(u - 1)(-q^2e^{-2\pi i\lambda} + qx^4 + q - e^{-2\pi i\lambda})}$$

$$\psi(0, 2; 2, 0) = -\frac{(q^2 - 1)(que^{-2\pi i\lambda} - 1)(q - 1)(uq^2e^{-2\pi i\lambda} - 1)}{q^2(qe^{-2\pi i\lambda} - 1)(e^{-2\pi i\lambda} - 1)(q - u)(u - 1)}$$

$$\psi(1, 0; 1, 0) = \frac{(q^2 - 1)(uq^2 - e^{-2\pi i\lambda})}{q(-e^{-2\pi i\lambda} + q)(q - u)}$$

$$\psi(1, 0; 0, 1) = -\frac{(q^3 - e^{-2\pi i\lambda})(qu - 1)}{q(-e^{-2\pi i\lambda} + q)(q - u)}$$

$$\psi(1, 1; 0, 2) = \frac{(qe^{-2\pi i\lambda} - 1)(qu - 1)(q - 1)(-e^{-2\pi i\lambda} + qu)}{q(-e^{-2\pi i\lambda} + q)(e^{-2\pi i\lambda} - 1)(q - u)(u - 1)}$$

$$\psi(1, 1; 1, 1) = -\left(\frac{(-e^{-2\pi i\lambda} + u)(q^2 - 1)(qe^{-2\pi i\lambda} - 1)^2(q^2 - e^{-2\pi i\lambda})(qu - 1)(u - 1)}{(-e^{-2\pi i\lambda} + q)(-e^{-2\pi i\lambda} + u)(q^2 - 1)(que^{-2\pi i\lambda} - 1)(q - 1)} - 1 \right) \\ \times \frac{(que^{-2\pi i\lambda} - 1)(q - 1)}{q(qe^{-2\pi i\lambda} - 1)(e^{-2\pi i\lambda} - 1)(q - u)(u - 1)}$$

$$\psi(1, 1; 2, 0) = -\frac{(-e^{-2\pi i\lambda} + q)(que^{-2\pi i\lambda} - 1)(qu - 1)(q - 1)}{q(qe^{-2\pi i\lambda} - 1)(e^{-2\pi i\lambda} - 1)(q - u)(u - 1)}$$

$$\psi(1, 2; 1, 2) = -\frac{(q^3e^{-2\pi i\lambda} - 1)(qu - 1)}{q(qe^{-2\pi i\lambda} - 1)(q - u)}$$

$$\psi(1, 2; 2, 1) = -\frac{(1/q - 1)(q^2 - 1)(uq^2e^{-2\pi i\lambda} - 1)}{(qe^{-2\pi i\lambda} - 1)(q - u)(q - 1)}$$

$$\begin{aligned}
\psi(2, 0; 0, 2) &= \frac{(q^2 - 1)(uq^2 - e^{-2\pi i\lambda})(q - 1)(-e^{-2\pi i\lambda} + qu)}{q^2(-e^{-2\pi i\lambda} + q)(e^{-2\pi i\lambda} - 1)(q - u)(u - 1)} \\
\psi(2, 0; 1, 1) &= -\frac{(q^2 - 1)^2(q^2 - e^{-2\pi i\lambda})(qu - 1)(-e^{-2\pi i\lambda} + qu)}{q^2(-e^{-2\pi i\lambda} + q)(qe^{-2\pi i\lambda} - 1)(q - u)(q - 1)(u - 1)} \\
\psi(2, 0; 2, 0) &= -\frac{(uq^2 - 1)(-e^{-2\pi i\lambda} + q)(q^2 - e^{-2\pi i\lambda})(qu - 1)}{q^2(qe^{-2\pi i\lambda} - 1)(e^{-2\pi i\lambda} - 1)(q - u)(u - 1)} \\
\psi(2, 1; 1, 2) &= -\frac{(-e^{-2\pi i\lambda} + u)(q^2 - 1)}{(qe^{-2\pi i\lambda} - 1)(q - u)} \\
\psi(2, 1; 2, 1) &= \frac{(-e^{-2\pi i\lambda} + q)(qu - 1)}{(qe^{-2\pi i\lambda} - 1)(q - u)} \\
\psi(2, 2; 2, 2) &= 1
\end{aligned}$$

Let $q \rightarrow q^2$ and $u \rightarrow \frac{1}{zq^2}$, $\psi(i_1, j_1, i_2, j_2) = S(i_1, j_1, i_2, j_2)$. Then

$$\begin{aligned}
S(0, 0; 0, 0) &= 1 \\
S(0, 1; 0, 1) &= -\frac{q^2(q^2e^{-2\pi i\lambda} - 1)(z - 1)}{(zq^4 - 1)(q^2 - e^{-2\pi i\lambda})} \\
S(0, 1; 1, 0) &= \frac{(q^4 - 1)(zq^2 - e^{-2\pi i\lambda})}{(zq^4 - 1)(q^2 - e^{-2\pi i\lambda})} \\
S(0, 2; 0, 2) &= \frac{(-q^2 + z)(z - 1)(-q^6x^4 + q^4e^{-2\pi i\lambda} + q^2e^{-2\pi i\lambda} - 1)}{(e^{-2\pi i\lambda} - 1)(q^2 - e^{-2\pi i\lambda})(q^6z^2 - q^4z - q^2z + 1)} \\
S(0, 2; 1, 1) &= \frac{(-e^{-2\pi i\lambda} + z)(q^2 + 1)(q^4 - 1)(q^4e^{-2\pi i\lambda} - 1)(z - 1)}{(q^2e^{-2\pi i\lambda} - 1)(q^2 - e^{-2\pi i\lambda})(q^6z^2 - q^4z - q^2z + 1)} \\
S(0, 2; 2, 0) &= \frac{(-e^{-2\pi i\lambda} + z)(q^2 - 1)^2(q^2 + 1)(-q^2e^{-2\pi i\lambda} + z)}{(e^{-2\pi i\lambda} - 1)(q^2e^{-2\pi i\lambda} - 1)(q^6z^2 - q^4z - q^2z + 1)}
\end{aligned}$$

$$S(1, 0; 1, 0) = -\frac{(q^4 - 1)(-q^2 + ze^{-2\pi i\lambda})}{(zq^4 - 1)(q^2 - e^{-2\pi i\lambda})}$$

$$S(1, 0; 0, 1) = \frac{(q^6 - e^{-2\pi i\lambda})(z - 1)}{(zq^4 - 1)(q^2 - e^{-2\pi i\lambda})}$$

$$S(1, 1; 0, 2) = \frac{q^2(ze^{-2\pi i\lambda} - 1)(q^4 - 1)(z - 1)(-q^6x^4 + q^4e^{-2\pi i\lambda} + q^2e^{-2\pi i\lambda} - 1)}{(q^2 + 1)(e^{-2\pi i\lambda} - 1)(q^2 - e^{-2\pi i\lambda})(q^4e^{-2\pi i\lambda} - 1)(q^6z^2 - q^4z - q^2z + 1)}$$

$$S(1, 1; 1, 1) = -\frac{q^{10}e^{-2\pi i\lambda}z^2 - q^8x^4z + q^8e^{-2\pi i\lambda}z^2 - 3q^8e^{-2\pi i\lambda}z + q^8e^{-2\pi i\lambda} - q^8z - q^6x^4z^2}{(-q^4e^{-2\pi i\lambda} + q^2x^4 + q^2 - e^{-2\pi i\lambda})(q^6z^2 - q^4z - q^2z + 1)}$$

$$-\frac{2q^6x^4z - 2q^6e^{-2\pi i\lambda}z^2 + q^6e^{-2\pi i\lambda}z + q^6e^{-2\pi i\lambda} - q^6z^2 + 2q^6z + 2q^4x^4z - q^4x^4 + q^4e^{-2\pi i\lambda}z^2}{(-q^4e^{-2\pi i\lambda} + q^2x^4 + q^2 - e^{-2\pi i\lambda})(q^6z^2 - q^4z - q^2z + 1)}$$

$$-\frac{q^4e^{-2\pi i\lambda}z - 2q^4e^{-2\pi i\lambda} + 2q^4z - q^4 - q^2x^4z + q^2e^{-2\pi i\lambda}z^2 - 3q^2e^{-2\pi i\lambda}z + q^2e^{-2\pi i\lambda} - q^2z + e^{-2\pi i\lambda}}{(-q^4e^{-2\pi i\lambda} + q^2x^4 + q^2 - e^{-2\pi i\lambda})(q^6z^2 - q^4z - q^2z + 1)}$$

$$S(1, 1; 2, 0) = \frac{q^2(-e^{-2\pi i\lambda} + z)(q^2 - 1)(q^2 - e^{-2\pi i\lambda})(z - 1)}{(e^{-2\pi i\lambda} - 1)(q^2e^{-2\pi i\lambda} - 1)(q^6z^2 - q^4z - q^2z + 1)}$$

$$S(1, 2; 1, 2) = \frac{(q^6e^{-2\pi i\lambda} - 1)(z - 1)}{(zq^4 - 1)(q^2e^{-2\pi i\lambda} - 1)}$$

$$S(1, 2; 2, 1) = -\frac{(q^4 - 1)(-q^2e^{-2\pi i\lambda} + z)}{(zq^4 - 1)(q^2e^{-2\pi i\lambda} - 1)}$$

$$S(2, 0; 0, 2) = -\frac{(ze^{-2\pi i\lambda} - 1)(q^2 - 1)^2(q^2 + 1)(-q^2 + ze^{-2\pi i\lambda})}{(e^{-2\pi i\lambda} - 1)(q^2 - e^{-2\pi i\lambda})(q^6z^2 - q^4z - q^2z + 1)}$$

$$S(2, 0; 1, 1) = -\frac{(ze^{-2\pi i\lambda} - 1)(q^2 - 1)(q^2 + 1)^2(q^4 - e^{-2\pi i\lambda})(z - 1)}{(-q^4e^{-2\pi i\lambda} + q^2x^4 + q^2 - e^{-2\pi i\lambda})(q^6z^2 - q^4z - q^2z + 1)}$$

$$S(2, 0; 2, 0) = -\frac{(-q^2 + z)(z - 1)(-q^6 + q^4e^{-2\pi i\lambda} + q^2e^{-2\pi i\lambda} - x^4)}{(e^{-2\pi i\lambda} - 1)(q^2e^{-2\pi i\lambda} - 1)(q^6z^2 - q^4z - q^2z + 1)}$$

$$S(2, 1; 1, 2) = \frac{(q^4 - 1)(zq^2e^{-2\pi i\lambda} - 1)}{(zq^4 - 1)(q^2e^{-2\pi i\lambda} - 1)}$$

$$S(2, 1; 2, 1) = -\frac{q^2(q^2 - e^{-2\pi i\lambda})(z - 1)}{(zq^4 - 1)(q^2e^{-2\pi i\lambda} - 1)}$$

$$S(2, 2; 2, 2) = 1$$

6.2 Computer Simulations

A previous arXiv version contained the follow images of computer simulations; however, it was pointed out to the first author (who created the simulations) by multiple readers that the histograms was not overlayed with the probability distribution function of the Tracy–Widom distribution.

In the current version, we elaborate on the meaning of the histograms.




Figure 2: The left image shows 4000 simulations of the stochastic–six vertex model. The right figure simulates the dynamic stochastic–six vertex model with the same values.

The histograms on the left was created from simulations for the values $b_1 = 0.55$, $b_2 = 0.5$ and $e^{2\pi i\lambda} = 1.05$ and show the height function at the point $(1000, 1000)$. The histogram on the right was created with the same values except that $e^{2\pi i\lambda} = 0$. This figure shows that even at the time value of 1000, the distributions are visually indistinguishable.

Running the simulation long enough to obtain a reasonable overlay with the Tracy–Widom distribution. With some experimentation, it takes approximately 400,000 time increments to obtain a visually appealing figure. Each simulation required approximately 80 hours to complete on the Texas A&M High Performance Resource Center, using the FASTER cluster. The dynamic version of the simulation would require approximately two years to complete, possibly due to inefficiencies in the first author’s program. The first author hopes to improve the computer program at a later time.




Figure 3: The height function of the stochastic six vertex model at $(400000, 400000)$ when $b_1 = 0.75, b_2 = 0.65$. This shows 48 simulations.

References

- [1] Amol Aggarwal. Dynamical stochastic higher spin vertex models. *Selecta Math. (N.S.)*, 24(3):2659–2735, 2018.
- [2] Richard Askey and James Wilson. A set of orthogonal polynomials that generalize the Racah coefficients or $6-j$ symbols. *SIAM J. Math. Anal.*, 10(5):1008–1016, 1979.
- [3] Mario Ayala, Gioia Carinci, and Frank Redig. Quantitative Boltzmann-Gibbs principles via orthogonal polynomial duality. *J. Stat. Phys.*, 171(6):980–999, 2018.
- [4] Mario Ayala, Gioia Carinci, and Frank Redig. Higher order fluctuation fields and orthogonal duality polynomials. *Electron. J. Probab.*, 26:Paper No. 27, 35, 2021.
- [5] O. Babelon, D. Bernard, and E. Billey. A quasi-Hopf algebra interpretation of quantum $3-j$ and $6-j$ symbols and difference equations. *Phys. Lett. B*, 375(1-4):89–97, 1996.

- [6] Alexei Borodin. Symmetric elliptic functions, IRF models, and dynamic exclusion processes. *J. Eur. Math. Soc. (JEMS)*, 22(5):1353–1421, 2020.
- [7] Alexei Borodin and Ivan Corwin. Dynamic ASEP, duality, and continuous q^{-1} -Hermite polynomials. *Int. Math. Res. Not. IMRN*, (3):641–668, 2020.
- [8] Alexei Borodin, Ivan Corwin, and Vadim Gorin. Stochastic six-vertex model. *Duke Math. J.*, 165(3):563–624, 2016.
- [9] Alexei Borodin, Ivan Corwin, and Tomohiro Sasamoto. From duality to determinants for q -TASEP and ASEP. *Ann. Probab.*, 42(6):2314–2382, 2014.
- [10] Gioia Carinci, Chiara Franceschini, Cristian Giardinà, Wolter Groenevelt, and Frank Redig. Orthogonal dualities of Markov processes and unitary symmetries. *SIGMA Symmetry Integrability Geom. Methods Appl.*, 15:Paper No. 053, 27, 2019.
- [11] Gioia Carinci, Chiara Franceschini, and Wolter Groenevelt. q -Orthogonal dualities for asymmetric particle systems. *Electron. J. Probab.*, 26:Paper No. 108, 38, 2021.
- [12] Gioia Carinci, Cristian Giardinà, Frank Redig, and Tomohiro Sasamoto. A generalized asymmetric exclusion process with $U_q(\mathfrak{sl}_2)$ stochastic duality. *Probab. Theory Related Fields*, 166(3-4):887–933, 2016.
- [13] Ivan Corwin, Promit Ghosal, and Konstantin Matetski. Stochastic PDE limit of the dynamic ASEP. *Comm. Math. Phys.*, 380(3):1025–1089, 2020.

- [14] Eugène Cremmer, Jean-Loup Gervais, and Jean-François Roussel. The quantum group structure of 2D gravity and minimal models. II. The genus-zero chiral bootstrap. *Comm. Math. Phys.*, 161(3):597–630, 1994.
- [15] Giovanni Felder and Alexander Varchenko. On representations of the elliptic quantum group $E_{\tau,\eta}(\mathfrak{sl}_2)$. *Comm. Math. Phys.*, 181(3):741–761, 1996.
- [16] Chiara Franceschini, Jeffrey Kuan, and Zhengye Zhou. Orthogonal polynomial duality and unitary symmetries of multi-species $\text{asep}(q, \theta)$ and higher-spin vertex models via $*$ -bialgebra structure of higher rank quantum groups, 2022.
- [17] George Gasper and Mizan Rahman. *Basic hypergeometric series*, volume 96 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, second edition, 2004. With a foreword by Richard Askey.
- [18] Wolter Groenevelt. Orthogonal stochastic duality functions from Lie algebra representations. *J. Stat. Phys.*, 174(1):97–119, 2019.
- [19] Wolter Groenevelt and Carel Wagenaar. A generalized dynamic asymmetric exclusion process: Orthogonal dualities and degenerations, 2023.
- [20] Sabine Jansen and Noemi Kurt. On the notion(s) of duality for Markov processes. *Probab. Surv.*, 11:59–120, 2014.
- [21] M. Jimbo, H. Konno, S. Odake, and J. Shiraishi. Quasi-Hopf twistors for elliptic quantum groups. *Transform. Groups*, 4(4):303–327, 1999.

- [22] A. N. Kirillov and N. Yu. Reshetikhin. Representations of the algebra $U_q(\mathfrak{sl}(2))$, q -orthogonal polynomials and invariants of links. In *Infinite-dimensional Lie algebras and groups (Luminy-Marseille, 1988)*, volume 7 of *Adv. Ser. Math. Phys.*, pages 285–339. World Sci. Publ., Teaneck, NJ, 1989.
- [23] Leonid I. Korogodski and Yan S. Soibelman. *Algebras of functions on quantum groups. Part I*, volume 56 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1998.
- [24] J. Kuan. Algebraic symmetry and self-duality of an open asep. *Math Phys Anal Geom*, 24(12), 2021.
- [25] Jeffrey Kuan. An algebraic construction of duality functions for the stochastic $U_q(A_n^{(1)})$ vertex model and its degenerations. *Comm. Math. Phys.*, 359(1):121–187, 2018.
- [26] Jeffrey Kuan. A multi-species ASEP(q, j) and q -TAZRP with stochastic duality. *Int. Math. Res. Not. IMRN*, (17):5378–5416, 2018.
- [27] Jeffrey Kuan. Stochastic fusion of interacting particle systems and duality functions, 2019.
- [28] Jeffrey Kuan. A short note on Markov duality in multi-species higher spin stochastic vertex models. *Electronic Communications in Probability*, 26(none):1 – 11, 2021.
- [29] Jeffrey Kuan. Two Dualities: Markov and Schur–Weyl. *International Mathematics Research Notices*, 2022(13):9633–9662, 02 2021.
- [30] Jeffrey Kuan and Zhengye Zhou. Asymptotics of two-point correlations in the multi-species q -tazrp, 2023.

- [31] A. Kuniba, V. V. Mangazeev, S. Maruyama, and M. Okado. Stochastic R matrix for $U_q(A_n^{(1)})$. *Nuclear Phys. B*, 913:248–277, 2016.
- [32] Yier Lin. Markov duality for stochastic six vertex model. *Electronic Communications in Probability*, 24(none):1 – 17, 2019.
- [33] William Mead. Private communication to the first author in Eindhoven, December 2022.
- [34] Gunter Schütz and Sven Sandow. Non-abelian symmetries of stochastic processes: Derivation of correlation functions for random-vertex models and disordered-interacting-particle systems. *Phys. Rev. E*, 49:2726–2741, Apr 1994.
- [35] Gunter M. Schütz. Duality relations for asymmetric exclusion processes. *J. Statist. Phys.*, 86(5-6):1265–1287, 1997.
- [36] Michael Wheeler. Private communication to both authors in Boston, November 2022.
- [37] Michael Wheeler. Private communication to the first author in New York City, May 2017.
- [38] Zhengye Zhou. Orthogonal polynomial stochastic duality functions for multi-species SEP(2j) and multi-species IRW. *Symmetry, Integrability and Geometry: Methods and Applications*, dec 2021.