
PROBABILISTIC AND CAUSAL SATISFIABILITY: THE IMPACT OF MARGINALIZATION

A PREPRINT

Julian Dörfler^{*1}, Benito van der Zander^{*2}, Markus Bläser^{†1}, and Maciej Liśkiewicz^{†2}

{mblaeser,jdoerfler}@cs.uni-saarland.de; {maciej.liskiewicz,b.vanderzander}@uni-luebeck.de

¹Saarland University, Germany

²University of Lübeck, Germany

May 14, 2024

Abstract

The framework of Pearl’s Causal Hierarchy (PCH) formalizes three types of reasoning: observational, interventional, and counterfactual, that reflect the progressive sophistication of human thought regarding causation. We investigate the computational complexity aspects of reasoning in this framework focusing mainly on satisfiability problems expressed in probabilistic and causal languages across the PCH. That is, given a system of formulas in the standard probabilistic and causal languages, does there exist a model satisfying the formulas? The resulting complexity changes depending on the level of the hierarchy as well as the operators allowed in the formulas (addition, multiplication, or marginalization).

We focus on formulas involving marginalization that are widely used in probabilistic and causal inference, but whose complexity issues are still little explored. Our main contribution are the exact computational complexity results showing that linear languages (allowing addition and marginalization) yield NP^{PP} -, PSPACE -, and NEXP -complete satisfiability problems, depending on the level of the PCH. Moreover, we prove that the problem for the full language (allowing additionally multiplication) is complete for the class $\text{succ-}\exists\mathbb{R}$ for languages on the highest, counterfactual level. Previous work has shown that the satisfiability problem is complete for $\text{succ-}\exists\mathbb{R}$ on the lower levels leaving the counterfactual case open. Finally, we consider constrained models that are restricted to a small polynomial size. The constraint on the size reduces the complexity of the interventional and counterfactual languages to NEXP -complete.

1 Introduction

The development of the modern causal theory in AI and empirical sciences has greatly benefited from an influential structured approach to inference about causal phenomena, which is based on a reasoning hierarchy named “Ladder of Causation”, also often referred to as the “Pearl’s Causal Hierarchy” (PCH) ([Shpitser and Pearl, 2008, Pearl, 2009, Bareinboim et al., 2022], see also [Pearl and Mackenzie, 2018] for a gentle introduction to the topic). This three-level framework formalizes various types of reasoning that reflect the progressive sophistication of human thought regarding causation. It arises from a collection of causal mechanisms that model the “ground truth” of *unobserved* nature formalized within a

^{*}Contributing equally first authors.

[†]Contributing equally last authors.

Structural Causal Model (SCM). These mechanisms are then combined with three patterns of reasoning concerning *observed* phenomena expressed at the corresponding layers of the hierarchy, known as *probabilistic* (also called *associational* in the AI literature), *interventional*, and *counterfactual* (for formal definitions of these concepts, see Sec. 2).

A basic term at the probabilistic/associational layer is expressed as a common probability, such as¹ $\mathbb{P}(x, y)$. This may represent queries like “How likely does a patient have both diabetes ($X = x$) and high blood pressure ($Y = y$)?” From basic terms, we can build more complex terms by using additions (linear terms) or even arbitrary polynomials (polynomial terms). Formulas at this layer consist of Boolean combinations of (in)equalities of basic, linear or, in the general case, polynomial terms. The interventional patterns extend the basic probability terms by allowing the use of Pearl’s do-operator [Pearl, 2009] which models an experiment like a Randomized Controlled Trial [Fisher, 1936]. For instance, $\mathbb{P}([x]y)$ which², in general differs from $\mathbb{P}(y|x)$, allows to ask hypothetical questions such as, e.g., “How likely it is that a patient’s headache will be cured ($Y = y$) if he or she takes aspirin ($X = x$)?”. An example formula at this layer is $\mathbb{P}([x]y) = \sum_z \mathbb{P}(y|x, z)\mathbb{P}(z)$ which estimates the causal effect of the intervention $do(X = x)$ (all patients take aspirin) on outcome variable $Y = y$ (headache cure). It illustrates the use of a prominent back-door adjustment to eliminate the confounding effect of a factor represented by variable Z [Pearl, 2009]. The basic terms at the highest level of the hierarchy enable to formulate queries related to counterfactual situations. For example, $\mathbb{P}([X=x]Y=y|(X=x', Y=y'))$ expresses the probability that, for instance, a patient who did not receive a vaccine ($X = x'$) and died ($Y = y'$) would have lived ($Y = y$) if he or she had been vaccinated ($X = x$).

The computational complexity aspects of reasoning about uncertainty in this framework have been the subject of intensive studies in the past decades. The research has resulted in a large number of significant achievements, especially in the case of probabilistic inference with the input probability distributions encoded by Bayesian networks [Pearl, 1988]. These problems are of great interest in AI and important results from this perspective include [Cooper, 1990, Dagum and Luby, 1993, Roth, 1996, Park and Darwiche, 2004]. However, despite intensive research, many fundamental problems in the field remain open.

The main focus of our work is on the computational complexity of *satisfiability* problems and their *validity* counterparts which enable formulating precise assumptions on data and implications of causal explanations. The problems take as input a Boolean combination of (in)equalities of terms at the PCH-layer of interest with the task to decide if there exists a satisfying SCM (encoding a probability distribution) for the input formula or if the formula is valid for all SCMs, respectively. For example, for binary random variables X and Y , the formula consisting of the single equality $\sum_x \sum_y \mathbb{P}((X=x) \wedge (X \neq x \vee Y=y) \wedge (X \neq x \vee Y \neq y)) = 0$ in the language of the probabilistic layer is satisfied since there exists an SCM in which it is true (in fact, the formula holds in any SCM). An SCM for inputs at this layer can be identified with the standard joint probability distribution, in our case, with $P(X=x, Y=y)$, for $x, y \in \{0, 1\}$. Interestingly, the instance can be seen as a result of reduction from the not-satisfiable Boolean formula $a \wedge (\bar{a} \vee b) \wedge (\bar{a} \vee \bar{b})$. Based on this example, one can construct a polynomial reduction from the complement of the SAT problem to the satisfiability problem for the language of the probabilistic layer. E.g., the no-instance $(\bar{a} \vee b) \wedge (\bar{a} \vee \bar{b})$ is reduced to $\sum_x \sum_y \mathbb{P}((X \neq x \vee Y=y) \wedge (X \neq x \vee Y \neq y)) = 0$ which is false for all distributions $P(X, Y)$.

The complexity of the studied satisfiability problems depends on the combination of two factors: (1) the PCH-layer to which the basic terms belong and (2) the operators which can be used to specify the (in)equalities of the input formula. The most basic operators are “+” and “.” and, meaningful in causality, the unary summation operator Σ used to express marginalization. Of interest is also conditioning, which will be discussed in our paper, as well. The main interest of our research is focused on the precise characterization of the computational complexity of satisfiability problems (and their validity counterparts) for

¹In our paper, we consider random variables over discrete, finite domains. By an event we mean a propositional formula over atomic events of the form $X = x$, such as $(X = x \wedge Y = y)$ or $(X = x \vee Y \neq y)$. Moreover, by $\mathbb{P}(Y=y, X=x)$, etc., we mean, as usually, $\mathbb{P}(X=x \wedge Y=y)$. Finally by $\mathbb{P}(x, y)$ we abbreviate $\mathbb{P}(Y=y, X=x)$.

²A common and popular notation for the post-interventional probability is $\mathbb{P}(Y=y|do(X=x))$. In this paper, however, we use the notation $\mathbb{P}([X=x]Y=y)$ since it is more convenient for our analysis.

languages of all PCH layers, combined with increasing the expressiveness of (in)equalities by enabling the use of more complex operators.

Related Work to our Study. In their seminal paper, Fagin, Halpern and Megiddo [1990] explore the language of the lowest probabilistic layer of PCH consisting of Boolean combinations of (in)equalities of *basic* and *linear* terms. Besides the complete axiomatization for the used logic, they show that the problem of deciding satisfiability is NP-complete indicating that the complexity is surprisingly no worse than that of propositional logic. The authors subsequently extend the language to include (in)equalities of *polynomial* terms, aiming to facilitate reasoning about conditional probabilities. While they establish the existence of a PSPACE algorithm for deciding if such a formula is satisfiable, they leave the exact complexity open. Recently, Mossé, Ibeling, and Icard, [Mossé et al., 2022] resolved this issue by demonstrating that deciding satisfiability is $\exists\mathbb{R}$ -complete, where $\exists\mathbb{R}$ represents the well-studied class defined as the closure of the Existential Theory of the Reals (ETR) under polynomial-time many-one reductions. Furthermore, for the higher, more expressive PCH layers Mossé et al. prove that for (in)equalities of polynomial terms both at the interventional and the counterfactual layer the decision problems still remain $\exists\mathbb{R}$ -complete.

The languages used in these studies, and also in other works as, e.g., [Nilsson, 1986, Georgakopoulos et al., 1988, Ibeling and Icard, 2020], are able to fully express probabilistic reasoning, resp., inferring interventional and counterfactual predictions. In particular, they allow to express *marginalization* which is a common paradigm in this field. However, since the languages *do not* include the unary summation operator Σ , the abilities of expressing marginalization are relatively limited. Thus, for instance, to express the marginal distribution of a random variable Y over a subset of (binary) variables $\{Z_1, \dots, Z_m\} \subseteq \{X_1, \dots, X_n\}$ as $\sum_{z_1, \dots, z_m} \mathbb{P}(y, z_1, \dots, z_m)$, an encoding without summation requires an extension $\mathbb{P}(y, Z_1=0, \dots, Z_m=0) + \dots + \mathbb{P}(y, Z_1=1, \dots, Z_m=1)$ of exponential size in m . Consequently, to analyze the complexity aspects of the problems under study, languages allowing standard notation for encoding marginalization using the Σ operator are needed. In [van der Zander et al., 2023], the authors present a first systematic study in this setting. They introduce a new natural complexity class, named $\text{succ-}\exists\mathbb{R}$, which can be viewed as a succinct variant of $\exists\mathbb{R}$, and show that the satisfiability for the (in)equalities of *polynomial* terms, both at the *probabilistic* and *interventional* layer, are complete for $\text{succ-}\exists\mathbb{R}$. They leave open the exact complexity for the *counterfactual* case. Moreover, the remaining variants (basic and linear terms) remain unexplored for all PCH layers.

Contributions and Structure. The previous research establishes that, from a computational perspective, many problems for the interventional and counterfactual reasoning are not harder than for pure probabilistic reasoning. In our work, we show that the situation changes drastically if, to express marginalization, the common summation operator is used. Below we highlight our main contributions, partially summarized also in the table below.

- For combinations of (in)equalities of *basic* and *linear* terms, unlike previous results, the compact summation for marginalization increases the complexity, depending on the level of the PCH: from NP^{PP} -, through PSPACE-, to NEXP-completeness (Sec. 3).

- The counterfactual satisfiability for (in)equalities of *polynomial* terms is $\text{succ-}\exists\mathbb{R}$ -complete, which solves the open problem in [van der Zander et al., 2023] (Sec. 3.2);

- Unlike the languages devoid of the marginalization operator, the crucial small-model property is no longer satisfied. This property says that any satisfiable formula has a model of size bounded polynomially in the input length. Satisfiability with marginalization and with the additional requirement that there is a small model is complete for NEXP for interventional and counterfactual reasoning (Sec. 4).

Terms	\mathcal{L}_1 (prob.)	\mathcal{L}_2 (interv.)	\mathcal{L}_3 (count.)
basic lin	NP (a)		
poly	$\text{NP}_{\text{real}}^{(d)} = \exists\mathbb{R}$ (b)		
basic & marg. lin & marg.	NP^{PP} (1)	PSPACE (2)	NEXP (3)
poly & marg.	$\text{NEXP}_{\text{real}}^{(e)} = \text{succ-}\exists\mathbb{R}$ (c, 4)		

Sources: (a) for \mathcal{L}_1 [Fagin et al., 1990], for \mathcal{L}_2 and \mathcal{L}_3 [Mossé et al., 2022], (b) [Mossé et al., 2022], (c) for \mathcal{L}_1 and \mathcal{L}_2 [van der Zander et al., 2023], (d) [Erickson et al., 2022], (e) [Bläser et al., 2024]. Our results (1)-(3): Theorem 1, the completeness result, and (4) for \mathcal{L}_3 : Theorem 2.

2 Preliminaries

In this section, we give definitions of the main concepts of the theory of causation, including the Structural Causal Model (SCM), provide syntax and semantics for the languages of the PCH, and discuss the complexity classes used in the paper. To illustrate the main ideas behind the causality notions, we will start with an example that, we hope, will make it easier to understand the formal definitions. In the example, we consider a (hypothetical) scenario involving three attributes represented by binary random variables: pneumonia modeled by $Z = 1$, drug treatment (e.g., with antibiotics) represented by $X = 1$, and recovery, with $Y = 1$ (and $Y = 0$ meaning mortality). Below we describe a SCM which models an unobserved true mechanism behind this setting and the canonical patterns of reasoning that can be expressed at appropriate layers of the hierarchy.

Structural Causal Model An SCM is defined as a tuple $(\mathcal{F}, P, \mathbf{U}, \mathbf{X})$ which is of *unobserved nature* from the perspective of a researcher who studies the scenario. The SCM models the ground truth for the distribution $P(\mathbf{U})$ of the population and the mechanism \mathcal{F} . In our example, the model assumes three independent binary random variables $\mathbf{U} = \{U_1, U_2, U_3\}$, with probabilities: $P(U_1=1) = 0.75, P(U_2=1) = 0.8, P(U_3=1) = 0.4$, and specifies the mechanism $\mathcal{F} = \{F_1, F_2, F_3\}$ for the evaluation of the three endogenous (observed) random variables Z, X, Y as follows: $Z := F_1(U_1) = 1 - U_1$; $X := F_2(Z, U_2) = (1 - Z)U_2$; $Y := F_3(X, U_1, U_3) = X(1 - U_1)(1 - U_3) + (1 - X)(1 - U_1)U_3 + U_1U_3$. Thus, our model determines the distribution $P(\mathbf{u})$, for $\mathbf{u} = (u_1, u_2, u_3)$, and the values for the observed variables, as can be seen above.

The unobserved random variable U_1 models all circumstances that lead to pneumonia and Z is a function of U_1 (which may be more complex in real scenarios). Getting a treatment depends on having pneumonia but also on other circumstances, like having similar symptoms due to other diseases, and this is modelled by U_2 . So X is a function of Z and U_2 . Finally, mortality depends on all circumstances that lead to pneumonia, getting the treatment, and on further circumstances like having other diseases, which are modelled by U_3 . So Y is a function of U_1, X , and U_3 . We always assume that the dependency graph of the SCM is acyclic. This property is also called *semi-Markovian*.

Layer 1 Empirical sciences rely heavily on the use of observed data, which are typically represented as probability distributions over observed (measurable) variables. In our example, this is the distribution P over Z, X , and Y . The remaining variables U_1, U_2, U_3 , as well as the mechanism \mathcal{F} , are of unobserved nature. Thus, in our scenario, a researcher gets the probabilities (shown to the right) $P(z, x, y) = \sum_{\mathbf{u}} \delta_{\mathcal{F}, \mathbf{u}}(z, x, y) \cdot P(\mathbf{u})$, where vectors $\mathbf{u} = (u_1, u_2, u_3) \in \{0, 1\}^3$ and $\delta_{\mathcal{F}, \mathbf{u}}(z, x, y) = 1$ if $F_1(u_1)=z, F_2(z, u_2)=x$, and $F_3(x, u_1, u_2)=y$; otherwise $\delta_{\mathcal{F}, \mathbf{u}}(z, x, y) = 0$. The relevant query in our scenario $P(Y=1|X=1)$ can be evaluated as $P(Y=1|X=1) = P(Y=1, X=1)/P(X=1) = 0.24/0.6 = 0.4$ which says that the probability for recovery ($Y=1$) is only 40% given that the patient took the drug ($X=1$). On the other hand, the query for $X=0$ can be evaluated as $P(Y=1|X=0) = P(Y=1, X=0)/P(X=0) = 0.16/0.4 = 0.4$ which may lead to the (wrong, see the next layer) opinion that the drug is irrelevant to recovery.

Layer 2 Consider a randomized drug trial in which each patient receives treatment, denoted as $do(X=1)$, regardless of pneumonia (Z) and other conditions (U_2). We model this by performing a hypothetical intervention in which we replace in \mathcal{F} the mechanism $F_2(Z, U_2)$ by the constant function 1 and leaving the remaining functions unchanged. If $\mathcal{F}_{X=1} =$

$\{F'_1=F_1, F'_2=1, F'_3=F_3\}$ denotes the new mechanism, then the *post-interventional* distribution $P([X=1]Z, Y)$ is specified as $P([X=1]z, y) = \sum_{\mathbf{u}} \delta_{\mathcal{F}_{X=1}, \mathbf{u}}(z, y) \cdot P(\mathbf{u})$, where $\delta_{\mathcal{F}_{X=1}, \mathbf{u}}$ denotes function δ as above, but for the new mechanism $\mathcal{F}_{X=1}$ (the distribution is shown on the

U_1	U_2	U_3	$P(\mathbf{u})$	Z	X	Y
0	0	0	0.03	1	0	0
0	0	1	0.02	1	0	1
0	1	0	0.12	1	0	0
0	1	1	0.08	1	0	1
1	0	0	0.09	0	0	0
1	0	1	0.06	0	0	1
1	1	0	0.36	0	1	0
1	1	1	0.24	0	1	1

Z	X	Y	$P(z, x, y)$
0	0	0	0.09
0	0	1	0.06
0	1	0	0.36
0	1	1	0.24
1	0	0	0.15
1	0	1	0.10

U_1	U_2	U_3	$P(\mathbf{u})$	Z	$X=1$	Y
0	0	0	0.03	1	1	1
0	0	1	0.02	1	1	0
0	1	0	0.12	1	1	1
0	1	1	0.08	1	1	0
1	0	0	0.09	0	1	0
1	0	1	0.06	0	1	1
1	1	0	0.36	0	1	0
1	1	1	0.24	0	1	1

Z	Y	$P([X=1]z, y)$
0	0	0.45
0	1	0.30
1	0	0.10
1	1	0.15

righthand side)³. To determine the causal effect of the drug on recovery, we compute, in an analogous way, the distribution $P([X=0]Z, Y)$ after the intervention $do(X=0)$, which means that all patients receive placebo. Then, comparing the value $P([X=1]Y=1) = 0.45$ with $P([X=0]Y=1) = 0.40$, we can conclude that $P([X=1]Y=1) - P([X=0]Y=1) > 0$. This can be interpreted as a positive (average) effect of the drug in the population which is in opposite to what has been inferred using the purely probabilistic reasoning of Layer 1. Note that it is not obvious how to compute the post-interventional distributions from the observed probability $P(Z, X, Y)$; Indeed, this is a challenging task in the field of causality.

Layer 3 The key phenomena that can be modeled and analyzed at this level are counterfactual situations. Imagine, e.g., in our scenario there is a group of patients who did not receive the treatment and died ($X=0, Y=0$). One may ask, what would be the outcome Y had they been given the treatment ($X=1$). In particular, one can ask what is the probability of recovery if we had given the treatment to the patients of this group. Using the formalism of Layer 3, we can express this as a counterfactual query: $P([X=1]Y=1|X=0, Y=0) = P([X=1](Y=1) \wedge (X=0, Y=0))/P(X=0, Y=0)$. Note that the event $[X=1](Y=1) \wedge (X=0, Y=0)$ incorporates simultaneously two counterfactual mechanisms: $\mathcal{F}_{X=1}$ and \mathcal{F} . This is the key difference to Layer 2, where we can only have one. We define the probability in this situation as follows:

$$P([X=x](Z=z, Y=y) \wedge (Z=z', X=x', Y=y')) = \sum_{\mathbf{u}} \delta_{\mathcal{F}_{X=x}, \mathbf{u}}(z, y) \cdot \delta_{\mathcal{F}, \mathbf{u}}(z', x', y') \cdot P(\mathbf{u}).$$

$X=0, Y=0$ is satisfied only for $(U_1, U_2, U_3) \in \{(0, 0, 0), (0, 1, 0), (1, 0, 0)\}$ (first table), and of them only $\{(0, 0, 0), (0, 1, 0)\}$ satisfies $[X=1]Y=1$ (third table). Thus, by marginalizing Z , we get $P([X=1]Y=1|X=0, Y=0) = 0.15/0.24 = 0.625$ which may be interpreted that more than 62% who did not receive treatment and died would have survived with treatment. Finally, we would like to note that, in general, the events of Layer 3 can be quite involved and incorporate simultaneously many counterfactual worlds.

2.1 The Languages of Causal Hierarchy

We give here an informal but reasonably precise description of syntax and semantics of the languages studied in this paper. For formal definitions, see Section A in Appendix.

We always consider discrete distributions and represent the values of the random variables as $Val = \{0, 1, \dots, c-1\}$. We denote by \mathbf{X} the set of variables used in a system and by capital letters X_1, X_2, \dots , we denote the individual variables which all share the same domain Val . A value of X_i is often denoted by x_i or a natural number. By an *atomic* event, we mean an event of the form $X = x$, where X is a random variable and x is a value in the domain of X . The language \mathcal{E}_{prop} of propositional formulas δ over atomic events is defined as the closure of such events under the Boolean operators \wedge and \neg . The atomic intervention is either empty \perp or of the form $X = x$. An intervention formula is a conjunction of atomic interventions. The language of post-interventional events, denoted as $\mathcal{E}_{post-int}$, consists the formulas of the form $[\alpha]\delta$ where α is an intervention and δ is in \mathcal{E}_{prop} . The counterfactual events, $\mathcal{E}_{counterfact}$, is the set $\mathcal{E}_{post-int}$ closed under \wedge and \neg .

The PCH consists of languages on three layers each of which is based on terms of the form $\mathbb{P}(\delta_i)$, with $i = 1, 2, 3$. For the observational (associational) language (Layer 1), we have $\delta_1 \in \mathcal{E}_{prop}$, for the interventional language (Layer 2), we have $\delta_2 \in \mathcal{E}_{post-int}$ and, for the counterfactual language (Layer 3), $\delta_3 \in \mathcal{E}_{counterfact}$. The expressive power and computational complexity properties of the languages depend largely on the operations that are allowed to apply on terms $\mathbb{P}(\delta_i)$. Allowing gradually more complex operators, we define the languages which are the subject of our studies. The terms for levels $i = 1, 2, 3$ are described as follows. The basic terms, denoted as \mathcal{T}_i^{base} , are probabilities $\mathbb{P}(\delta_i)$ as, e.g., $\mathbb{P}(X_1=x_1 \vee X_2=x_2)$ or $\mathbb{P}([X_1=x_1]X_2=x_2)$. From basic terms, we build more complex linear terms \mathcal{T}_i^{lin} by using additions and polynomial terms \mathcal{T}_i^{poly} by using arbitrary polynomials. By $\mathcal{T}_i^{base(\Sigma)}$, $\mathcal{T}_i^{lin(\Sigma)}$, and $\mathcal{T}_i^{poly(\Sigma)}$ we denote the corresponding sets of terms when adding a unary marginalization operator of the form $\sum_x \mathbf{t}$ for a term \mathbf{t} . In the summation, we have a dummy variable x which ranges over all values $0, 1, \dots, c-1$. The summation $\sum_x \mathbf{t}$ is a purely syntactical concept which represents the sum $\mathbf{t}[0/x] + \mathbf{t}[1/x] + \dots + \mathbf{t}[c-1/x]$, where by $\mathbf{t}[v/x]$, we mean the expression

³A common and popular notation for the post-interventional probability is $P(Z, Y|do(X=1))$. In this paper, we use the notation $P([X=1]Z, Y)$ since it is more convenient for analyses involving counterfactuals.

in which all occurrences of x are replaced with value v . For example, for $Val = \{0, 1\}$, the expression $\sum_x \mathbb{P}(Y=1, X=x)$ semantically represents $\mathbb{P}(Y=1, X=0) + \mathbb{P}(Y=1, X=1)$.

Now, let $Lab = \{\text{base}, \text{base}(\Sigma), \text{lin}, \text{lin}(\Sigma), \text{poly}, \text{poly}(\Sigma)\}$ denote the labels of all variants of languages. Then for each $* \in Lab$ and $i = 1, 2, 3$ we define the languages \mathcal{L}_i^* of Boolean combinations of inequalities in a standard way by the grammars: $\mathbf{f} ::= \mathbf{t} \leq \mathbf{t}' \mid \neg \mathbf{f} \mid \mathbf{f} \wedge \mathbf{f}$ where \mathbf{t}, \mathbf{t}' are terms in \mathcal{T}_i^* . Although the languages and their operations can appear rather restricted, all the usual elements of probabilistic and causal formulas can be encoded in a natural way (see Section A in Appendix for more discussion).

To define the semantics we use SCMs as in [Pearl, 2009, Sec. 3.2]. An SCM is a tuple $\mathfrak{M} = (\mathcal{F}, P, \mathbf{U}, \mathbf{X})$, with exogenous variables \mathbf{U} and endogenous variables $\mathbf{X} = \{X_1, \dots, X_n\}$. $\mathcal{F} = \{F_1, \dots, F_n\}$ consists of functions such that F_i calculates the value of variable X_i from the values (\mathbf{x}, \mathbf{u}) as $F_i(\mathbf{pa}_i, \mathbf{u}_i)$, where⁴, $\mathbf{Pa}_i \subseteq \mathbf{X}$ and $\mathbf{U}_i \subseteq \mathbf{U}$. P specifies a probability distribution of all exogenous variables \mathbf{U} . Without loss of generality, we assume that domains of exogenous variables are also discrete and finite.

For any basic intervention formula $[X_i = x_i]$ (which, in our notation, means $do(X_i = x_i)$), we denote by $\mathcal{F}_{X_i = x_i}$ the function obtained from \mathcal{F} by replacing F_i with the constant function $F_i(\mathbf{pa}_i, \mathbf{u}_i) := x_i$. We generalize this definition for any intervention α in a natural way and denote as \mathcal{F}_α the resulting functions. For any $\varphi \in \mathcal{E}_{prop}$, we write $\mathcal{F}, \mathbf{u} \models \varphi$ if φ is satisfied for values of \mathbf{X} calculated from the values \mathbf{u} . For any intervention α , we write $\mathcal{F}, \mathbf{u} \models [\alpha]\varphi$ if $\mathcal{F}_\alpha, \mathbf{u} \models \varphi$. And for all $\psi, \psi_1, \psi_2 \in \mathcal{E}_{counterfact}$, we write (i) $\mathcal{F}, \mathbf{u} \models \neg \psi$ if $\mathcal{F}, \mathbf{u} \not\models \psi$ and (ii) $\mathcal{F}, \mathbf{u} \models \psi_1 \wedge \psi_2$ if $\mathcal{F}, \mathbf{u} \models \psi_1$ and $\mathcal{F}, \mathbf{u} \models \psi_2$. Finally, for $\psi \in \mathcal{E}_{counterfact}$, let $S_\mathfrak{M} = \{\mathbf{u} \mid \mathcal{F}, \mathbf{u} \models \psi\}$. We define $\llbracket \mathbf{e} \rrbracket_\mathfrak{M}$, for some expression \mathbf{e} , recursively in a natural way, starting with basic terms as follows $\llbracket \mathbb{P}(\psi) \rrbracket_\mathfrak{M} = \sum_{\mathbf{u} \in S_\mathfrak{M}(\psi)} P(\mathbf{u})$ and, for $\delta \in \mathcal{E}_{prop}$, $\llbracket \mathbb{P}(\psi|\delta) \rrbracket_\mathfrak{M} = \llbracket \mathbb{P}(\psi \wedge \delta) \rrbracket_\mathfrak{M} / \llbracket \mathbb{P}(\delta) \rrbracket_\mathfrak{M}$, assuming that the expression is undefined if $\llbracket \mathbb{P}(\delta) \rrbracket_\mathfrak{M} = 0$. For two expressions \mathbf{e}_1 and \mathbf{e}_2 , we define $\mathfrak{M} \models \mathbf{e}_1 \leq \mathbf{e}_2$, if and only if, $\llbracket \mathbf{e}_1 \rrbracket_\mathfrak{M} \leq \llbracket \mathbf{e}_2 \rrbracket_\mathfrak{M}$. The semantics for negation and conjunction are defined in the usual way, giving the semantics for $\mathfrak{M} \models \varphi$ for any formula φ in \mathcal{L}_3^* .

2.2 Satisfiability for PCH Languages and Relevant Complexity Classes

The (decision) satisfiability problems for languages of PCH, denoted by $\text{SAT}_{\mathcal{L}_i^*}$, with $i = 1, 2, 3$ and $* \in Lab$, take as input a formula φ in the languages \mathcal{L}_i^* and ask whether there exists a model \mathfrak{M} such that $\mathfrak{M} \models \varphi$. Analogously, the validity problems for \mathcal{L}_i^* consists in deciding whether, for a given φ , $\mathfrak{M} \models \varphi$ holds for all models \mathfrak{M} . From the definitions, it is obvious that variants of the problems for the level i are at least as hard as the counterparts at the lower level.

To measure the computational complexity of $\text{SAT}_{\mathcal{L}_i^*}$, a central role play the following, well-known Boolean complexity classes $\text{NP}, \text{PSPACE}, \text{NEXP}$, and EXPSPACE (for formal definitions see, e.g., Arora and Barak [2009]). Recent research has shown that the precise complexity of several natural satisfiability problems can be expressed in terms of the classes over the real numbers $\exists\mathbb{R}$ and $\text{succ-}\exists\mathbb{R}$. Recall, that the existential theory of the reals (ETR) is the set of true sentences of the form

$$\exists x_1 \dots \exists x_n \varphi(x_1, \dots, x_n), \quad (1)$$

where φ is a quantifier-free Boolean formula over the basis $\{\vee, \wedge, \neg\}$ and a signature consisting of the constants 0 and 1, the functional symbols $+$ and \cdot , and the relational symbols $<, \leq$, and $=$. The sentence is interpreted over the real numbers in the standard way. The theory forms its own complexity class $\exists\mathbb{R}$ which is defined as the closure of the ETR under polynomial time many-one reductions [Grigoriev and Vorobjov, 1988, Canny, 1988, Schaefer, 2009]. A succinct variant of ETR, denoted as succ-ETR , and the corresponding class $\text{succ-}\exists\mathbb{R}$, have been introduced in van der Zander et al. [2023]. succ-ETR is the set of all Boolean circuits C that encode a true sentence φ as in (1) as follows. Assume that C computes a function $\{0, 1\}^N \rightarrow \{0, 1\}^M$. Then $\{0, 1\}^N$ represents the node set of the tree underlying φ and $C(i)$ is an encoding of the description of node i , consisting of the label of i , its parent, and its two children. The variables in φ are x_1, \dots, x_{2N} . As in the case of $\exists\mathbb{R}$, to $\text{succ-}\exists\mathbb{R}$ belong all languages which are polynomial time many-one reducible to succ-ETR .

⁴We consider recursive models, that is, we assume the endogenous variables are ordered such that variable X_i (i.e. function F_i) is not affected by any X_j with $j > i$.

For two computational problems A, B , we will write $A \leq_P B$ if A can be reduced to B in polynomial time, which means A is not harder to solve than B . A problem A is complete for a complexity class \mathcal{C} , if $A \in \mathcal{C}$ and, for every other problem $B \in \mathcal{C}$, it holds $B \leq_P A$. By $\text{co-}\mathcal{C}$ we denote the class of all problems A such that its complements \bar{A} belong to \mathcal{C} .

3 The Computational Complexity of Satisfiability for Languages of the Causal Hierarchy

The computational complexity of probabilistic causal satisfiability problems has been a subject of intensive studies for languages which do not allow explicitly marginalization via summation operator Σ . Very recently van der Zander et al. [2023] addressed the problem for polynomial languages. Below, we summarise these results (informally presented in the Introduction):

- (a) $\text{SAT}_{\mathcal{L}_i}^{\text{base}}$ and $\text{SAT}_{\mathcal{L}_i}^{\text{lin}}$ are NP-complete, for $i = 1, 2, 3$ [Fagin et al., 1990, Mossé et al., 2022],
- (b) $\text{SAT}_{\mathcal{L}_i}^{\text{poly}}$ are $\exists\mathbb{R}$ -complete, for $i = 1, 2, 3$ [Mossé et al., 2022], and
- (c) $\text{SAT}_{\mathcal{L}_i}^{\text{poly}(\Sigma)}$ are $\text{succ-}\exists\mathbb{R}$ -complete, for $i = 1, 2$ [van der Zander et al., 2023]⁵.

Accordingly, the validity problems for the languages above are co-NP- , $\text{co-}\exists\mathbb{R}\text{-}$ resp., $\text{co-succ-}\exists\mathbb{R}\text{-}$ complete.

3.1 Linear Languages

In this section, we investigate the complexity of the satisfiability for the languages of the three levels of PCH that do not allow multiplication, that is, $\mathcal{L}_1^*, \mathcal{L}_2^*, \mathcal{L}_3^*$, with $*$ $\in \{\text{base}(\Sigma), \text{lin}(\Sigma)\}$. Due to Fagin et al. [1990] and Mossé et al. [2022], we know that the satisfiability problems across all the levels are NP-complete if compact summation, i.e., marginalization is not allowed.

Summations increase the complexity, depending on the level of the PCH in which they are introduced. On the probabilistic level, since the primitives can contain Boolean formulas, one can count the number of all satisfying assignments of a Boolean formula by summing over all possible values for the random variables in the formula. Determining this count is the canonical PP-complete problem, so evaluating the equations given a model is PP-hard. From this, together with the need of finding a model, we conclude that $\text{SAT}_{\mathcal{L}_1}^{\text{lin}(\Sigma)}$ is NP^{PP} -complete. (All technical details, precise statements, and proofs can be found in Section B.)

In causal formulas, one can perform interventions to set the value of a variable, which possibly changes the value of other variables. This correspond to the choice of a Boolean variable value by an existential or universal quantifier, since in a Boolean formula with multiple quantifiers, the value chosen by each quantifier can depend on the values chosen by earlier quantifiers. Thus, an interventional equation can encode a quantified Boolean formula. This makes $\text{SAT}_{\mathcal{L}_2}^{\text{lin}(\Sigma)}$ PSPACE-hard. It is even PSPACE-complete, since one can evaluate the interventions one at a time without storing the entire model because each primitive can only contain one intervention.

On the counterfactual level, one can perform multiple interventions, and thus compare different functions of the model to each other. Hence, the formulas can only be evaluated if the entire, exponential-sized model is known. Thus deciding the satisfiability requires exponential time and nondeterminism to find the model, making $\text{SAT}_{\mathcal{L}_3}^{\text{lin}(\Sigma)}$ NEXP-complete.

We summarize this in the following theorem, with the detailed proofs given in the appendix in Section B.1.

Theorem 1. *The complexities of the satisfiability problems for the linear languages can be characterized as follows:*

- (1) $\text{SAT}_{\mathcal{L}_1}^{\text{base}(\Sigma)}$ and $\text{SAT}_{\mathcal{L}_1}^{\text{lin}(\Sigma)}$ are NP^{PP} -complete,
- (2) $\text{SAT}_{\mathcal{L}_2}^{\text{base}(\Sigma)}$ and $\text{SAT}_{\mathcal{L}_2}^{\text{lin}(\Sigma)}$ are PSPACE-complete, and
- (3) $\text{SAT}_{\mathcal{L}_3}^{\text{base}(\Sigma)}$ and $\text{SAT}_{\mathcal{L}_3}^{\text{lin}(\Sigma)}$ are NEXP-complete.

⁵They show this for the variant where the primitives are also allowed to be conditional probabilities. We extend this to our variant in Theorem 2

The hardness of $\text{SAT}_{\mathcal{L}_1}^{\text{lin}(\Sigma)}$ depends on negations in the Boolean formulas, which make it difficult to count all satisfying assignments. Without negations, marginalization just removes variables, e.g. $\sum_x \mathbb{P}(X=x \wedge Y=y)$ becomes $\mathbb{P}(Y=y)$:

Proposition 1. $\text{SAT}_{\mathcal{L}_1}^{\text{base}(\Sigma)}$ and $\text{SAT}_{\mathcal{L}_1}^{\text{lin}(\Sigma)}$ are NP-complete if the primitives in $\mathcal{E}_{\text{post-int}}$ are restricted to not contain negations.

3.2 succ- $\exists\mathbb{R}$ -completeness of Satisfiability Problems

In [van der Zander et al., 2023] the authors prove that $\text{SAT}_{\mathcal{L}_1}^{\text{poly}(\Sigma)}$ and $\text{SAT}_{\mathcal{L}_2}^{\text{poly}(\Sigma)}$ are complete for succ- $\exists\mathbb{R}$ whenever the basic terms are allowed to also contain conditional probabilities. They, however, leave open the exact complexity status of $\text{SAT}_{\mathcal{L}_3}^{\text{poly}(\Sigma)}$. Here we show that the problem is in succ- $\exists\mathbb{R}$, with and without conditional probabilities, which proves its succ- $\exists\mathbb{R}$ -completeness.

Lemma 1. $\text{SAT}_{\mathcal{L}_3}^{\text{poly}(\Sigma)} \in \text{succ-}\exists\mathbb{R}$. This also holds true if we allow the basic terms to contain conditional probabilities.

This follows from the $\text{NEXP}_{\text{real}}$ machine model of [Bläser et al., 2024] combined with the algorithm proving $\text{SAT}_{\mathcal{L}_3}^{\text{poly}} \in \exists\mathbb{R} = \text{NP}_{\text{real}}$ from Mossé et al. [2022]. In Theorem 2 below, we conclude the completeness result for $\text{SAT}_{\mathcal{L}_3}^{\text{poly}(\Sigma)}$ strengthening slightly the hardness results for $\text{SAT}_{\mathcal{L}_1}^{\text{poly}(\Sigma)}$ and $\text{SAT}_{\mathcal{L}_2}^{\text{poly}(\Sigma)}$ of van der Zander et al. [2023] for languages which disallow conditional probabilities.

Theorem 2. $\text{SAT}_{\mathcal{L}_1}^{\text{poly}(\Sigma)}$, $\text{SAT}_{\mathcal{L}_2}^{\text{poly}(\Sigma)}$, and $\text{SAT}_{\mathcal{L}_3}^{\text{poly}(\Sigma)}$ are succ- $\exists\mathbb{R}$ -complete. This also holds true if we allow the basic terms to contain conditional probabilities.

4 Probabilistic and Causal Reasoning with a Small Model

An important feature of $\text{SAT}_{\mathcal{L}_i}^{\text{poly}}$ instances, over all levels $i = 1, 2, 3$, is the small model property which says that for every satisfiable formula there is a model whose size is bounded polynomially with respect to the length of the input. This property appeared to be useful to prove the memberships of $\text{SAT}_{\mathcal{L}_i}^{\text{poly}}$ in NP and in $\exists\mathbb{R}$ [Fagin et al., 1990, Ibeling and Icard, 2020, Mossé et al., 2022]. Interestingly, our membership proofs of $\text{SAT}_{\mathcal{L}_1}^{\text{lin}(\Sigma)}$ in NPP as well as of $\text{SAT}_{\mathcal{L}_2}^{\text{lin}(\Sigma)}$ in PSPACE (Theorem 1) rely on the property as well.

Apart from these advantages, the small model property is interesting in itself. For example, on the probabilistic layer, a polynomial formula φ (without the summation) over variables X_1, \dots, X_n is satisfiable, if and only if, there exists a Bayesian network $\mathcal{B} = (\mathcal{G}, P_{\mathcal{B}})$, with DAG \mathcal{G} over nodes X_1, \dots, X_n and $P_{\mathcal{B}}$ specifying set of conditional probability tables for each variable X_i , such that φ is true in \mathcal{B} and the total size of the tables is bounded by a polynomial of $|\varphi|$.

These have motivated the introduction of the small-model problem, denoted as $\text{SAT}_{sm, \mathcal{L}_1}^{\text{poly}(\Sigma)}$, which is defined like $\text{SAT}_{\mathcal{L}_1}^{\text{poly}(\Sigma)}$ with the additional constraint that a satisfying distribution should only have polynomially large support, that is, only polynomially many entries in the exponentially large table of probabilities are nonzero [Bläser et al., 2024]. The authors achieve this by extending an instance with an additional unary input $p \in \mathbb{N}$ and requiring that the satisfying distribution has a support of size at most p . We define $\text{SAT}_{sm, \mathcal{L}_2}^{\text{poly}(\Sigma)}$ and $\text{SAT}_{sm, \mathcal{L}_3}^{\text{poly}(\Sigma)}$ in the same way. Formally, we use the following:

Definition 1. The decision problems $\text{SAT}_{sm, \mathcal{L}_i}^{\text{poly}(\Sigma)}$, with $i = 1, 2, 3$, take as input a formula $\varphi \in \mathcal{L}_i^{\text{poly}(\Sigma)}$ and a unary encoded number $p \in \mathbb{N}$ and ask whether there exists a model $\mathfrak{M} = (\mathcal{F}, P, \mathbf{U} = \{U_1, \dots, U_m\}, \mathbf{X})$ such that $\mathfrak{M} \models \varphi$ and $\#\{(u_1, \dots, u_m) : P(U_1 = u_1, \dots, U_m = u_m) > 0\} \leq p$.

Bläser et al. [2024] have shown that $\text{SAT}_{sm, \mathcal{L}_1}^{\text{poly}(\Sigma)}$ is complete for a new complexity class $\exists\mathbb{R}^{\Sigma}$, a class that extends the existential theory of reals with summation operators. This is not a succinct class, so it is different from succ- $\exists\mathbb{R}$, in fact they show

$$\exists\mathbb{R} \subseteq \exists\mathbb{R}^{\Sigma} \subseteq \text{PSPACE} \subseteq \text{NEXP} \subseteq \text{succ-}\exists\mathbb{R} = \text{NEXP}_{\text{real}},$$

where $\text{NEXP}_{\text{real}}$ is the class of problems decided by nondeterministic real Random Access Machines (RAMs) in exponential time (see [Bläser et al., 2024] for the exact details). However once we allow for interventions the complexity increases to NEXP -completeness. Note however that this is likely still weaker than the $\text{NEXP}_{\text{real}}$ -completeness of $\text{SAT}_{\mathcal{L}_2}^{\text{poly}(\Sigma)}$ and $\text{SAT}_{\mathcal{L}_3}^{\text{poly}(\Sigma)}$.

Theorem 3. $\text{SAT}_{sm, \mathcal{L}_2}^{\text{poly}(\Sigma)}$ and $\text{SAT}_{sm, \mathcal{L}_3}^{\text{poly}(\Sigma)}$ are NEXP -complete.

This follows because a causal model consists of a probabilistic part $\mathbb{P}(\mathbf{u})$ and a deterministic, causal part \mathcal{F} . In the small model, $\mathbb{P}(\mathbf{u})$ is restricted to have polynomial size, but \mathcal{F} can still have exponential size. In $\text{SAT}_{sm, \mathcal{L}_1}^{\text{poly}(\Sigma)}$, one cannot reason about \mathcal{F} , but $\text{SAT}_{sm, \mathcal{L}_2}^{\text{poly}(\Sigma)}$ and $\text{SAT}_{sm, \mathcal{L}_3}^{\text{poly}(\Sigma)}$ can access the full power of \mathcal{F} . Since NEXP subsumes $\exists\mathbb{R}$, reasoning about polynomial many real numbers (probabilities) cannot increase the complexity.

5 Discussion

The standard, well-studied class $\exists\mathbb{R}$ is defined as the closure of the ETR problem under polynomial time many-one reductions. The class $\text{succ-}\exists\mathbb{R}$, which can be viewed as a succinct variant of $\exists\mathbb{R}$, was proposed in [van der Zander et al., 2023]. Based on this structure of complexity classes, we present the computational complexities of satisfiability problems for languages at all the levels of the PCH. Our new completeness results nicely extend and complement the previous achievements by Fagin et al. [1990], Mossé et al. [2022], van der Zander et al. [2023] and Bläser et al. [2024]. The main focus of our research was on languages allowing the use of marginalization which is expressed in the languages by a summation operator Σ over the domain of the random variables. This captures the standard notation commonly used in probabilistic and causal inference.

A very interesting feature of the satisfiability problems for the full, polynomial languages is the following property. For both variants, with and without summation operators, while the expressive powers of the corresponding languages differ, the complexities of the corresponding satisfiability problems at all three levels of the PCH are the same. We have, namely, that all $\text{SAT}_{\mathcal{L}_i}^{\text{poly}(\Sigma)}$ problems are complete for $\text{succ-}\exists\mathbb{R}$ and all $\text{SAT}_{\mathcal{L}_i}^{\text{poly}}$ are $\exists\mathbb{R}$ -complete. Interestingly, the same holds for linear languages *without* marginalization, too: due to Fagin et al. [1990] and Mossé et al. [2022], we know that $\text{SAT}_{\mathcal{L}_i}^{\text{lin}}$ are NP -complete for all levels i . We find that the situation changes drastically in the case of linear languages *allowing* the summation operator Σ . One of our main results characterizes the complexities of $\text{SAT}_{\mathcal{L}_1}^{\text{lin}(\Sigma)}$, $\text{SAT}_{\mathcal{L}_2}^{\text{lin}(\Sigma)}$, and $\text{SAT}_{\mathcal{L}_3}^{\text{lin}(\Sigma)}$ problems as NP^{PP} , PSPACE , and NEXP -complete, resp. The analogous completeness results are true for $\text{SAT}_{\mathcal{L}_i}^{\text{base}(\Sigma)}$.

Another interesting feature is that the completeness results for linear languages are expressed in terms of standard Boolean classes while the completeness of satisfiability for languages involving polynomials over the probabilities requires classes over the reals.

References

- Antonis Achilleos. NEXP -completeness and universal hardness results for justification logic. In *International Computer Science Symposium in Russia*, pages 27–52. Springer, 2015.
- Sanjeev Arora and Boaz Barak. *Computational complexity: a modern approach*. Cambridge University Press, 2009.
- Elias Bareinboim, Juan D. Correa, Duligur Ibeling, and Thomas Icard. *On Pearl’s Hierarchy and the Foundations of Causal Inference*, pages 507–556. Association for Computing Machinery, New York, NY, USA, 2022.
- Markus Bläser, Julian Dörfler, Maciej Liśkiewicz, and Benito van der Zander. The existential theory of the reals with summation operators. *Manuscript*, 2024.
- John Canny. Some algebraic and geometric computations in PSPACE . In *Proceedings of the twentieth annual ACM symposium on Theory of computing*, pages 460–467. ACM, 1988.
- Gregory F Cooper. The computational complexity of probabilistic inference using bayesian belief networks. *Artificial intelligence*, 42(2-3):393–405, 1990.

- Paul Dagum and Michael Luby. Approximating probabilistic inference in Bayesian belief networks is NP-hard. *Artificial Intelligence*, 60(1):141–153, 1993.
- Jeff Erickson, Ivor Van Der Hoog, and Tillmann Miltzow. Smoothing the gap between NP and ER. *SIAM Journal on Computing*, pages FOCS20–102, 2022.
- Ronald Fagin, Joseph Y Halpern, and Nimrod Megiddo. A logic for reasoning about probabilities. *Information and computation*, 87(1-2):78–128, 1990.
- Ronald Aylmer Fisher. Design of experiments. *British Medical Journal*, 1(3923):554, 1936.
- George Georgakopoulos, Dimitris Kavvadias, and Christos H Papadimitriou. Probabilistic satisfiability. *Journal of Complexity*, 4(1):1–11, 1988.
- Dima Grigoriev and Nicolai Vorobjov. Solving systems of polynomial inequalities in subexponential time. *J. Symb. Comput.*, 5(1/2):37–64, 1988.
- Duligur Ibeling and Thomas Icard. Probabilistic reasoning across the causal hierarchy. In *The 34th AAAI Conference on Artificial Intelligence, AAAI 2020*, pages 10170–10177. AAAI Press, 2020.
- Harry R. Lewis. Complexity results for classes of quantificational formulas. *Journal of Computer and System Sciences*, 21(3):317 – 353, 1980.
- Michael L Littman, Judy Goldsmith, and Martin Mundhenk. The computational complexity of probabilistic planning. *Journal of Artificial Intelligence Research*, 9:1–36, 1998.
- Milan Mossé, Duligur Ibeling, and Thomas Icard. Is causal reasoning harder than probabilistic reasoning? *The Review of Symbolic Logic*, pages 1–26, 2022.
- Nils J Nilsson. Probabilistic logic. *Artificial Intelligence*, 28(1):71–87, 1986.
- James D Park and Adnan Darwiche. Complexity results and approximation strategies for MAP explanations. *Journal of Artificial Intelligence Research*, 21:101–133, 2004.
- Judea Pearl. *Probabilistic reasoning in intelligent systems: networks of plausible inference*. Morgan Kaufmann, 1988.
- Judea Pearl. *Causality*. Cambridge University Press, 2009. ISBN 0-521-77362-8.
- Judea Pearl and Dana Mackenzie. *The book of why: the new science of cause and effect*. Basic books, 2018.
- James Renegar. On the computational complexity and geometry of the first-order theory of the reals. Part I: Introduction. Preliminaries. The geometry of semi-algebraic sets. The decision problem for the existential theory of the reals. *Journal of symbolic computation*, 13(3):255–299, 1992.
- Dan Roth. On the hardness of approximate reasoning. *Artificial Intelligence*, 82(1-2):273–302, 1996.
- Marcus Schaefer. Complexity of some geometric and topological problems. In *International Symposium on Graph Drawing*, pages 334–344. Springer, 2009.
- Ilya Shpitser and Judea Pearl. Complete identification methods for the causal hierarchy. *Journal of Machine Learning Research*, 9(Sep):1941–1979, 2008.
- Benito van der Zander, Markus Bläser, and Maciej Liśkiewicz. The hardness of reasoning about probabilities and causality. In *Proc. Joint Conference on Artificial Intelligence (IJCAI 2023)*, 2023.
- Junzhe Zhang, Jin Tian, and Elias Bareinboim. Partial counterfactual identification from observational and experimental data. In *International Conference on Machine Learning*, pages 26548–26558. PMLR, 2022.

A Syntax and Semantics of the Languages of PCH

We always consider discrete distributions in the probabilistic and causal languages studied in this paper. We represent the values of the random variables as $Val = \{0, 1, \dots, c-1\}$ and denote by \mathbf{X} the set of random variables used in a system. By capital letters X_1, X_2, \dots , we denote the individual variables and assume, w.l.o.g., that they all share the same domain Val . A value of X_i is often denoted by x_i or a natural number. In this section, we describe syntax and semantics of the languages starting with probabilistic ones and then we provide extensions to the causal systems.

By an *atomic* event, we mean an event of the form $X = x$, where X is a random variable and x is a value in the domain of X . The language \mathcal{E}_{prop} of propositional formulas over atomic events is defined as the closure of such events under the Boolean operators \wedge and \neg . To specify the syntax of interventional and counterfactual events we define the intervention and extend the syntax of \mathcal{E}_{prop} to $\mathcal{E}_{post-int}$ and $\mathcal{E}_{counterfact}$, respectively, using the following grammars:

$$\begin{aligned} \mathcal{E}_{prop} \text{ is defined by } \mathbf{p} &::= X = x \mid \neg \mathbf{p} \mid \mathbf{p} \wedge \mathbf{p} \\ \mathcal{E}_{int} \text{ is defined by } \mathbf{i} &::= \top \mid X = x \mid \mathbf{i} \wedge \mathbf{i} \\ \mathcal{E}_{post-int} \text{ is defined by } \mathbf{p_i} &::= [\mathbf{i}] \mathbf{p} \\ \mathcal{E}_{counterfact} \text{ is defined by } \mathbf{c} &::= \mathbf{p_i} \mid \neg \mathbf{c} \mid \mathbf{c} \wedge \mathbf{c}. \end{aligned}$$

Note that since \top means that no intervention has been applied, we can assume that $\mathcal{E}_{prop} \subseteq \mathcal{E}_{post-int}$.

The PCH consists of three languages $\mathcal{L}_1, \mathcal{L}_2, \mathcal{L}_3$, each of which is based on terms of the form $\mathbb{P}(\delta)$. For the (observational or associational) language \mathcal{L}_1 , we have $\delta \in \mathcal{E}_{prop}$, for the (interventional) language \mathcal{L}_2 , we have $\delta \in \mathcal{E}_{post-int}$ and for the (counterfactual) language \mathcal{L}_3 , $\delta \in \mathcal{E}_{counterfact}$. The expressive power and computational complexity properties of the languages depend largely on the operations that we are allowed to apply to the basic terms. Allowing gradually more complex operators, we describe the languages which are the subject of our studies below. We start with the description of the languages \mathcal{T}_i^* of terms, with $i = 1, 2, 3$, using the following grammars⁶

$$\begin{array}{lcl} \mathcal{T}_i^{\text{base}} & \mathbf{t} ::= \mathbb{P}(\delta_i) & \left| \quad \mathcal{T}_i^{\text{base}(\Sigma)} \quad \mathbf{t} ::= \mathbb{P}(\delta_i) \mid \sum_x \mathbf{t} \right. \\ \mathcal{T}_i^{\text{lin}} & \mathbf{t} ::= \mathbb{P}(\delta_i) \mid \mathbf{t} + \mathbf{t} & \left| \quad \mathcal{T}_i^{\text{lin}(\Sigma)} \quad \mathbf{t} ::= \mathbb{P}(\delta_i) \mid \mathbf{t} + \mathbf{t} \mid \sum_x \mathbf{t} \right. \\ \mathcal{T}_i^{\text{poly}} & \mathbf{t} ::= \mathbb{P}(\delta_i) \mid \mathbf{t} + \mathbf{t} \mid -\mathbf{t} \mid \mathbf{t} \cdot \mathbf{t} & \left| \quad \mathcal{T}_i^{\text{poly}(\Sigma)} \quad \mathbf{t} ::= \mathbb{P}(\delta_i) \mid \mathbf{t} + \mathbf{t} \mid -\mathbf{t} \mid \mathbf{t} \cdot \mathbf{t} \mid \sum_x \mathbf{t} \right. \end{array}$$

where δ_1 are formulas in \mathcal{E}_{prop} , $\delta_2 \in \mathcal{E}_{post-int}$, $\delta_3 \in \mathcal{E}_{counterfact}$.

The probabilities of the form $\mathbb{P}(\delta_i)$ are called *primitives* or *basic terms*. In the summation operator \sum_x , we have a dummy variable x which ranges over all values $0, 1, \dots, c-1$. The summation $\sum_x \mathbf{t}$ is a purely syntactical concept which represents the sum $\mathbf{t}^{[0/x]} + \mathbf{t}^{[1/x]} + \dots + \mathbf{t}^{[c-1/x]}$, where by $\mathbf{t}^{[v/x]}$, we mean the expression in which all occurrences of x are replaced with value v . For example, for $Val = \{0, 1\}$, the expression $\sum_x \mathbb{P}(Y=1, X=x)$ semantically represents $\mathbb{P}(Y=1, X=0) + \mathbb{P}(Y=1, X=1)$. We note that the dummy variable x is not a (random) variable in the usual sense and that its scope is defined in the standard way.

In the table above, the terms in $\mathcal{T}_i^{\text{base}}$ are just basic probabilities with the events given by the corresponding languages \mathcal{E}_{prop} , $\mathcal{E}_{post-int}$, or $\mathcal{E}_{counterfact}$. Next, we extend terms by being able to compute sums of probabilities and by adding the same term several times, we also allow for weighted sums with weights given in unary. Note that this is enough to state all our hardness results. All matching upper bounds also work when we allow for explicit weights given in binary. In the case of $\mathcal{T}_i^{\text{poly}}$, we are allowed to build polynomial terms in the primitives. On the right-hand side of the table, we have the same three kind of terms, but to each of them, we add a marginalization operator as a building block.

The polynomial calculus $\mathcal{T}_i^{\text{poly}}$ was originally introduced by Fagin, Halpern, and Megiddo [Fagin et al., 1990] (for $i = 1$) to be able to express conditional probabilities by clearing denominators. While this works for $\mathcal{T}_i^{\text{poly}}$, this does not work in the case of $\mathcal{T}_i^{\text{poly}(\Sigma)}$, since clearing denominators with exponential sums creates expressions that are too large. But we

⁶In the given grammars we omit the brackets for readability, but we assume that they can be used in a standard way.

could introduce basic terms of the form $\mathbb{P}(\delta_i|\delta)$ with $\delta \in \mathcal{E}_{prop}$ explicitly. All our hardness proofs work without conditional probabilities but all our matching upper bounds are still true with explicited conditional probabilities. Expression as $\mathbb{P}(X=1) + \mathbb{P}(Y=2) \cdot \mathbb{P}(Y=3)$ is a valid term in $\mathcal{T}_1^{\text{poly}}$ and $\sum_z \mathbb{P}([X=0](Y=1, Z=z))$ and $\sum_z \mathbb{P}([X=0]Y=1, Z=z)$ are valid terms in the language $\mathcal{T}_3^{\text{poly}(\Sigma)}$, for example.

Now, let $Lab = \{\text{base}, \text{base}(\Sigma), \text{lin}, \text{lin}(\Sigma), \text{poly}, \text{poly}(\Sigma)\}$ denote the labels of all variants of languages. Then for each $* \in Lab$ and $i = 1, 2, 3$ we define the languages \mathcal{L}_i^* of Boolean combinations of inequalities in a standard way:

$$\mathcal{L}_i^* \text{ is defined by } \mathbf{f} ::= \mathbf{t} \leq \mathbf{t}' \mid \neg \mathbf{f} \mid \mathbf{f} \wedge \mathbf{f}, \quad \text{where } \mathbf{t}, \mathbf{t}' \text{ are terms in } \mathcal{T}_i^*.$$

Although the language and its operations can appear rather restricted, all the usual elements of probabilistic and causal formulas can be encoded. Namely, equality is encoded as greater-or-equal in both directions, e.g. $\mathbb{P}(x) = \mathbb{P}(y)$ means $\mathbb{P}(x) \geq \mathbb{P}(y) \wedge \mathbb{P}(y) \geq \mathbb{P}(x)$. The number 0 can be encoded as an inconsistent probability, i.e., $\mathbb{P}(X=1 \wedge X=2)$. In a language allowing addition and multiplication, any positive integer can be easily encoded from the fact $\mathbb{P}(\top) \equiv 1$, e.g. $4 \equiv (1+1)(1+1) \equiv (\mathbb{P}(\top) + \mathbb{P}(\top))(\mathbb{P}(\top) + \mathbb{P}(\top))$. If a language does not allow multiplication, one can show that the encoding is still possible. Note that these encodings barely change the size of the expressions, so allowing or disallowing these additional operators does not affect any complexity results involving these expressions.

To define the semantics of the languages, we use a structural causal model (SCM) as in [Pearl, 2009, Sec. 3.2]. An SCM is a tuple $\mathfrak{M} = (\mathcal{F}, P, \mathbf{U}, \mathbf{X})$, such that $\mathbf{V} = \mathbf{U} \cup \mathbf{X}$ is a set of variables partitioned into exogenous (unobserved) variables $\mathbf{U} = \{U_1, U_2, \dots\}$ and endogenous variables \mathbf{X} . The tuple $\mathcal{F} = \{F_1, \dots, F_n\}$ consists of functions such that function F_i calculates the value of variable X_i from the values (\mathbf{x}, \mathbf{u}) of other variables in \mathbf{V} as $F_i(\mathbf{pa}_i, \mathbf{u}_i)$ ⁷, where $\mathbf{Pa}_i \subseteq \mathbf{X}$ and $\mathbf{U}_i \subseteq \mathbf{U}$. P specifies a probability distribution of all exogenous variables \mathbf{U} . Since variables \mathbf{X} depend deterministically on the exogenous variables via functions F_i , \mathcal{F} and P obviously define the joint probability distribution of \mathbf{X} . Throughout this paper, we assume that domains of endogenous variables \mathbf{X} are discrete and finite. In this setting, exogenous variables \mathbf{U} could take values in any domains, including infinite and continuous ones. A recent paper [Zhang et al., 2022] shows, however, that any SCM over discrete endogenous variables is equivalent for evaluating post-interventional probabilities to an SCM where all exogenous variables are discrete with finite domains. As a consequence, throughout this paper, we assume that domains of exogenous variables \mathbf{U} are discrete and finite, too.

For any basic \mathcal{E}_{int} -formula $X_i = x_i$ (which, in our notation, means $do(X_i = x_i)$), we denote by $\mathcal{F}_{X_i=x_i}$ the function obtained from \mathcal{F} by replacing F_i with the constant function $F_i(\mathbf{v}) := x_i$. We generalize this definition for any interventions specified by $\alpha \in \mathcal{E}_{int}$ in a natural way and denote as \mathcal{F}_α the resulting functions. For any $\varphi \in \mathcal{E}_{prop}$, we write $\mathcal{F}, \mathbf{u} \models \varphi$ if φ is satisfied for values of \mathbf{X} calculated from the values \mathbf{u} . For $\alpha \in \mathcal{E}_{int}$, we write $\mathcal{F}, \mathbf{u} \models [\alpha]\varphi$ if $\mathcal{F}_\alpha, \mathbf{u} \models \varphi$. And for all $\psi, \psi_1, \psi_2 \in \mathcal{E}_{counterfact}$, we write (i) $\mathcal{F}, \mathbf{u} \models \neg\psi$ if $\mathcal{F}, \mathbf{u} \not\models \psi$ and (ii) $\mathcal{F}, \mathbf{u} \models \psi_1 \wedge \psi_2$ if $\mathcal{F}, \mathbf{u} \models \psi_1$ and $\mathcal{F}, \mathbf{u} \models \psi_2$. Finally, for $\psi \in \mathcal{E}_{counterfact}$, let $S_\mathfrak{M} = \{\mathbf{u} \mid \mathcal{F}, \mathbf{u} \models \psi\}$. We define $\llbracket \mathbf{e} \rrbracket_\mathfrak{M}$, for some expression \mathbf{e} , recursively in a natural way, starting with basic terms as follows $\llbracket \mathbb{P}(\psi) \rrbracket_\mathfrak{M} = \sum_{\mathbf{u} \in S_\mathfrak{M}(\psi)} P(\mathbf{u})$ and, for $\delta \in \mathcal{E}_{prop}$, $\llbracket \mathbb{P}(\psi|\delta) \rrbracket_\mathfrak{M} = \llbracket \mathbb{P}(\psi \wedge \delta) \rrbracket_\mathfrak{M} / \llbracket \mathbb{P}(\delta) \rrbracket_\mathfrak{M}$, assuming that the expression is undefined if $\llbracket \mathbb{P}(\delta) \rrbracket_\mathfrak{M} = 0$. For two expressions \mathbf{e}_1 and \mathbf{e}_2 , we define $\mathfrak{M} \models \mathbf{e}_1 \leq \mathbf{e}_2$, if and only if, $\llbracket \mathbf{e}_1 \rrbracket_\mathfrak{M} \leq \llbracket \mathbf{e}_2 \rrbracket_\mathfrak{M}$. The semantics for negation and conjunction are defined in the usual way, giving the semantics for $\mathfrak{M} \models \varphi$ for any formula φ in \mathcal{L}_3^* .

B Technical Details and Proofs

B.1 Proofs of Section 3

In this section we prove the completeness results in Theorem 1: the first result will follow from Lemma 2 and 3, Result (2) from Lemma 4 and 5, and (3) will follow from Lemma 6. Finally, we show the proof of Proposition 1.

First we show that a sum in the probabilistic language can be partitioned into a sum over probabilities and a sum over purely logical terms.

⁷We consider recursive models, that is, we assume the endogenous variables are ordered such that variable X_i (i.e. function F_i) is not affected by any X_j with $j > i$.

Fact 1. Let $\delta \in \mathcal{E}_{prop}$ be a propositional formula over variables X_{i_1}, \dots, X_{i_l} . A sum $\sum_{x_{i_1}} \dots \sum_{x_{i_l}} \mathbb{P}(\delta)$ is equivalent to

$$\sum_{\hat{x}_1} \dots \sum_{\hat{x}_n} p_{\hat{x}_1 \dots \hat{x}_n} \sum_{x_{i_1}} \dots \sum_{x_{i_l}} \delta_{\hat{x}_1 \dots \hat{x}_n}(x_{i_1}, \dots, x_{i_l})$$

where the range of the sums is the entire domain, $p_{\hat{x}_1 \dots \hat{x}_n}$ is the probability of $\mathbb{P}(X_1 = \hat{x}_1 \wedge \dots \wedge X_n = \hat{x}_n)$ and $\delta_{\hat{x}_1 \dots \hat{x}_n}(x_{i_1}, \dots, x_{i_l})$ a function that returns 1 if the implication $(X_1 = \hat{x}_1 \wedge \dots \wedge X_n = \hat{x}_n) \rightarrow \delta(x_{i_1}, \dots, x_{i_l})$ is a tautology and 0 otherwise.

Proof. $\sum_{x_{i_1}} \dots \sum_{x_{i_l}} \mathbb{P}(\delta)$ is equivalent to

$$\begin{aligned} & \sum_{x_{i_1}} \dots \sum_{x_{i_l}} \sum_{\hat{x}_1} \dots \sum_{\hat{x}_n} \mathbb{P}((X_1 = \hat{x}_1 \wedge \dots \wedge X_n = \hat{x}_n) \wedge \delta) \\ &= \sum_{\hat{x}_1} \dots \sum_{\hat{x}_n} \sum_{x_{i_1}} \dots \sum_{x_{i_l}} \mathbb{P}((X_1 = \hat{x}_1 \wedge \dots \wedge X_n = \hat{x}_n) \wedge \delta) \\ &= \sum_{\hat{x}_1} \dots \sum_{\hat{x}_n} \sum_{x_{i_1}} \dots \sum_{x_{i_l}} p_{\hat{x}_1 \dots \hat{x}_n} \delta_{\hat{x}_1 \dots \hat{x}_n}(x_{i_1}, \dots, x_{i_l}) \\ &= \sum_{\hat{x}_1} \dots \sum_{\hat{x}_n} p_{\hat{x}_1 \dots \hat{x}_n} \sum_{x_{i_1}} \dots \sum_{x_{i_l}} \delta_{\hat{x}_1 \dots \hat{x}_n}(x_{i_1}, \dots, x_{i_l}), \end{aligned}$$

which completes the proof. \square

Since the second sums only depend on $\hat{x}_1 \dots \hat{x}_n$ and not on $p_{\hat{x}_1 \dots \hat{x}_n}$, they can be calculated without knowing the probability distribution. Due to the dependency on $\hat{x}_1 \dots \hat{x}_n$ the expression cannot be simplified further in general. However, if δ contains no constant events like $X_{i_j} = 0$ but only events $X_{i_j} = x_{i_j}$ depending on the summation variables x_{i_j} , the sum always includes one iteration where the event occurs and $c - 1$ iterations where it does not. Thus the sum $\sum_{x_{i_1}} \dots \sum_{x_{i_l}} \delta_{\hat{x}_1 \dots \hat{x}_n}(x_{i_1}, \dots, x_{i_l})$ is constant and effectively counts the number of satisfying assignments to the formula δ . Since the sum $\sum_{\hat{x}_1} \dots \sum_{\hat{x}_n} p_{\hat{x}_1 \dots \hat{x}_n}$ is always 1, the original sum $\sum_{x_{i_1}} \dots \sum_{x_{i_l}} \mathbb{P}(\delta)$ also counts the number of satisfying assignments.

Lemma 2. $\text{SAT}_{\mathcal{L}_1}^{\text{lin}(\Sigma)}$ is in NP^{PP} .

Proof. First we need to show that satisfiable instances have solutions of polynomial size.

We write each (sum of a) primitive in the arithmetic expressions as

$$\sum_{\hat{x}_1} \dots \sum_{\hat{x}_n} p_{\hat{x}_1 \dots \hat{x}_n} \sum_{x_{i_1}} \dots \sum_{x_{i_l}} \delta_{\hat{x}_1 \dots \hat{x}_n}(x_{i_1}, \dots, x_{i_l})$$

according to Fact 1.

The value of all $p_{\hat{x}_1 \dots \hat{x}_n}$ can be encoded as a vector $\vec{p} \in \mathbb{R}^{c^n}$. For each $\hat{x}_1, \dots, \hat{x}_n$, the sum $\sum_{x_{i_1}} \dots \sum_{x_{i_l}} \delta_{\hat{x}_1 \dots \hat{x}_n}(x_{i_1}, \dots, x_{i_l})$ is a constant integer $\leq c^l$. Suppose there are m such sums or primitives in the instance, whose values can be encoded as a matrix $A \in \mathbb{R}^{m \times c^n}$.

Then the value of every sum in the instance is given by the vector $A\vec{p} \in \mathbb{R}^m$.

There exists a non-negative vector $\vec{q} \in \mathbb{Q}^{c^n}$ containing at most m non-zero entries with $A\vec{q} = A\vec{p}$ (Lemma 2.5 in Fagin et al. [1990]). By including a constraint $\sum_{\hat{x}_1} \dots \sum_{\hat{x}_n} p_{\hat{x}_1 \dots \hat{x}_n} = 1$ when constructing the matrix A , we can ensure that all values in \vec{q} are valid probabilities. \vec{q} is a polynomial sized solution and can be guessed nondeterministically.

For this solution, the sum $\sum_{\hat{x}_1} \dots \sum_{\hat{x}_n} p_{\hat{x}_1 \dots \hat{x}_n}$ can then be evaluated (over all non-zero entries $p_{\hat{x}_1 \dots \hat{x}_n}$) in polynomial time. Each sum $\sum_{x_{i_1}} \dots \sum_{x_{i_l}} \delta_{\hat{x}_1 \dots \hat{x}_n}(x_{i_1}, \dots, x_{i_l})$ can be evaluated using the PP oracle because a PP oracle is equivalent to a #P oracle which can count the number of satisfying assignments of δ . \square

Now we can tackle the hardness of $\text{SAT}_{\mathcal{L}_1}^{\text{base}(\Sigma)}$:

Lemma 3. $\text{SAT}_{\mathcal{L}_1}^{\text{base}(\Sigma)}$ is NP^{PP} -hard.

Proof. The canonical NP^{PP} -complete problem E-MajSat is deciding the satisfiability of a formula $\psi : \exists x_1 \dots x_n \#y_1 \dots y_n \phi \geq 2^{n-1}$, i.e. deciding whether the Boolean formula ϕ has a majority of satisfying assignments to Boolean variables y_i after choosing Boolean variables x_i existentially [Littman et al., 1998].

We will reduce this to $\text{SAT}_{\mathcal{L}_1}^{\text{base}(\Sigma)}$. Let there be $2n$ random variables $X_1, \dots, X_n, Y_1, \dots, Y_n$ associated with the Boolean variables. Assume w.l.o.g. that these random variables have domain $\{0, 1\}$. Let ϕ' be the formula ϕ after replacing Boolean variable x_i with $X_i = 0$ and y_i with $Y_i = y_i$.

Consider the probabilistic inequality $\varphi : \sum_{y_1} \dots \sum_{y_n} \mathbb{P}(\phi') \geq 2^{n-1}$ whereby 2^{n-1} is encoded as $\sum_{x_1} \dots \sum_{x_{n-1}} \mathbb{P}(\top)$.

The left hand side is equivalent to

$$\sum_{\hat{x}_1} \dots \sum_{\hat{x}_n} \sum_{\hat{y}_1} \dots \sum_{\hat{y}_n} p_{\hat{x}_1 \dots \hat{x}_n, \hat{y}_1 \dots \hat{y}_n} \sum_{y_1} \dots \sum_{y_n} \delta_{\hat{x}_1 \dots \hat{x}_n, \hat{y}_1 \dots \hat{y}_n}(y_1, \dots, y_n)$$

according to Fact 1.

Since the last sum sums over all values of y_i , it counts the number of satisfying assignments to ϕ given \hat{x}_i . Writing this count as $\#_y(\phi(\hat{x}_1 \dots \hat{x}_n))$, the expression becomes: $\sum_{\hat{x}_1} \dots \sum_{\hat{x}_n} \sum_{\hat{y}_1} \dots \sum_{\hat{y}_n} p_{\hat{x}_1 \dots \hat{x}_n, \hat{y}_1 \dots \hat{y}_n} \#_y(\phi(\hat{x}_1 \dots \hat{x}_n))$.

Since $\#_y(\phi(\hat{x}_1 \dots \hat{x}_n))$ does not depend on \hat{y} , we can write the expression as $\sum_{\hat{x}_1} \dots \sum_{\hat{x}_n} p_{\hat{x}_1 \dots \hat{x}_n} \#_y(\phi(\hat{x}_1 \dots \hat{x}_n))$ whereby $p_{\hat{x}_1 \dots \hat{x}_n} = \sum_{\hat{y}_1} \dots \sum_{\hat{y}_n} p_{\hat{x}_1 \dots \hat{x}_n, \hat{y}_1 \dots \hat{y}_n}$.

If ψ is satisfiable, there is an assignment $\hat{x}_1, \dots, \hat{x}_n$ with $\#_y(\phi(\hat{x}_1 \dots \hat{x}_n)) \geq 2^{n-1}$. If we set $p_{\hat{x}_1 \dots \hat{x}_n} = 1$ and every other probability $p_{\hat{x}'_1 \dots \hat{x}'_n} = 0$, φ is satisfied.

If φ is satisfiable, let $x_1^{\text{max}}, \dots, x_n^{\text{max}}$ be the assignment that maximizes $\#_y(\phi(x_1^{\text{max}} \dots x_n^{\text{max}}))$. Then $2^{n-1} \leq \sum_{\hat{x}_1} \dots \sum_{\hat{x}_n} p_{\hat{x}_1 \dots \hat{x}_n} \#_y(\phi(\hat{x}_1 \dots \hat{x}_n)) \leq \sum_{\hat{x}_1} \dots \sum_{\hat{x}_n} p_{\hat{x}_1 \dots \hat{x}_n} \#_y(\phi(x_1^{\text{max}} \dots x_n^{\text{max}})) = (\sum_{\hat{x}_1} \dots \sum_{\hat{x}_n} p_{\hat{x}_1 \dots \hat{x}_n}) \#_y(\phi(x_1^{\text{max}} \dots x_n^{\text{max}})) = \#_y(\phi(x_1^{\text{max}} \dots x_n^{\text{max}}))$. Hence ψ is satisfied for $x_1^{\text{max}} \dots x_n^{\text{max}}$. \square

Lemma 4. $\text{SAT}_{\mathcal{L}_2}^{\text{lin}(\Sigma)}$ is in PSPACE.

Proof. We need to show that Algorithm 1 is correct and in PSPACE. The basic idea of the algorithm is that rather than guessing a model and evaluating each sum with its interventions⁸, we enumerate all possible interventions (and resulting values) and increment each sum that includes the intervention. Thereby, rather than storing the functions and interventions, we only need to store and update the value of the sums.

By definition, each sum $\sum_{\mathbf{y}_i} \mathbb{P}([\alpha_i]\delta_i)$ in the input can be written as $\sum_{\mathbf{u}} p_{\mathbf{u}} \sum_{\mathbf{y}_i: \mathcal{F}, \mathbf{u} \models [\alpha_i]\delta_i} 1$, where the second sum does not depend on $p_{\mathbf{u}}$. As in Lemma 2, one can write the probabilities as a single vector $\vec{p}_{\mathbf{u}}$, each sum $\sum_{\mathbf{y}_i: \mathcal{F}, \mathbf{u} \models [\alpha_i]\delta_i} 1$ as row in a matrix A , such that an entry of the row is 1 if $\mathcal{F}, \mathbf{u} \models [\alpha_i]\delta_i$ holds and 0 otherwise. Then one obtains the value of all sums as product $A\vec{p}_{\mathbf{u}}$. A small model property follows that there are only polynomial many, rational probabilities $p_{\mathbf{u}}$. These can be guessed nondeterministically⁹.

Next, we combine the terms $\sum_{\mathbf{u}} p_{\mathbf{u}}$ of all sums (implicitly). For example, two sums $\sum_{\mathbf{y}_i} \mathbb{P}([\alpha_i]\delta_i) + \sum_{\mathbf{y}_j} \mathbb{P}([\alpha_j]\delta_j)$ can be rewritten as

$$\begin{aligned} \sum_{\mathbf{y}_i} \mathbb{P}([\alpha_i]\delta_i) + \sum_{\mathbf{y}_j} \mathbb{P}([\alpha_j]\delta_j) &= \sum_{\mathbf{u}} p_{\mathbf{u}} \sum_{\mathbf{y}_i: \mathcal{F}, \mathbf{u} \models [\alpha_i]\delta_i} 1 + \sum_{\mathbf{u}} p_{\mathbf{u}} \sum_{\mathbf{y}_j: \mathcal{F}, \mathbf{u} \models [\alpha_j]\delta_j} 1 \\ &= \sum_{\mathbf{u}} p_{\mathbf{u}} \left(\sum_{\mathbf{y}_i: \mathcal{F}, \mathbf{u} \models [\alpha_i]\delta_i} 1 + \sum_{\mathbf{y}_j: \mathcal{F}, \mathbf{u} \models [\alpha_j]\delta_j} 1 \right). \end{aligned}$$

The algorithm performs this sum over \mathbf{u} in line 5 and calculates the next sums in the subfunction. We can and will ignore the actual values \mathbf{u} . Relevant is only that the value of the sums is multiplied by $p_{\mathbf{u}}$ and that the functions \mathcal{F} might change in each iteration.

⁸For example, a sum like $\sum_x \mathbb{P}([X=x]Y=y)$ performs multiple interventions on X , which is difficult to evaluate. Sums containing only a single intervention could be evaluated trivially.

⁹The algorithm just guesses them without doing any rewriting of the equations. This paragraph just explains why this guessing is possible.

Input: $\text{SAT}_{\mathcal{L}_2}^{\text{lin}(\Sigma)}$ instance
Output: Is the instance satisfiable?

- 1 Guess small model probabilities $p_{\mathbf{u}}$;
- 2 Guess a causal order X_1, \dots, X_n ;
- 3 Rewrite each sum $\sum_{\mathbf{y}_i} \mathbb{P}([\alpha_i] \delta_i)$ in the input as $\sum_{\mathbf{u}} p_{\mathbf{u}} \sum_{\mathbf{y}_i: \mathcal{F}, \mathbf{u} \models [\alpha_i] \delta_i} 1$;
- 4 Initialize a counter c_i to zero for each such sum
- 5 **for** $p_{\mathbf{u}} > 0$ **do**
- 6 Guess values x_1, \dots, x_n ;
- 7 Simulate-Interventions($1, \{\}, x_1, \dots, x_n$)
- 8 **end**
- 9 Replace the sums by c_i and verify whether the (in-)equalities are satisfied;

- 10 **Function** *Simulate-Interventions*($i, \alpha, x_1, \dots, x_n$)
- Input:** Current variable X_i ; set of interventions α ; values x_1, \dots, x_n
- 11 **if** $i > n$ **then**
- 12 **for each** sum counter c_j **do**
- 13 **for all possible values** \mathbf{y}_j **of the sum** **do**
- 14 **if** α_j (after inserting \mathbf{y}_i) is $[\{X_i = x_i \text{ for } i \in \alpha\}]$ and x_1, \dots, x_n satisfy δ_j
 (after inserting \mathbf{y}_i) **then**
- 15 increment c_j by $p_{\mathbf{u}}$
- 16 **end**
- 17 **end**
- 18 **end**
- 19 **else**
- 20 Simulate-Interventions($i + 1, \alpha, x_1, \dots, x_n$) ;
- 21 **for value** v **do**
- 22 Let $x'_1, \dots, x'_n := x_1, \dots, x_n$;
- 23 $x'_i := v$;
- 24 **if** $v \neq x_i$ **then**
- 25 guess new values x'_{i+1}, \dots, x'_n
- 26 **end**
- 27 Simulate-Interventions($i + 1, \alpha \cup \{i\}, x'_1, \dots, x'_n$) ;
- 28 **end**
- 29 **end**

Algorithm 1: Solving $\text{SAT}_{\mathcal{L}_2}^{\text{lin}(\Sigma)}$

Recall that the functions $\mathcal{F} = (F_1, \dots, F_n)$ determine the values of the endogenous variables, that is the value of variable X_i is given by $x_i = F_i(\mathbf{u}, x_1, \dots, x_{i-1})$. Thereby the functions (i.e. variables) have a causal order $X_1 \prec \dots \prec X_n$, such that the value x_i only depends on variables X_j with $j < i$. In reverse, this means that each intervention on a variable X_i can only change variables X_j with $j > i$.

Rather than storing the functions $\mathcal{F} = (F_1, \dots, F_n)$, the algorithm only stores the values $x_i = F_i(\mathbf{u}, x_1, \dots, x_{i-1})$, i.e. the values of the endogenous variables. The algorithm knows these values as well as the causal order, since they can be guessed non-deterministically.

The subfunction Simulate-Interventions then performs all possible interventions recursively, intervening first on variable X_1 , then X_2, \dots , until X_n . The parameter i means an intervention on variable X_i , α is the set of all previous interventions, and x_1, \dots, x_n the current values.

In line 20, it proceeds to the next variable, without changing the current variable X_i (simulating all possible interventions includes intervening on only a subset of variables). In line 21, it enumerates all values x'_i for variable X_i . If $x_i = x'_i$, then the intervention does nothing. If $x_i \neq x'_i$, then the intervention might change all variables X_j with $j > i$ (because the function F_j might depend on X_i and change its value). This is simulated by guessing the new values x'_j . Thereby, we get the new values without considering the functions.

In the last call, line 27, it has completed a set of interventions α . The function then searches every occurrence of the interventions α in the (implicitly expanded) input formula.

That is, for each sum $\sum_{\mathbf{y}_j: \mathcal{F}, \mathbf{u} \models [\alpha_j] \delta_j} 1$, it counts how often $\alpha_j = \alpha$ occurs in the sum while the values x_i satisfy δ_j .¹⁰

Since each intervention is enumerated only once, in the end, it obtains for all sums their exact value. It can then verify whether the values satisfy the (in-)equalities of the input.

If a satisfying model exists, the algorithm confirms it, since it can guess the probabilities and the values of the functions. In reverse, if the algorithm returns true, a satisfying model can be constructed. The probabilities directly give a probability distribution $P(\mathbf{u})$. The functions F_i can be constructed because, for each set of values $\mathbf{u}, x_1, \dots, x_{i-1}$, only a single value for x_i is guessed, which becomes the value of the function.

Algorithm 1 runs in nondeterministical polynomial space and thus in PSPACE. \square

Lemma 5. $\text{SAT}_{\mathcal{L}_2}^{\text{base}(\Sigma)}$ is PSPACE hard.

Proof. We reduce from the canonical PSPACE-complete problem QBF. Let $Q_1 x_1 Q_2 x_2 \dots Q_n x_n \psi$ be a quantified Boolean formula with arbitrary quantifiers $Q_1, \dots, Q_n \in \{\exists, \forall\}$. We introduce Boolean random variables $\mathbf{X} = \{X_1, \dots, X_n\}$ to represent the values of the variables and denote by $\mathbf{Y} = \{Y_1, \dots, Y_k\} \subseteq \mathbf{X}$ the universally quantified variables.

By Lemma ??, we can enforce an ordering $X_1 \prec X_2 \prec \dots \prec X_n$ of variables, i.e. variable X_i can only depend on variables X_j with $j < i$.

Our only further constraint is

$$\sum_{\mathbf{y}} \mathbb{P}([\mathbf{y}] \psi') = 2^k \quad (2)$$

where ψ' is obtained from ψ by replacing positive literals x_i by $X_i = 1$ and negative literals \bar{x}_i by $X_i = 0$.

Suppose the constructed $\text{SAT}_{\mathcal{L}_2}^{\text{lin}(\Sigma)}$ formula is satisfied by a model \mathfrak{M} . We show that $Q_1 x_1 Q_2 x_2 \dots Q_n x_n \psi$ is satisfiable. Each probability implicitly sums over all possible values \mathbf{u} of the exogenous variables. Fix one such \mathbf{u} with positive probability. Combined with $X_1 \prec \dots \prec X_n$ this implies all random variables now deterministically depend only on any of the previous variables. Equation (2) enforces $\mathbb{P}([\mathbf{y}] \psi') = 1$ for every choice of \mathbf{y} and thus simulates the \mathbf{Y} being universally quantified. As the existential variables x_i , we then choose the value x_i of X_i which can only depend on X_j with $j < i$. The formula ψ' and thus ψ is then satisfied due to $\mathbb{P}([\mathbf{y}] \psi') = 1$.

On the other hand, suppose $Q_1 x_1 Q_2 x_2 \dots Q_n x_n \psi$ is satisfiable. We create a deterministic model \mathfrak{M} as follows: The value of existentially quantified variables X_i is then computed by the function $F_i(x_1, \dots, x_{i-1})$ defined as the existentially chosen value when the previous variables are set to x_1, \dots, x_{i-1} . The values of the universally quantified variables do not matter since we intervene on them before every occurrence. This satisfies the required order of variables and since ψ is satisfied we have $\mathbb{P}([\mathbf{y}] \psi') = 1$ for every choice of the universally quantified variables \mathbf{y} , thus satisfying equation (2). \square

Lemma 6. $\text{SAT}_{\mathcal{L}_3}^{\text{base}(\Sigma)}$ and $\text{SAT}_{\mathcal{L}_3}^{\text{lin}(\Sigma)}$ are NEXP-complete.

Proof. We will reduce satisfiability of a Schönfinkel-Bernays sentence to satisfiability of $\text{SAT}_{\mathcal{L}_3}^{\text{base}(\Sigma)}$. The class of Schönfinkel-Bernays sentences (also called Effectively Propositional Logic, EPR) is a fragment of first-order logic formulas where satisfiability is decidable. Each sentence in the class is of the form $\exists \mathbf{x} \forall \mathbf{y} \psi$ whereby ψ can contain logical operations \wedge, \vee, \neg , variables \mathbf{x} and \mathbf{y} , equalities, and relations $R_i(\mathbf{x}, \mathbf{y})$ which depend on a set of variables, but ψ cannot contain any quantifier or functions. Determining whether a Schönfinkel-Bernays sentence is satisfiable is an NEXP-complete problem Lewis [1980] even if all variables are restricted to binary values Achilleos [2015].

We will represent Boolean values as the value of the random variables, with 0 meaning FALSE and 1 meaning TRUE. We will assume that $c = 2$, so that all random variables are binary, i.e. $\text{Val} = \{0, 1\}$.

¹⁰Rather than incrementing the counter c_j by 1 and then multiplying the final result by $p_{\mathbf{u}}$, we increment it directly by $p_{\mathbf{u}}$.

In the proof, we will write (in)equalities between random variables as $=$ and \neq . In the binary setting, $X = Y$ is an abbreviation for $(X = 0 \wedge Y = 0) \vee (X = 1 \wedge Y = 1)$, and $X \neq Y$ an abbreviation for $\neg(X = Y)$. To abbreviate interventions, we will write $[w]$ for $[W = w]$, $[\mathbf{w}]$ for interventions on multiple variables $[\mathbf{W} = \mathbf{w}]$, and $[\mathbf{v} \setminus \mathbf{w}]$ for interventions on all endogenous variables except \mathbf{W} .

We use random variables $\mathbf{X} = \{X_1, \dots, X_n\}$ and $\mathbf{Y} = \{Y_1, \dots, Y_n\}$ for the quantified Boolean variables \mathbf{x}, \mathbf{y} in the sentence $\exists \mathbf{x} \forall \mathbf{y} \psi$. For each distinct k -ary relation $R_i(z_1, \dots, z_k)$ in the formula, we define a random variable R_i and variables Z_i^1, \dots, Z_i^k for the arguments. For the j -th occurrence of that relation $R_i(t_{ij}^1, \dots, t_{ij}^k)$ with $t_{ij}^l \in \{x_1, \dots, x_n, y_1, \dots, y_n\}$, we define another random variable R_i^j .

We use the following constraint to ensure that R_i only depends on its arguments:

$$\sum_{\mathbf{v}} \mathbb{P}([z_i^1, \dots, z_i^k] R_i \neq [\mathbf{v} \setminus r_i] R_i) = 0 \quad (3)$$

Thereby $\sum_{\mathbf{v}}$ refers to summing over all values of all endogenous variables¹¹ in the model and the constraint says that an intervention on Z_i^1, \dots, Z_i^k gives the same result for R_i as an intervention on Z_i^1, \dots, Z_i^k and the remaining variables, excluding R_i .

We use the following constraint to ensure that R_i^j only depends on its arguments:

$$\sum_{\mathbf{v}} \mathbb{P}([t_{ij}^1, \dots, t_{ij}^k] R_i^j \neq [\mathbf{v} \setminus r_i^j] R_i^j) = 0 \quad (4)$$

and that R_i^j and R_i have an equal value for equal arguments:

$$\sum_{t_{ij}^1, \dots, t_{ij}^k} \mathbb{P}([T_{ij}^1 = t_{ij}^1, \dots, T_{ij}^k = t_{ij}^k] R_i^j \neq [Z_i^1 = t_{ij}^1, \dots, Z_i^k = t_{ij}^k] R_i) = 0. \quad (5)$$

We add the following constraint for each X_i to ensure that the values of \mathbf{X} are not affected by the values of \mathbf{Y} :

$$\sum_{\mathbf{v}} \sum_{\mathbf{y}'} \mathbb{P}([\mathbf{v} \setminus x_i] X_i \neq [\mathbf{v} \setminus (x_i, \mathbf{y}), \mathbf{Y} = \mathbf{y}'] X_i) = 0 \quad (6)$$

Here the first sum sums over all values \mathbf{v} of all endogenous variables \mathbf{V} (including X_i and \mathbf{Y}), and the second sum sums over values for variables \mathbf{Y} . The intervention $[\mathbf{v} \setminus x_i]$ intervenes on all variables except X_i and sets the values \mathbf{y} to the values of the first sum. The intervention $[\mathbf{v} \setminus (x_i, \mathbf{y}), \mathbf{Y} = \mathbf{y}']$ intervenes on all variables except X_i and sets the values \mathbf{y} to the values \mathbf{y}' of the second sum. The constraint thus ensures that the value of X_i does not change when changing \mathbf{Y} from \mathbf{y} to \mathbf{y}' .

Let ψ' be obtained from ψ by replacing equality and relations on the Boolean values with the corresponding definitions for the random variables:

$$\sum_{\mathbf{y}} \mathbb{P}([\mathbf{y}] \psi') = 2^n \quad (7)$$

Suppose the $\text{SAT}_{\mathcal{L}_3}^{\text{base}(\Sigma)}$ instance is satisfied by a model \mathfrak{M} . We need to show $\exists \mathbf{x} \forall \mathbf{y} \psi$ is satisfiable. Each probability $\mathbb{P}(\dots)$ implicitly sums over all possible values \mathbf{u} of the exogenous variables. The values \mathbf{x} of the variables \mathbf{X} might change together with the values \mathbf{u} , however, any values \mathbf{x} that are taken at least once can be used to satisfy $\exists \mathbf{x} \forall \mathbf{y} \psi$: If there was any \mathbf{x} that would not satisfy ψ for all values of \mathbf{y} , $\mathbb{P}([\mathbf{y}] \psi')$ would be less than 1 for these values of \mathbf{x} (determined by \mathbf{u}) and \mathbf{y} , and Equation (7) would not be satisfied.

For each relation R_i , we choose the values given by the random variable R_i . Each occurrence $R_i^j(t_{ij}^1, \dots, t_{ij}^k)$ has a value that is given by $[Z_i^1 = t_{ij}^1, \dots, Z_i^k = t_{ij}^k] R_i$ in the model \mathfrak{M} . Due to Equations (5) and Equations (4), that is the same value as $[T_i^1 = t_{ij}^1, \dots, T_i^k = t_{ij}^k] R_i^j$, which is the value used in $[\mathbf{y}] \psi'$. Since $[\mathbf{y}] \psi'$ is satisfied, so is ψ and $\exists \mathbf{x} \forall \mathbf{y} \psi$.

¹¹All variables include variables Z_i^1, \dots, Z_i^k .

Suppose $\exists \mathbf{x} \forall \mathbf{y} \psi$ is satisfiable. We create a deterministic model \mathfrak{M} as follows: The value of random variables \mathbf{X} is set to the values chosen by $\exists \mathbf{x}$. The relation random variables R_i are functions depending on random variables Z_i^1, \dots, Z_i^k that return the value of the relation $R_i(z_1, \dots, z_k)$. The relation random variables R_i^j on arguments $T_{ij}^1, \dots, T_{ij}^k$ return the value of the relation $R_i(t_{ij}^1, \dots, t_{ij}^k)$. This satisfies Equation 3 and 4 because the functions only depends on their arguments, and Equation 5 because the functions result from the same relation (so the functions are dependent, but causally independent, which yields a non-faithful model. But the equations do not test for faithfulness or dependences). All other random variables can be kept constant, which satisfies Equation 6 (despite being constant in the model, the causal interventions can still change their values). Finally, Equation 7 holds, because ψ is satisfied for all \mathbf{Y} .

The problem can be solved in NEXP because expanding all sums of a $\text{SAT}_{\mathcal{L}_3}^{\text{lin}(\Sigma)}$ instance creates a $\text{SAT}_{\mathcal{L}_3}^{\text{lin}}$ instance of exponential size, which can be solved non-deterministically in a time polynomial to the expanded size as shown by Mossé et al. [2022]. \square

Finally, we give the proof of Proposition 1.

Proof of Proposition 1. In the proof of Lemma 2, we require a PP-oracle to evaluate the sum $\sum_{x_{i_1}} \dots \sum_{x_{i_l}} \delta_{\hat{x}_1 \dots \hat{x}_n}(x_{i_1}, \dots, x_{i_l})$, which counts the number of assignments x_{i_1}, \dots, x_{i_l} that make $(X_1 = \hat{x}_1 \wedge \dots \wedge X_n = \hat{x}_n) \rightarrow \delta(x_{i_1}, \dots, x_{i_l})$ a tautology for given \hat{x}_i .

Without negations, there are also no disjunctions, so δ consists of a conjunction of terms that compare some variable to some constant, $X_i = j$, or to some variable $X_i = x_{i_j}$ used in the summation. For any constant $X_i = j$, we check whether $\hat{x}_i = j$. If that is false, the number of satisfying assignments to δ is zero. From any condition $X_i = x_{i_j}$, we learn that x_{i_j} has to be equal to \hat{x}_i . If there is any contradiction, i.e., $X_i = x_{i_j} \wedge X_k = x_{i_j}$ and $\hat{x}_i \neq \hat{x}_k$, the number of assignments is also zero. This determines the value of each x_{i_j} occurring in δ and makes δ a tautology.

It leaves the value of x_{i_j} not occurring in δ undefined, but those do not affect δ , and can be chosen arbitrarily, so the number of assignments is just multiplied by the size of the domain c for each not-occurring variable.

This can be evaluated in polynomial time, so the complexity is reduced to $\text{NP}^P = \text{NP}$.

The problems remain NP-hard, since a Boolean formula can be encoded in the language $\mathcal{L}_1^{\text{base}}$ of combinations of Boolean inequalities. Each Boolean variable x is replaced by $P(X) > 0$ for a corresponding random variable X . For example, a 3-SAT instance like $(x_{i_{1,1}} \vee \neg x_{i_{1,2}} \vee x_{i_{1,3}}) \wedge (x_{i_{2,1}} \vee \neg x_{i_{2,2}} \vee \neg x_{i_{2,3}}) \wedge \dots$ can be encoded as $(P(X_{i_{1,1}}) > 0 \vee \neg P(X_{i_{1,2}}) > 0 \vee P(X_{i_{1,3}}) > 0) \wedge (P(X_{i_{2,1}}) > 0 \vee \neg P(X_{i_{2,2}}) > 0 \vee \neg P(X_{i_{2,3}}) > 0) \wedge \dots$, which clearly has the same satisfiability. \square

B.2 Proofs of Section 3.2

Mossé et al. [2022] already prove $\text{SAT}_{\mathcal{L}_3}^{\text{poly}} \in \exists \mathbb{R}$ when the $\mathcal{L}_3^{\text{poly}}$ -formula is allowed to neither contain subtractions nor conditional probabilities. We slightly strengthen this result to allow both of them.

Lemma 7. $\text{SAT}_{\mathcal{L}_3}^{\text{poly}} \in \exists \mathbb{R}$. *This also holds true if we allow the basic terms to contain conditional probabilities.*

Proof. [Mossé et al., 2022] show that $\text{SAT}_{\mathcal{L}_3}^{\text{poly}}$ without subtraction or conditional probabilities is in $\exists \mathbb{R}$. Their algorithm is given in the form of a NP-reduction from $\text{SAT}_{\mathcal{L}_3}^{\text{poly}}$ to ETR and using the closure of $\exists \mathbb{R}$ under NP-reductions. In particular given a $\mathcal{L}_3^{\text{poly}}$ -formula φ , they replace each event $\mathbb{P}(\epsilon)$ by the sum $\sum_{\delta \in \Delta^+ : \delta \models \epsilon} \mathbb{P}(\delta)$ where $\Delta^+ \subseteq \mathcal{E}_{\text{counterfact}}$ is a subset of size at most $|\varphi|$. They add the constraint $\sum_{\delta \in \Delta^+} \mathbb{P}(\delta) = 1$ and then replace each of the $\mathbb{P}(\delta)$ by a variable constrained to be between 0 and 1 to obtain a ETR-formula. Note that the final ETR-formula allows for subtraction, so φ is allowed to have subtractions as well. Remains to show how to deal with conditional probabilities. We define conditional probabilities $\mathbb{P}(\delta|\delta')$ to be undefined if $\mathbb{P}(\delta') = 0$ (this proof works similarly for other definitions). In φ replace $\mathbb{P}(\delta|\delta')$ by $\frac{\mathbb{P}(\delta)}{\mathbb{P}(\delta')}$. The resulting ETR-formula then contains some divisions. To remove some

division $\frac{\alpha}{\beta}$, we use Tsaitin's trick and replace $\frac{\alpha}{\beta}$ by a fresh variable z . We then add the constraints $\alpha = z \cdot \beta$ and $\beta \neq 0$ to the formula. \square

Now we are ready to show that $\text{SAT}_{\mathcal{L}_3}^{\text{poly}(\Sigma)}$ can be solved in NEXP over the Reals.

Proof of Lemma 1. By Lemma 7 we know that $\text{SAT}_{\mathcal{L}_3}^{\text{poly}}$ is in $\exists\mathbb{R}$. Thus, by [Erickson et al., 2022], there exists an NP_{real} algorithm, call it A , which for a given $\mathcal{L}_3^{\text{poly}}$ -formula decides if is satisfiable.

To solve the $\text{SAT}_{\mathcal{L}_3}^{\text{poly}(\Sigma)}$ problem, a $\text{NEXP}_{\text{real}}$ algorithm expands firstly all sums of a given instance and creates an equivalent $\text{SAT}_{\mathcal{L}_3}^{\text{poly}}$ instance of size bounded exponentially in the size of the initial input formula. Next, the algorithm A is used to decide if the expanded instance is satisfiable. \square

B.3 Proofs of Section 4

Input: $\mathcal{L}_3^{\text{poly}(\Sigma)}$ formula φ and a unary number $p \in \mathbb{N}$

Output: Is the instance satisfiable with a model of size at most p ?

- 1 Guess a causal order X_1, \dots, X_n ;
- 2 Explicitly expand all exponential sums in φ ;
- 3 For each probability $\mathbb{P}(\delta_i)$ occurring in φ introduce a variable P_i , initialized to 0;
- 4 Introduce symbolic variables z_1, \dots, z_p ;
- 5 **for** $i \in \{1, \dots, p\}$ **do**
- 6 Guess a deterministic SCM \mathfrak{M} using the causal order X_1, \dots, X_n . **for each** $\mathbb{P}(\delta_i)$ **that**
 is satisfied by \mathfrak{M} **do**
- 7 Symbolically increment P_j by z_i .
- 8 **end**
- 9 **end**
- 10 Replace all probabilities $\mathbb{P}(\delta_i)$ by the values of P_i and check whether the resulting ETR-formula (in the variables z_1, \dots, z_p) is satisfiable using Renegar's algorithm;

Algorithm 2: Solving $\text{SAT}_{sm, \mathcal{L}_3}^{\text{poly}(\Sigma)}$

We prove the NEXP-completeness of the interventional and counterfactual satisfiability problems with small model property as follows.

Proof of Theorem 3. We show that $\text{SAT}_{sm, \mathcal{L}_2}^{\text{poly}(\Sigma)}$ is NEXP hard. This is done using a variation of the reduction from Lemma 6. We again reduce from the satisfiability of Schönfinkel-Barnay sentences. The random variables are defined in the same way as in Lemma 6. Using Lemma ??, we require that each model has the variable ordering $X_1 \prec \dots \prec X_n \prec Y_1 \prec \dots \prec Y_n$ and then the variables R_i and R_i^j in some arbitrary, but fixed order. This allows to enforce a deterministic model by adding the constraint $\sum_{\mathbf{t}} \sum_{\mathbf{v}} (\mathbb{P}([\mathbf{t}|\mathbf{v}])^2 - \mathbb{P}([\mathbf{t}|\mathbf{v}]))^2 = 0$ for each subset T of variables containing a (possibly empty) consecutive subset of variables starting at X_1 in our variable ordering. This enforces every probability in this model to be either 0 or 1 and further enforces that no variable depends on any exogenous variables, but only the endogenous variables. This determinism allows us to rephrase the additional equations from Lemma 6 using purely interventional terms:

We use the following constraint to ensure that R_i only depends on its arguments:

$$\sum_{\mathbf{v}} (\mathbb{P}([z_i^1, \dots, z_i^k] R_i = r_i) - \mathbb{P}([\mathbf{v} \setminus r_i] R_i = r_i))^2 = 0 \quad (8)$$

Thereby $\sum_{\mathbf{v}}$ refers to summing over all values of all endogenous variables in the model and the constraint says that an intervention on Z_i^1, \dots, Z_i^k gives the same result for R_i as an intervention on Z_i^1, \dots, Z_i^k and the remaining variables, excluding R_i .

We use the following constraint to ensure that R_i^j only depends on its arguments:

$$\sum_{\mathbf{v}} (\mathbb{P}([t_1, \dots, t_k] R_i^j = r_i^j) - \mathbb{P}([\mathbf{v} \setminus r_i^j] R_i^j = r_i^j))^2 = 0 \quad (9)$$

and that R_i^j and R_i have an equal value for equal arguments:

$$\sum_{t_1, \dots, t_k} (\mathbb{P}([T_1=t_1, \dots, T_k=t_k]R_i^j=r_i) - \mathbb{P}([Z_i^1=t_1, \dots, Z_i^k=t_k]R_i=r_i))^2 = 0. \quad (10)$$

We add the following constraint for each X_i to ensure that the values of \mathbf{X} are not affected by the values of \mathbf{Y} :

$$\sum_{\mathbf{v}} \sum_{\mathbf{y}'} (\mathbb{P}([\mathbf{v} \setminus x_i]X_i=x_i) - \mathbb{P}([\mathbf{v} \setminus (x_i, \mathbf{y}), \mathbf{Y}=\mathbf{y}']X_i=x_i))^2 = 0 \quad (11)$$

Let ψ' be obtained from ψ by replacing equality and relations on the Boolean values with the corresponding definitions for the random variables:

$$\sum_{\mathbf{y}} \mathbb{P}([\mathbf{y}]\psi') = 2^n \quad (12)$$

The correctness now follows in the same way as Lemma 6. On one hand, the model \mathfrak{M} constructed there if $\exists \mathbf{x} \forall \mathbf{y} \psi$ is satisfiable is already deterministic and thus fulfils all the above constraints. On the other hand if there is a model \mathfrak{M} satisfying the above constraints, all possible values of \mathbf{u} of the exogenous variables result in the exact same probabilities which, following the original proof, lead to $\exists \mathbf{x} \forall \mathbf{y} \psi$ to be satisfiable.

Furthermore $\text{SAT}_{sm, \mathcal{L}_3}^{\text{poly}(\Sigma)} \in \text{NEXP}$. We use the same trick from Lemma 4 and instead of viewing the model as changing with \mathbf{u} , we instead consider for each value \mathbf{u} a separate (deterministic) model. Due to the small-model property we then only have to consider polynomially many models and each model has a straight-forward representation using exponentially many bits, directly encoding all the functions in \mathcal{F} as explicit tables. Algorithm 2 uses this observation to solve $\text{SAT}_{sm, \mathcal{L}_3}^{\text{poly}(\Sigma)}$ nondeterministically in exponential time. It constructs an ETR-formula with polynomially many variables, polynomially many polynomials of polynomial degree, but potentially exponentially many monomials. Each coefficient of the formula can be represented using a polynomial bit length. Renegar's algorithm (Renegar [1992]) can thus solve the ETR-formula deterministically in time $O(\text{poly}(|\varphi|)^{O(p)})$ which is exponential in the input length. \square