A CLASS OF EXPLICIT SOLUTIONS FOR THE FERMAT PROBLEM FOR TETRAHEDRA

ANASTASIOS N. ZACHOS

ABSTRACT. We present a class of explicit solutions for the problem of minimization of the function $f(x, y, z) = \sum_{i=1}^{4} \sqrt{(x - x_i)^2 + (y - y_i)^2 + (z - z_i)^2}$, which gives the location of the unique stationary (Fermat-Torricelli) point for four non-collinear and non-coplanar points $A_i = (x_i, y_i, z_i)$, determining tetrahedra, which are derived by a proper class of isosceles tetrahedra having four equal edges and two equal opposite edges. This class of explicit solutions contains Mehlhos and Glastier's explicit solutions (theoretical constructions) obtained in [23] and [15], respectively.

1. INTRODUCTION

In 1643, Fermat posed an optimization problem, which has not been studied by Ancient Greeks. The problem states as follows: Given three points in the Euclidean plane, find a point having the minimum sum of distances from these three points.

The solution of the Fermat problem is called the Fermat point. There are two types of solutions for the Fermat problem in the Euclidean plane.

First type. The corner angles of the triangle formed by these three fixed points are less than 120 degrees. This type was discovered by Torricelli (a student of Gallileo) and it is called Torricelli solution. The Torricelli solution refers to the Fermat-Torricelli point, which is strictly inside the triangle and its side of the triangle is seen by 120 degrees. This is the isogonal (equiangular) property of the Fermat-Torricelli point. The Fermat Torricelli point is the intersection point of three circles circumscribed around regular triangles located outside on each side of the triangle. Hence, the Fermat-Torricelli point is constructed for three noncollinear points in \mathbb{R}^2 using ruler and compass (Euclidean construction).

Second type. One of the corner angles of the triangle is greater or equal to 120 degrees. This type was discovered by Cavallieri and it is called Cavallieri solution. The Cavallieri solution refers to the Fermat-Cavalieri point, which is the vertex of the obtuse corner angle of the triangle.

Extending the Fermat problem for four non-collinear points in \mathbb{R}^2 , we consider the following two cases:

Case 1 If these four points form a convex quadrilateral, then the Fermat (Torricelli) point is the intersection of the two diagonals, This result was proved by Fagnano.

Case 2 If these four points form a non-convex quadrilateral, then the Fermat (Cavallieri) point is the vertex of the non-convex angle. This result is easily derived by using triangle inequalities.

²⁰¹⁰ Mathematics Subject Classification. Primary 51M14,51M20; Secondary 51M16. Key words and phrases. Fermat problem, Fermat-Torricelli point, tetrahedra.

ANASTASIOS N. ZACHOS

The unsolvability of the Fermat problem by Euclidean constructions using ruler and compass for five points in \mathbb{R}^2 has been proved by Cockayne and Melzak in [5] by applying Galois theory in a specific example $P_1P_2P_3P_4P_5$ for $P_1 = (0,0)$, $P_2 = (0,-1)$, $P_3 = (0,1)$, $P_4 = (a,b)$, $P_5 = (a,-b)$. They derive an eighth degree polynomial equation with respect to the distance P_1P_{min} , where P_{min} is the Fermat-Torricelli point. The coefficients of these polynomials depend on the integers a, b. They observed that the eight degree polynomial equation contains a Galois group over the field of rationals, which does not have order 2^k , for k a positive integer. Hence, this equation is not solvable by radicals. Therefore, P_1P_{min} is not constructible using ruler and compass. Furthermore, the unsolvability of Fermat problem by Euclidean constructions has been proved for five points in \mathbb{R}^2 using different examples by Bajaj in [5] and by Mehlhos in [23].

Let $A_1A_2A_3A_4$ be a tetrahedron and let $A_i = (x_i, y_i, z_i)$, i = 1, 2, 3, 4. Then the Fermat's Problem states as follow:

Problem 1. Find (x, y, z) in \mathbb{R}^3 , that minimizes:

$$f(x, y, z) = \sum_{i=1}^{4} \sqrt{(x - x_i)^2 + (y - y_i)^2 + (z - z_i)^2}.$$
 (1.1)

Definition 1. The solution to Problem 1 is called the Fermat point of a tetrahedron $A_1A_2A_3A_4$ and it is denoted by A_0 .

The two types of solutions for the Fermat Problem (Problem 1) are:

Case 1 A_0 is strictly inside of $A_1A_2A_3A_4$. A_0 is called the Fermat-Torricelli point and it is a generalization of Torricelli's solution in \mathbb{R}^3 .

Case 2 A_0 coincides with one of the vertices $\{A_1, A_2, A_3, A_4\}$. A_0 is called the Fermat-Cavallieri point and it is a generalization of Cavallieri'solution in \mathbb{R}^3 .

It is worth mentioning that Synge in correspondence with Coxeter were the first who suggested to call a generalization of solution of Case 1 for a finite number of given points in \mathbb{R}^n $(n \ge 2)$ Torricelli-Fermat point or Fermat-Torricelli point ([30, p. 2]).

There are many references to the Fermat problem and called by different names. The Fermat-Weber problem refers to the Fermat problem for the location of industries and created a new science that is called "Location Science" ([37]). The optimum location for the production of a good based on the fixed locations of the market and a finite number of raw material sources, which do not belong to the same line and determine the least-cost production location by computing the total costs of transporting raw materials from these sites to the production site and product from the production site to the market.

The Fermat-Steiner problem also refers to the tree solution for three non-collinear points to the plane, which was suggested by Courant and Robbins in [8] and adopted by Gueron and Tessler in [14] and has been applied by many researchers in the fields of Mathematical Chemistry and Mathematical Biology and especially in the geometric folding problem of proteins and aminoacids ([26]).

We adopt Synge's Coxeter suggestion for the Fermat solution, which is the Fermat-Torricelli point for case 1 and use the Fermat-Cavalleri point for the case 2.

In 2000, Mehlhos gave an elegant proof for the unsolvability of the Fermat problem by considering for four non-coplanar and non-collinear points $A_1 = (0, -1, 0)$, $A_2 = (0, 1, 0), A_3 = (1, 0, 0), A_4 = (1, 0, 1)$ in \mathbb{R}^3 without using Galois theory([23,

Theorem 2,pp. 153-155]). He observed that the Fermat-Torricelli point $A_0 =$ (x,0,z) belongs to the x-z-plane and by turning $\triangle A_1A_0A_2$ by 90° in the x - z-plane A_0 is the intersection point of the two diagonals of the new derived convex quadrilateral $A'_1A'_2A_3A_4$. The angle of rotation is called the twist angle and it is very useful for the computation of the Fermat-Torricelli point. Thus, by applying Fagnano's result A_0 is the Fermat-Torricelli point of $A'_1A'_2A_3A_4$. By using as variable the angle $\varphi = \angle A_3 O x$ and by turning the figure with the angle φ such that $A'_1A'_2$ belong to the z-axis and by substituting $\sin \varphi = x$ and $\cos \varphi = \sqrt{1-x^2}$ to the system of equations of the two diagonals he derived the quartic equation $8x^4 - 4x^3 - 7x^2 + 2x + 1 = 0$, which gives $8x^3 + 4x^2 - 3x - 1 = 0$, for $x \neq 1$. Thus, we are able to use Cardano-Ferrari's formulas, in order to find the solution of this cubic equation. Then, by substituting $x = t - \frac{1}{6}$, we get $t^3 - \frac{11}{24}t - \frac{23}{432} = 0$ or $432t^3 - 198t - 23 = 0$. Taking into account the rational root theorem, this equation does not have rational roots, but three real roots (one positive and two negative), because the determinant $D = 4(-198)^3 + 27(-23)^2 = -31035285 < 0$. Hence, the positive real solution consists of a cubic root, which does not give a rational number. Therefore, $\sin \varphi$ is not constructive, which yields that $A_0 = \left(\frac{\cos^2 \varphi}{1 - \sin \varphi}, 0, \frac{\cos \varphi \sin \varphi}{1 - \sin \varphi}\right)$ cannot be constructed using ruler and compass. We consider Mehlhos approach as the first example that deals with an explicit solution of the Fermat Problem for a tetrahedron such that four edges of the tetrahedron are seen by four equal angles and the remaining two opposite (non-neighboring) edges are seen by two equal angles.

In 2014, Uteshev succeeded in finding an explicit analytical solution for the Fermat problem for a triangle in \mathbb{R}^2 and managed to express the coordinates of the Fermat-Torricelli point w.r to the coordinates of the three fixed points A_1, A_2, A_3 . The key idea is that he used duality in the Fermat problem and introduced a dual Fermat-Torricelli problem even for unequal positive numbers (weights), that correspond to the vertices of the triangle ([33]). The value of the objective function $f(x, y) = \sum_{i=1}^{4} b_i \sqrt{(x - x_i)^2 + (y - y_i)^2}$ does not change if the weights b_1, b_2, b_3 become distances (side lengths of a dual triangle) and the distances $\sqrt{(x - x_i)^2 + (y - y_i)^2}$ become weights, for i = 1, 2, 3. Unfortunately, the notion of duality does not seem to work for the Fermat problem for tetrahedra, because the Fermat-Torricelli problem is in general not constructible using ruler and compass, even for the unweighted case.

We consider six main attempts to find the Fermat-Torricelli point for tetrahedra in \mathbb{R}^3 :

(i) Sturm's method of intersection of algebraic surfaces of order higher than two ([29]) at the Fermat-Torricelli point. Egervary showed that the order of algebraic surfaces can be reduced to second order ([11]).

(ii) Weiszfeld introduced an algorithm, which gives a fixed point iteration method by solving the system of the three equations $\frac{\partial f}{\partial x} = 0$, $\frac{\partial f}{\partial y} = 0$, $\frac{\partial f}{\partial z} = 0$, of the objective function (1.1) w.r to x, y, z ([35], [36]). Kuhn proved the convergence of Weiszfeld algorithm in [18].

(iii) Synge's dynamic approach was given in [30]. Synge ([30]) was the first who used a dynamic process that contains an infinite number of steps (not a Euclidean construction) for the Fermat-Torricelli point using "spindles" (spherical segments). He considered around the two opposite edges of the tetrahedron and created two isosceles triangles containing two angles $\pi - \angle A_1 A_0 A_2$ and $\pi - \angle A_3 A_0 A_4$ and by

rotating two circular arcs having as chords the edges $A_1A_2 A_3A_4$, he showed that there is a common value such that $\angle A_1A_0A_2 = \angle A_1A_0A_2$, which yields a unique touching point (Fermat-Torricelli point) of the two "spindles."

(iv) An ϵ approximation method of ovals (m-ellipsoids) back to Descartes. An ellipsoid is a triaxial quadratic surface, which is given in Cartesian coordinates by the equation $(\frac{x}{a})^2 + (\frac{y}{b})^2 + (\frac{z}{c})^2 = 1$ Straud generalized the "thread" construction for an ellipsoid analogous to the taut pencil and string construction of the ellipse ([16, pp. 19-22]), with respect to two fixed confocal points. In a private letter (1638) Descartes invited Fermat to investigate properties of ellipses with four confocal points. These "ellipses" are called 'egglipses' or Descartes ovals or multiconics ([10]). The concentration of tetrafocal ellipsoids to a specific value for the objective functions f(x, y, z) lead to the unique Fermat-Torricelli point. In general, the Fermat problem is an NP hard problem to compute the Fermat-Torricelli point. Chandrasekaran and Tamir invented an ellipsoid method, which leads to a polynomial method to construct an approximate solution for the Fermat problem of fixed accuracy in \mathbb{R}^n . Thus, the Fermat problem may be solved in polynomial time ([7]).

(v) Explicit solutions of Glastier, Kupitz-Martini, Mehlhos. Let $A_1 = (0, -1, 0)$, $A_2 = (0, 1, 0)$, $A_3 = (1, 0, 0)$, $A_4 = (1, 0, 1)$ be a tetrahedron in \mathbb{R}^3 .

First type of explicit solution (Regular tetrahedra) In 1993, Glastier proved that the centroid G and the Fermat-Torricelli point A_0 of a regular tetrahedron $A_1A_2A_3A_4$ coincide, because the following balancing condition of unit vectors hold ([15]):

$$\frac{\vec{XA_1}}{|XA_1|} + \frac{\vec{XA_2}}{|XA_2|} + \frac{\vec{XA_3}}{|XA_3|} + \frac{\vec{XA_4}}{|XA_4|} = 0,$$

for X = G and $X = A_0$. By squaring both sides of the equation, the equiangular property of the Fermat-Torricelli point for regular tetrahedra is obtained:

$$\angle A_i A_0 A_j = \cos^{-1}(-1/3) \approx 109^{\circ}28',$$

for $i, j = 1, 2, 3, 4, i \neq j$. Furthermore, he verified a result well-known in chemistry regarding the molecular structure of methane CH_4 , by observing that the six bond angles HCH in the methane molecule CH_4 are equal to $\approx 109^{\circ}28'$ and by applying the Fermat solution for a regular tetrahedron ([15]). The four positions of hydrogen H yield a regular tetrahedron and the carbon C is located at the corresponding Fermat-Torricelli point.

Second type of explicit solution (Isosceles tetrahedra) A tetrahedron is called isosceles if all its two faces (triangles) are congruent. The centroid (barycenter) of an isosceles tetrahedron coincides with the Fermat-Torricelli point and with the center of the circumscribed sphere. Various characterizations for isosceles tetrahedra are given by Bogatyi in Arnold's Problems ([4, pp. 188-192]) and by Kupitz-Martini ([19]).

Third type of explicit solution (Glastier) Let $OA_1A_2A_3$ be a tetrahedron, such that $O = (0, 0, 0), A_1 = (1, 0, 0), A_2 = (0, 1, 0), A_3 = (0, 0, 1)$. The Fermat-Torricelli point can be assumed to be placed at $A_0 = (x, x, x)$. Thus, the objective function f(x, y, z) takes the form $f(x, x, x) = \sqrt{3}x + 3\sqrt{2x^2 + (x - 1)^2}$. By solving $\frac{\partial f}{\partial x} = 0$, we get $12x^2 - 8x + 1 = 0$, which gives $x = \frac{1}{6}$. We note that the Fermat-Torricelli point $A_0 = (\frac{1}{6}, \frac{1}{6}, \frac{1}{6})$ possesses the equiangular property $\angle A_i A_0 A_j = \arccos(-\frac{1}{3})$, for $i, j = 1, 2, 3, 4, i \neq j$. An alternative proof is to consider a point $A_5 = (-\frac{1}{3}, -\frac{1}{3}, -\frac{1}{3})$ that lies on the ray formed by A_0O . Hence, $A_5A_1A_2A_3$ is a regular tetrahedron,

which satisfies the equation

$$\frac{\vec{XA_1}}{|XA_1|} + \frac{\vec{XA_2}}{|XA_2|} + \frac{\vec{XA_3}}{|XA_3|} + \frac{\vec{XA_5}}{|XA_5|} = 0$$
$$\frac{\vec{XA_1}}{|XA_1|} + \frac{\vec{XA_2}}{|XA_2|} + \frac{\vec{XA_3}}{|XA_3|} + \frac{\vec{XO}}{|XO|} = 0.$$

Hence, we get $X = A_0$.

or

Fourth type of explicit solution (Mehlhos) Let $A_1 = (0, -1, 0)$, $A_2 = (0, 1, 0)$, $A_3 = (1, 0, 0)$, $A_4 = (1, 0, 1)$ be a tetrahedron in \mathbb{R}^3 . By following the same process that was used by Mehlos to show the unsolvability of the Fermat problem for tetrahedra with ruler and compass, Mehlhos derived the following explicit solution

$$A_0 = \left(\frac{\cos^2\varphi}{1-\sin\varphi}, 0, \frac{\cos\varphi\sin\varphi}{1-\sin\varphi}\right),\,$$

where $\varphi = \angle A_3 O x$. This is the first explicit solution for the Fermat-Torricelli problem, such that the equiangular property of the Fermat-Torricelli point does not hold for an non-regular tetrahedron. The non-isogonal property of the Fermat-Torricelli point for tetrahedra has been possibly known to Sturm and Lindelof ([29],[28],[22]) showed by Synge in [30] and proved by Abu-Abas, Abu-Saymeh and Hajja for higher dimensional simplexes in [1] and [2]).

(vi) Characterizations of the solutions for the Fermat problem for tetrahedra. Sturm, Kupitz-Martini, Eriksson, Noda, Sakai, Morimoto derived various characterizations for the Fermat-Torricelli point for tetrahedra in \mathbb{R}^3 ([29], [19][13], [25]).

The existence of the Fermat-Torricelli point A_0 in \mathbb{R}^3 is easily derived by a well known theorem of Weirstrass ([31, Corollary, p 111]):

If the function f(x, y, z) is continuous for every $(x, y, z) \in \mathbb{R}^3$ and $\lim f(x, y, z) = \infty$, for $x^2 + y^2 + z^2 \to \infty$, then the unconstrained problem is solvable. Therefore, the objective function f(x, y, z) is solvable and a solution (Fermat point) exists. The uniqueness of the Fermat point is obtained by the convexity of the Euclidean norm (distance). A complete proof for the convexity of distances in hyperbolic spaces is given by Thurston in his classical book [32, Theorem 2.5.8.pp. 90-94]. The proof is the same in \mathbb{R}^n ([32, Exercise 2.5.13 Case(b).p. 94]).

Sturm and Lindelof were the first, who gave a complete characterization of the solutions of the Fermat-Torricelli problem for m given points in \mathbb{R}^n ([28],[22]).

We need the following three results, which deal with the uniqueness and the characterization of solutions in the three dimensional case ([32, Theorem 2.5.8.pp. 90-94], [6, Theorem 18.3, pp. 237],[27, Theorem, p. 863], [23, Property 3, p. 154] or [30, Formulas (6,2), (6.3), p. 8]).

Let A, B two points in \mathbb{R}^3 .

Theorem 1. (Thurston's strict convexity of Euclidean distance functions) The distance function d(A, B) considered as a map $d : \mathbb{R}^3 \times \mathbb{R}^3 \to \mathbb{R}$, is convex. The composition $d \circ \gamma$ is strictly convex for any line γ in $\mathbb{R}^3 \times \mathbb{R}^3$, whose projections to the two factors are distinct lines.

We denote by $\vec{u}(j,i) \equiv \frac{\overrightarrow{A_jA_i}}{\|A_jA_i\|}$ the unit vector from A_j to A_i for i, j = 0, 1, 2, 3, 4.

Theorem 2. (I) If $\|\sum_{j=1, j\neq i}^{4} \vec{u}(j, i)\| > 1$, for each i = 1, 2, 3, 4, then (a) A_0 is strictly inside of the tetrahedron $A_1A_2A_3A_4$,

ANASTASIOS N. ZACHOS

 $\begin{array}{l} (b) \sum_{i=1}^{4} \vec{u}(0,i) = \vec{0} \\ (Fermat-Torricelli \ solution). \\ (II) \ If \|\sum_{j=1, j \neq i}^{4} \vec{u}(j,i)\| \leq 1 \ for \ some \ i = 1,2,3,4, \ then \ A_0 = A_i. \\ (Fermat-Cavallieri \ solution). \end{array}$

Theorem 3. (Spira-Synge's relations) The following relations hold:

$$\cos \angle A_1 A_0 A_2 = \cos \angle A_3 A_0 A_4, \tag{1.2}$$

$$\cos \angle A_2 A_0 A_3 = \cos \angle A_1 A_0 A_4, \tag{1.3}$$

$$\cos \angle A_1 A_0 A_3 = \cos \angle A_2 A_0 A_4 \tag{1.4}$$

and

$$1 + \cos \angle A_1 A_0 A_2 + \cos \angle A_1 A_0 A_3 + \cos \angle A_2 A_0 A_4 = 0. \tag{1.5}$$

In this paper, we focus on the fifth approach and we introduce a class of explicit solutions for the Fermat problem for tetrahedra in \mathbb{R}^3 , which contains the explicit solutions given by Mehlhos and Glastier.

2. Explicit solution for the Fermat problem for isosceles tetrahedra and almost platonic tetrahedra

We start by giving three definitions for three types of tetrahedra.

Definition 2. A Platonic tetrahedron is a regular polyhedron in which all edges are equal.

Definition 3. An isosceles tetrahedron is a non-regular polyhedron in which each pair of opposite edges are equal.

Definition 4. An almost Platonic tetrahedron is an isosceles tetrahedron having four equal edges and a pair of equal opposite edges.

Let $A_1A_2A_3A_4$ be an isosceles tetrahedron and let A_0 be the Fermat-Torricelli point. We denote by S(O; R) the circumscribed sphere with center O and radius R, where $A_i \in S(O; R)$ for i = 1, 2, 3, 4.

We denote by P_{jk}^i the orthogonal projection of A_i to the plane defined by $\triangle A_i O A_k$.

We denote by $\angle(ij) \equiv \angle A_i O A_j$. We denote by $\angle(i, jk) \equiv \angle A_i O P_{jk}^i$, for i, j, k = 1, 2, 3, 4. We denote by $\omega_{jk}^i = \angle P_{jk}^i O A_1$. We set $A_1 O = A_2 O = A_3 O = A_4 O = R$.

Proposition 1. If $A_1A_2 = A_3A_4 = a$, $A_1A_3 = A_2A_4 = b$, $A_1A_4 = A_2A_3 = c$, then $A_0 = O$.

Proof. By using the cosine law in $\triangle A_1 O A_2$ and $\triangle A_3 O A_4$, we get

$$a^2 = 2R^2 - 2R^2 \cos \angle (12) \tag{2.1}$$

and

$$a^2 = 2R^2 - 2R^2 \cos \angle (34). \tag{2.2}$$

By subtracting (2.2) from (2.1), we get $\angle(12) = \angle(34)$. By using the cosine law in $\triangle A_1OA_3$ and $\triangle A_2OA_4$, we get

$$b^2 = 2R^2 - 2R^2 \cos \angle (13) \tag{2.3}$$

and

$$b^2 = 2R^2 - 2R^2 \cos \angle (24). \tag{2.4}$$

By subtracting (2.3) from (2.4), we get \angle (13) = \angle (24). By using the cosine law in $\triangle A_2OA_3$ and $\triangle A_1OA_4$, we get

$$c^2 = 2R^2 - 2R^2 \cos \angle (23) \tag{2.5}$$

and

$$c^2 = 2R^2 - 2R^2 \cos \angle (14). \tag{2.6}$$

By subtracting (2.5) from (2.6), we get \angle (23) = \angle (14).

Without loss of generality, we express the unit vectors $\vec{u}(0, i)$ for i = 1, 2, 3, 4, in the following form:

$$\vec{u}(0,1) = (1,0,0),$$
 (2.7)

$$\vec{u}(0,2) = (\cos \angle (12), \sin \angle (12), 0),$$
(2.8)

$$\vec{u}(0,3) = (\cos \angle (3,12) \cos \omega_{12}^3, \cos \angle (3,12) \sin \omega_{12}^3, \sin \angle (3,12)), \qquad (2.9)$$

$$\vec{u}(0,4) = (\cos \angle (4,12) \cos \omega_{12}^4, \cos \angle (4,12) \sin \omega_{12}^4, \sin \angle (4,12)).$$
(2.10)

By taking the inner products $\vec{u}(0,1) \cdot \vec{u}(0,3)$, $\vec{u}(0,1) \cdot \vec{u}(0,4)$, $\vec{u}(0,2) \cdot \vec{u}(0,3)$, $\vec{u}(0,3) \cdot \vec{u}(0,3)$, we get:

$$\cos \angle (3, 12) \cos \omega_{12}^3 = \cos \angle (13),$$
 (2.11)

$$\cos \angle (4, 12) \cos \omega_{12}^4 = \cos \angle (14),$$
 (2.12)

$$\cos \angle (12) \cos \angle (3,12) \cos \omega_{12}^3 + \sin \angle (12) \cos \angle (3,12) \sin \omega_{12}^3 = \cos \angle (23), \quad (2.13)$$

$$\cos \angle (12) \cos \angle (4, 12) \cos \omega_{12}^4 + \sin \angle (12) \cos \angle (4, 12) \sin \omega_{12}^4 = \cos \angle (24), \quad (2.14)$$

$$\cos \angle (3,12) \cos \omega_{12}^3 \cos \angle (4,12) \cos \omega_{12}^4 + \\ \cos \angle (3,12) \sin \omega_{12}^3 \cos \angle (4,12) \sin \omega_{12}^4 + \sin \angle (3,12) \sin \angle (4,12) = \cos \angle (34).(2.15)$$

By substituting (2.11) in (2.13), solving w.r. to $\cos \angle (3, 12) \sin \omega_{12}^3$ and by squaring both parts of the derived equation and (2.11), respectively and by adding these two equations, we eliminate ω_{12}^3 and by setting $u = \cos \angle (12)$, $v = \cos \angle (23)$, $w = \cos \angle (13)$ in the derived equation, we obtain:

$$\cos^2(3,12) = \frac{v^2 + w^2 - 2uvw}{1 - u^2},\tag{2.16}$$

Similarly, by substituting (2.12) in (2.14), solving w.r. to $\cos \angle (4, 12) \sin \omega_{12}^4$ and by squaring both parts of the derived equation and (2.12), respectively and by adding these two equations we eliminate ω_{12}^4 and by setting $u = \cos \angle (12)$, $v = \cos \angle (23)$, $w = \cos \angle (13)$ in the derived equation, we obtain:

$$\cos^2(4,12) = \frac{u^2 + v^2 - 2uvw}{1 - u^2}.$$
(2.17)

By setting $u = \cos \angle (12)$, $v = \cos \angle (23)$, $w = \cos \angle (13)$ and by substituting (2.16), (2.17), (2.11), (2.11) in (2.15) and taking into account that $\angle (3, 12) = -\angle (4, 12)$, we get:

$$u = vw + \frac{(uw - v)(uv - w)}{1 - u^2} - \frac{1 - u^2 - v^2 - w^2 + 2uvw}{1 - u^2}.$$
 (2.18)

Hence, by solving w.r to u, we obtain that:

$$u = -1 - v - w$$

or

$$1 + \cos \angle (12) + \cos \angle (13) + \cos \angle (24) = 0,$$

which gives

$$\vec{u}(0,1) + \vec{u}(0,2) + \vec{u}(0,3) + \vec{u}(0,4) = 0.$$

Therefore $O = A_0$, because the Fermat-Torricelli point A_0 is unique.

Corollary 1. If $A_1A_2A_3A_4$ is an almost platonic tetrahedron, then $A_0 = O$.

Proof. It is a direct consequence of Proposition 1 for a = b or a = c or b = c. \Box

Corollary 2. If $A_1A_2A_3A_4$ is a platonic tetrahedron, then $A_0 = O$.

Proof. It is a direct consequence of Proposition 1 for a = b = c.

Definition 5. The natural Fermat-Torricelli point in \mathbb{R}^3 (Fig 1) is the Fermat-Torricelli point of an almost Platonic tetrahedron with four equal edges $a\sqrt{3}$ and two opposite equal edges $a\sqrt{2}$, where a = R (radius of circumscribed sphere) is a positive real number.

Proposition 2. If A_0 is the natural Fermat-Torricelli point in \mathbb{R}^3 of an almost Platonic tetrahedron $A_1A_2A_3A_4$, such that $A_1A_2 = A_3A_4 = a\sqrt{2}$, then $\angle A_1A_0A_2 = \angle A_3A_0A_4 = 90^\circ$ and $\angle A_1A_0A_3 = \angle A_2A_0A_4 = \angle A_1A_0A_4 = \angle A_2A_0A_3 = 120^\circ$.

Proof. By applying the cosine law in $\triangle A_1 A_0 A_2$, $\triangle A_1 A_0 A_2$, we get:

$$\cos \angle A_1 A_0 A_2 = \cos \angle A_2 A_0 A_4 = 0$$

which implies that:

$$\angle A_1 A_0 A_2 = \angle A_3 A_0 A_4 = 90^{\circ}. \tag{2.19}$$

By substituting (2.19) in (1.5), we derive that:

$$\cos \angle A_1 A_0 A_3 = \cos \angle A_2 A_0 A_3 = -\frac{1}{2},$$



FIGURE 1. The natural Fermat-Torricelli point in \mathbb{R}^3 : $\angle A_1 A_0 A_2 = \angle A_3 A_0 A_4 = 90^\circ$ and $\angle A_1 A_0 A_3 = \angle A_2 A_0 A_4 = \angle A_1 A_0 A_4 = \angle A_2 A_0 A_3 = 120^\circ$.

which yields

$$\angle A_1 A_0 A_3 = \angle A_2 A_0 A_4 = \angle A_1 A_0 A_4 = \angle A_2 A_0 A_3 = 120^{\circ}.$$

Remark 1. The natural Fermat-Torricelli point coincides with the Fermat-Torricelli point in \mathbb{R}^2 for three non-collinear points $\{A_1A_2A_3\}$, which form a triangle with angles $\angle A_iA_0A_j < 120^\circ - \epsilon$, for ϵ a positive real number and i, j = 1, 2, 3 (fig 2). We note that the two right angles that appeared in the three dimensional case vanish, but the other angles $\angle A_1A_0A_2 = \angle A_2A_0A_3 = \angle A_1A_0A_3 = 120^\circ$ angles remain (Theorem of Torricelli) and the equiangular property holds in the two dimensional case.

We may expand or contract isosceles tetrahedra, almost platonic tetrahedra and regular tetrahedra to non regular tetrahedra with respect to their Fermat-Torricelli point A_0 whose Fermat-Torricelli point A'_0 remains the same $A_0 = A'_0$. Let $A_1A_2A_3A_4$ be an isosceles tetrahedron or an almost platonic tetrahedron or a regular tetrahedron and A'_i be a point that lies on the ray A_iA_0 for i = 1, 2, 3, 4, such that $A'_i \in [A_0, A_i]$ or $A_i \in [A_0, A'_i]$.

Proposition 3. If $\|\sum_{j=1, j \neq i}^{4} \frac{A'_{j}A'_{i}}{|A'_{j}A'_{i}|}\| > 1$, for each i = 1, 2, 3, 4, then $A'_{0} = A_{0}$.

Proof. By applying Theorem 2, we obtain the balancing condition of unit vectors with respect to the Fermat-Torricelli point A'_0



FIGURE 2. The natural Fermat-Torricelli point coincides with the Fermat-Torricelli point in $\mathbb{R}^2 \angle A_1 A_0 A_2 = \angle A_2 A_0 A_3 = \angle A_1 A_0 A_3 = 120^\circ$.

$$\sum_{i}^{4} \frac{\vec{A_{0}'A_{i}'}}{|A_{0}'A_{i}'|} = 0$$

or

$$\sum_{i}^{4} \frac{\vec{A_0A_i}}{|A_0A_i|} = 0$$

which gives $A'_0 = A_0$.

3. A CLASS OF EXPLICIT SOLUTIONS USING QUADRATIC EQUATIONS

We introduce a class of explicit solutions for the Fermat problem for tetrahedra derived by almost platonic tetrahedra

Let $A_1A_2A_3A_4$ be a tetrahedron and A_0 be the Fermat-Torricelli point inside $A_1A_2A_3A_4$.

We denote by a_{ij} the distance $|A_iA_j|$, for i, j = 0, 1, 2, 3, 4.

We set $x = a_{01}, y = a_{02}, z = a_{03}, d = a_{04}, \omega = \angle A_1 A_0 A_2, \phi = \angle A_2 A_0 A_3.$

Let $A'_1A'_2A'_3A'_4$ be an almost platonic tetrahedron, such that each vertex A'_i lies on the ray A_iA_0 or $A'_i \in [A_0, A_i]$ for i = 1, 2, 3, 4. Without loss of generality, we may assume that $\omega = \angle A_1A_0A_2 = \angle A_3A_0A_4$ and $\phi = \angle A_1A_0A_3 = \angle A_2A_0A_3 = \angle A_1A_0A_4 = \angle A_2A_0A_4$.

Given a_{12} and ω , we choose the values of a_{13} and a_{34} , such that the following inequalities are satisfied:

$$\frac{\sqrt{2}}{2} \frac{a_{12}}{1 - \cos \omega} \sin \arccos(\frac{-1 - \cos \omega}{2}) < a_{13} < \frac{\sqrt{2}}{2} \frac{a_{12}}{1 - \cos \omega}$$

and

$$z\sin\omega < a_{34} < z\sin\omega$$

where z is a positive root of the quadratic equation

$$z^{2} - 2\frac{\sqrt{2}}{2}\frac{a_{12}}{1 - \cos\omega}\cos\phi z + (\frac{\sqrt{2}}{2}\frac{a_{12}}{1 - \cos\omega})^{2} - a_{13}^{2} = 0.$$

Theorem 4. If $a_{23} = a_{13}$, then

$$x(a_{12},\omega) = y(a_{12},\omega) = \frac{\sqrt{2}a_{12}}{2(1-\cos\omega)},$$
(3.1)

$$z(\phi, x(a_{12}, \omega), a_{13}) = x(a_{12}, \omega) \cos \phi + \sqrt{a_{13}^2 - x^2(a_{12}, \omega) \sin^2 \phi}$$
(3.2)

 $d(z(\phi, x(a_{12}, \omega), a_{13}), \omega, a_{34}) = z(\phi, x(a_{12}, \omega), a_{13}) \cos \omega + \sqrt{a_{34}^2 - z^2(\phi, x(a_{12}, \omega), a_{13}) \sin^2 \omega}$ (3.3)

$$a_{14} = a_{24} = \sqrt{x^2(a_{12},\omega) + d^2(z(\phi, x(a_{12},\omega), a_{13}), \omega, a_{34}) - 2x(a_{12},\omega)d(z(\phi, x(a_{12},\omega), a_{13}), \omega, a_{34})\cos\phi}.(3.4)$$

Proof. By substituting $\omega = \angle A_1 A_0 A_2 = \angle A_3 A_0 A_4$ and $\phi = \angle A_1 A_0 A_3 = \angle A_2 A_0 A_3 = \angle A_1 A_0 A_4 = \angle A_2 A_0 A_4$, in (1.5), we obtain:

$$\cos\phi = \frac{-1 - \cos\omega}{2} < 0. \tag{3.5}$$

By applying the cosine law in $\triangle A_2 A_0 A_3$, $\triangle A_1 A_0 A_4$, $\triangle A_1 A_0 A_3$, $\triangle A_2 A_0 A_4$, $\triangle A_1 A_0 A_2$, $\triangle A_3 A_0 A_4$, we get:

$$\cos\phi = \frac{y^2 + z^2 - a_{23}^2}{2yz},\tag{3.6}$$

$$\cos\phi = \frac{x^2 + d^2 - a_{14}^2}{2xd} \tag{3.7}$$

$$\cos\phi = \frac{x^2 + z^2 - a_{13}^2}{2xz} \tag{3.8}$$

$$\cos\phi = \frac{y^2 + d^2 - a_{24}^2}{2yd} \tag{3.9}$$

$$\cos\omega = \frac{x^2 + y^2 - a_{12}^2}{2xy} \tag{3.10}$$

$$\cos\omega = \frac{z^2 + d^2 - a_{34}^2}{2zd} \tag{3.11}$$

By subtracting (3.6) from (3.8) and taking into account that $a_{31} = a_{23}$, we derive that x = y or $0 < yx = z^2 - a_{23}^2$, which gives x = y or $z > a_{23}$. The inequality $z > a_{23}$ is impossible, because (3.6) yields $z^2 < y^2 + z^2 < a_{23}^2$ (ϕ is an obtuse angle),

which implies that $z < a_{23}$. Therefore, we obtain x = y. By substituting x = y in (3.10), we obtain (3.1).

By substituting (3.1) in (3.8), we get:

$$z^{2} - 2x(a_{12},\omega)\cos\phi z + x^{2}(a_{12},\omega) - a_{13}^{2} = 0.$$
(3.12)

The lower and upper bound of a_{13}

$$\frac{\sqrt{2}}{2} \frac{a_{12}}{1 - \cos \omega} \sin \arccos(\frac{-1 - \cos \omega}{2}) < a_{13} < \frac{\sqrt{2}}{2} \frac{a_{12}}{1 - \cos \omega}$$

yields the positive root (3.2)

By substituting (3.2) in (3.11), we get:

$$d^{2} - 2z(\phi, x(a_{12}, \omega), a_{13})\cos\omega d + z^{2}(\phi, x(a_{12}, \omega), a_{13}) - a_{34}^{2} = 0$$
(3.13)

The lower and upper bound of a_{34}

$$z \sin \omega < a_{34} < z$$

yields the positive root (3.3) of the quadratic equation (3.13). By substituting (3.1), (3.3) in (3.7), (3.9), we derive (3.4).

Example 1. Let $\omega = 105^{\circ}$, $a_{12} = 3$, $a_{23} = a_{13} = 4$, $a_{34} = 4.5$ By substituting these values in (3.5), (3.1), (3.2), (3.3), (3.4), we get:

 $\phi = 111.752^{\circ}$, $a_{14} = a_{24} = 3.96628$, x = y = 1.89071, z = 2.89323 and d = 2.85566. Therefore, we obtain a tetrahedron, which is derived by an almost platonic tetrahedron with two equal angles 105° and 111.752° having each vertex at a prescribed ray, which is formed by the Fermat-Torricelli point A_0 and a vertex of the almost platonic tetrahedron.

Example 2. Derivation of Mehlhos explicit solution in [23]. Let $\omega = 106.654^{\circ}$, $a_{12} = 2, a_{23} = a_{13} = \sqrt{2}, a_{34} = 1$. By substituting these values in (3.5), (3.1), (3.2), (3.3), (3.4), we get:

 $\phi = 110.898^{\circ}, a_{14} = a_{24} = \sqrt{3}, x = y = 1.24679, z = 0.35732$ and d = 0.837174.

4. Concluding Remarks

Various properties of closed polyhedra whose vertices exist on prescribed rays that meet at a point have been given by A.D. Alexandrov in [3, Chapter 9]. By enriching this intersection point with the minimum property of Fermat's problem for tetrahedra, we are able to prescribe tetrahedra derived by almost platonic tetrahedra having each vertex at a prescribed ray, such that the corresponding Fermat-Torricelli point remains the same. Theorem 4 may offer a class of explicit solutions to the Fermat problem for tetrahedra, which partially answers a question posed by Ivanov-Tuzhilin's preprint for open problems in [17] that deals with the derivation of theoretical constructions for Fermat points for four points in \mathbb{R}^3 .

References

- Z. Abu-Abbas, M. Hajja, A note on the Fermat point of a Tetrahedron. Math. Gaz. 79,484, (1995), 117-118.
- [2] S. Abu-Saymeh, M. Hajja, On the Fermat-Torricelli points of tetrahedra and of higher dimensional simplexes. Math. Mag. 70, 5 (1997), 372-378.
- [3] A.D. Alexandrov, Convex Polyhedra, Springer Monographs in Mathematics. Berlin: Springer (2005).
- [4] V. Arnold, Arnold's Problems. Berlin Springer (2004).
- [5] C. Bajaj, The algebraic degree of geometric optimization problems. Discrete Comput. Geom. 3, 1-2 (1988), 177-191.
- [6] V. Boltyanski, H. Martini, V. Soltan, Geometric Methods and Optimization Problems, Kluwer, Dordrecht/Boston/London (1999).
- [7] R. Chandrasekaran, A. Tamir, Algebraic optimization: The Fermat-Weber location problem. Math. Program., Ser. A 46 (2) (1990), 219-224.
- [8] R. Courant, H. Robbins, What is mathematics? An elementary approach to ideas and methods. 2nd. ed. New York, NY: Oxford University Press (1996).
- [9] H.S.M. Coxeter, Introduction to geometry. 2nd ed. New York etc.: John Wiley and Sons, Inc. XVI, (1969).
- [10] C. Groß, T.-K. Strempel, On generalizations of conics and on a generalization of the Fermat-Torricelli problem. Am. Math. Mon. 105 (1998), No. 8, 732-743.
- [11] E. Egerváry, Über ein Minimumproblem der Elementargeometrie. J. reine angew. Math. 178 (1938), 174-186.
- [12] F. Eriksson, The Fermat-Torricelli problem a classical beauty in modern dress. (Swedish). Normat 39 (1991), No. 2, 64-75.
- [13] F. Eriksson, The Fermat-Torricelli problem once more. Math. Gaz. 81, 490 (1997), 37-44.
- [14] S. Gueron and R. Tessler, The Fermat-Steiner problem, Amer. Math. Monthly 109 (2002) 443-451.
- [15] P. Glaister, Tetrahedra Fermat points and centroids. Mathematical Gazette, 77 (1993), 360-361.
- [16] D. Hilbert, S. Cohn-Vossen, Geometry and the Imagination. New York: Chelsea, 1999.
- [17] A.O. Ivanov, A. Tuzhilin, What Spaces Permit Fermat Points Construction and Melzak Algorithm?, preprint https://dcglab.uniyar.ac.ru/sites/default/files/papers/problems/ -WhatSpacesPermitFermatPointsConstructionandMelzakAlgorithm.pdf, 2012.
- [18] H. Kuhn, A note on Fermat's problem Math. Programming 4 (1973), 98-107.
- [19] Y. Kupitz, H. Martini, The Fermat-Torricelli point and isosceles tetrahedra. J. Geom. 49, 1-2 (1994), 150-162.
- [20] Y.S. Kupitz, H. Martini, Geometric aspects of the generalized Fermat-Torricelli problem, Intuitive Geometry, Bolyai Society Math Studies 6 (1997), 55–127.
- [21] Y. Kupitz, H. Martini, Equifacial tetrahedra and a famous location problem. Math. Gaz. 83, 498 (1999), 464-467.
- [22] L.L.Lindelof, Sur les maxima et minima d'une fonction des rayons vecteurs mens d'un point mobile a plusieurs centres fixes, Acta Soc. Sc. Fenn. 8 (1867), 191-203.
- [23] St. Mehlhos, Simple counter examples for the unsolvability of the Fermat- and Steiner-Weberproblem by compass and ruler. Beitr. Algebra Geom. 41, 1 (2000), 151-158.
- [24] E.J. Cockayne, Z.A. Melzak, Euclidean constructibility in graph-minimization problems. Math. Mag. 42 (1969), 206-208.
- [25] R. Noda, T. Sakai, M. Morimoto, Generalized Fermat's problem. Can. Math. Bull. 34, 1 (1991), 96-104.
- [26] J. Mcgregor Smith, Y. Jang, M.k. Kim, Steiner Minimal Trees, Twist Angles, and the Protein Folding problem, Proteins: Structure, Function and Bioniformatics 66 (2007), 889-902.
- [27] R. Spira, The isogonic and Erdős-Mordell points of a simplex. Am. Math. Mon. 78 (1971), 856-864.
- [28] R. Sturm, Ueber den Punkt kleinster Entfernungsumme von gegebenen Punkten. Kronecker J. XCVII (1884).
- [29] R. Sturm, Existenzbeweis des Punktes kleinster Entfernungssumme von vier gegebenen Punkten., J. für Math (1913). 143, 241-249.

ANASTASIOS N. ZACHOS

- [30] J.L. Synge, The Torricelli-Fermat point Generalised, Preprint, DIAS-STP-87-38, https://dair.dias.ie/id/eprint/824.
- [31] V.M. Tikhomirov, Stories about maxima and minima. Mathematical World 1. Providence, RI: American Mathematical Society; Washington, DC: Mathematical Association of America, (1990).
- [32] W.P. Thurston, *Three-dimensional geometry and topology. Vol. 1.* Princeton Mathematical Series. 35. Princeton, NJ: Princeton University Press (1997).
- [33] A. Yu. Uteshev, Analytical solution for the generalized Fermat-Torricelli problem. Am. Math. Mon. 121, No. 4 (2014), 318-331.
- [34] A. Weber, Theory of the location of industries, Univ. of Chicago Press, 1929.
- [35] E. Weiszfeld, Sur le point lequel la somme des distances de n points donnes est minimum, Tohoku Math. J.43, (1937), 355–386.
- [36] E. Weiszfeld, F. Plastria, On the point for which the sum of the distances to n given points is minimum, An. Oper, Res. 167 (2009),7–41.
- [37] G. O. Wesolowsky, The Weber problem history and perspectives, J. Location Sci. 1 (1993), 5-23.

UNIVERSITY OF PATRAS, DEPARTMENT OF MATHEMATICS, GR-26500 RION, GREECE *Email address:* azachos@gmail.com