

DECOMPOSITION NUMBERS IN THE PRINCIPAL BLOCK AND SYLOW NORMALISERS

GUNTER MALLE AND NOELIA RIZO

To the memory of Gary Seitz

ABSTRACT. If G is a finite group and p is a prime number, we investigate the relationship between the p -modular decomposition numbers of characters of height zero in the principal p -block of G and the p -local structure of G . In particular we prove that, under certain conditions on the non-abelian composition factors of G , $d_{\chi 1_G} \neq 0$ for all irreducible characters χ of degree prime to p in the principal p -block of G if, and only if, the normaliser of a Sylow p -subgroup of G has a normal p -complement.

1. INTRODUCTION

Let G be a finite group, p a prime number, $\text{Irr}(G)$ the set of irreducible ordinary characters of G and $\text{IBr}(G)$ the set of irreducible p -Brauer characters of G . Then the restriction χ° of any $\chi \in \text{Irr}(G)$ to the set of p -regular elements of G can be written as

$$\chi^\circ = \sum_{\varphi \in \text{IBr}(G)} d_{\chi\varphi} \varphi,$$

where the $d_{\chi\varphi}$ are uniquely determined non-negative integers. These are called the p -modular *decomposition numbers* of G , and a great deal of literature is devoted to understand them.

In [NT, NT2] Navarro and Tiep initiated the investigation on relations between p -decomposition numbers and properties of Sylow p -normalisers, considering two different settings. In [NT] they conjectured that if $p > 3$ then $d_{\chi 1_G} \neq 0$ for all $\chi \in \text{Irr}_{p'}(G)$, that is, for all irreducible characters of G of degree prime to p , if and only if G has self-normalising Sylow p -subgroups, and that this happens if and only if $d_{\chi 1_G} = 1$ for all $\chi \in \text{Irr}_{p'}(G)$. Note that irreducible characters of degree prime to p lie in p -blocks of maximal defect, so this situation can only happen when the principal block is the only Brauer p -block of maximal defect of G . It is then natural to wonder what can be said when this is not the case and one restricts attention to the principal p -block $B_0(G)$. In this sense, in [NT2] they conjectured that $d_{\chi 1_G} \neq 0$ for all $\chi \in \text{Irr}(B_0(G))$ if and only if G has a normal p -complement. In this paper we consider the intersection of the two conditions in [NT]

Date: May 15, 2024.

2010 Mathematics Subject Classification. Primary 20C15, 20C20, 20D20, 20C33.

Key words and phrases. decomposition numbers, principal blocks, Sylow normalisers.

The first author gratefully acknowledges financial support by the DFG — Project-ID 286237555. The second author is supported by Ministerio de Ciencia e Innovación (Grant PID2019-103854GB-I00 and Grant PID2022-137612NBI00 funded by MCIN/AEI/10.13039/501100011033 and “ERDF A way of making Europe”) and a CDEIGENT grant CIDEIG/2022/29 funded by Generalitat Valenciana.

and [NT2], namely what happens if $d_{\chi_{1G}} = 1$ just for all $\chi \in \text{Irr}_{p'}(B_0(G))$. Our main result is:

Theorem A. *Let G be a finite group and $p > 3$ be a prime. Assume that all non-abelian simple composition factors of G of order divisible by p satisfy Property $(*)$ below. The following are equivalent:*

- (i) *For every $\chi \in \text{Irr}_{p'}(B_0(G))$ we have $d_{\chi_{1G}} \neq 0$.*
- (ii) *For every $\chi \in \text{Irr}_{p'}(B_0(G))$ we have $d_{\chi_{1G}} = 1$.*
- (iii) *For $P \in \text{Syl}_p(G)$ we have $\mathbf{N}_G(P) = P \times K$ for some $K \leq G$.*

Moreover, if G is p -solvable, this equivalence holds for every prime p .

Notice that, as pointed out in [NT], all irreducible characters of odd degree of the alternating group $\mathfrak{A}_5$ contain the trivial character in their 2-modular reduction, and similarly, all irreducible characters of the Ree group ${}^2G_2(27)$ of non-zero 3-defect contain the trivial character in their 3-modular reduction, while the respective Sylow p -normalisers have no normal p -complement, so the equivalence in Theorem A fails for non-solvable groups with $p \leq 3$.

Theorem A involves the following property that a finite non-abelian simple group might, or might not, satisfy:

Property $(*)$. *Let S be non-abelian simple and $p > 3$ a prime dividing $|S|$. Then for all almost simple groups H with socle S and $|H : S|$ a p -power, there exists $\chi \in \text{Irr}_{p'}(B_0(H))$ such that $d_{\chi_{1H}} = 0$.*

We prove Property $(*)$ does hold for many simple groups: for sporadic groups, alternating groups, and simple groups of Lie type in characteristic different from p . We also show it for some groups of Lie type in characteristic p . (The general case of groups of Lie type in their defining characteristic was also left open in [NT] and [NT2].) As the knowledge on p -decomposition numbers for groups of Lie type in their own characteristic is too weak at present we refrain from making a general conjecture and just leave it as a question as to whether Property $(*)$ holds for all non-abelian finite simple groups.

Structure of the paper. In Section 2 we prove Property $(*)$ for S a sporadic simple group, an alternating group, a simple group of Lie type in characteristic different from p as well as for some groups of Lie type in characteristic p . In Section 3 we show that Theorem A holds for p -solvable groups and in Theorem 3.4 we reduce the general case to the validity of Property $(*)$ on composition factors, thus completing the proof of Theorem A.

Acknowledgements. The second author would like to thank Gabriel Navarro for bringing this topic to her attention.

2. ALMOST SIMPLE GROUPS

In this section, we discuss instances of Property $(*)$ from the introduction.

2.1. Alternating and sporadic groups. We start out with simple groups not of Lie type.

Proposition 2.1. *Property $(*)$ holds for S a sporadic simple group or the Tits group.*

Proof. Let S be as in the assumption and $H \geq S$ as in Property (*). Since $p > 2$ this means $H = S$. If S has abelian Sylow p -subgroups, by [NT2, Prop. 3.2] there exists $\chi \in \text{Irr}(B_0(H))$ such that $d_{\chi 1_H} = 0$. By one direction of Brauer's height zero conjecture [KM13], we have $\chi \in \text{Irr}_{p'}(B_0(H))$ as required. Else, by inspection in [GAP], S has just one p -block of maximal defect and then by [NT, Lemma 3.2], there is $\chi \in \text{Irr}_{p'}(H) = \text{Irr}_{p'}(B_0(H))$ with $d_{\chi 1_H} = 0$. $\square$

Proposition 2.2. *Property (*) holds for S an alternating group.*

Proof. Let $S = \mathfrak{A}_n$ with $n \geq 5$. As $p > 2$ again we have $H = S$. Set $G := \mathfrak{S}_n$. Recall that the irreducible characters of $\mathfrak{S}_n$ are naturally labelled by partitions of n . If p divides n , then let $\chi = \chi^{(n-2, 1^2)} \in \text{Irr}(B_0(G))$, as in [NT2, Prop. 3.1]. Then χ° is the sum of two irreducible Brauer characters of degrees $n-2$ and $(n-2)(n-3)/2$ and hence it is of degree prime to p and does not contain any irreducible Brauer character of degree 1. Now, any $\theta \in \text{Irr}(S)$ under χ lies in $\text{Irr}_{p'}(B_0(S))$ and $d_{\theta 1_S} = 0$, as wanted.

So we may assume that p does not divide n . Let $n = a_k p^k + \dots + a_1 p + a_0$ be the p -adic expansion of n . Since p does not divide n , we have $a_0 > 0$. Suppose first that $a_0 > 1$. Consider $\chi = \chi^{(a_0, 1^{n-a_0})}$. By Peel's theorem (see [Ja78, Thm 24.1]), $\chi^\circ \in \text{IBr}(G)$. Hence, it is enough to show that $\chi(1) \neq 1$, $\chi(1)$ is p' , and $\chi \in \text{Irr}(B_0(G))$. By the hook length formula, χ has degree

$$\chi(1) = \frac{n!}{n \cdot (a_0 - 1)!(n - a_0)!} = \binom{n-1}{n-a_0}.$$

Then $\chi(1) \neq 1$ since $1 < a_0 < n$. Moreover

$$n-1 = a_k p^k + \dots + a_1 p + (a_0 - 1)$$

and

$$n - a_0 = a_k p^k + \dots + a_1 p,$$

so by Lucas' theorem we have

$$\chi(1) = \binom{n-1}{n-a_0} \equiv \prod_{i=1}^k \binom{a_i}{a_i} \binom{a_0-1}{0} \equiv 1 \pmod{p}.$$

Thus, $\chi(1)$ is not divisible by p . Finally since the p -core of $\lambda = (a_0, 1^{n-a_0})$ is (a_0) , we have that χ lies in the principal p -block by the Nakayama conjecture, as desired. Now take $\theta \in \text{Irr}(S)$ under χ , so $\theta \in \text{Irr}_{p'}(B_0(S))$ and $d_{\theta 1_S} = 0$.

Finally, suppose that $a_0 = 1$, so p divides $n-1$. In this case consider $\chi = \chi^{(n-3, 2, 1)}$, so χ lies in $B_0(G)$. This is the character in the proof of [NT, Lemma 3.1(ii)], so we are done in this case as well. $\square$

2.2. Groups of Lie type in non-defining characteristic. For groups of Lie type in cross characteristic we consider the following set-up. Let $\mathbf{G}$ be a simple linear algebraic group of adjoint type over an algebraically closed field of characteristic r and $F : \mathbf{G} \rightarrow \mathbf{G}$ a Steinberg map, with group of fixed points $G := \mathbf{G}^F$. It is well-known that any simple group of Lie type can be obtained as $S = [G, G]$ for $\mathbf{G}, F$ chosen suitably. Moreover, if $\mathbf{G}_{\text{sc}}$ denotes a simply connected covering of $\mathbf{G}$, with corresponding Steinberg map also denoted F , then $S \cong \mathbf{G}_{\text{sc}}^F / \mathbf{Z}(\mathbf{G}_{\text{sc}}^F)$, if S is not the Tits simple group, which was already discussed in Proposition 2.1. Let $(\mathbf{G}^*, F)$ be dual to $(\mathbf{G}, F)$ and $G^* := \mathbf{G}^{*F}$.

We let $\mathbf{B} \leq \mathbf{G}$ denote an F -stable Borel subgroup of $\mathbf{G}$, and set $B := \mathbf{B}^F$.

We recall that automorphisms of prime order $p \geq 5$ of simple groups of Lie type are either field automorphisms, or diagonal automorphisms for groups of types $\mathrm{PSL}_n(\epsilon q)$, with $\epsilon \in \{\pm 1\}$.

Proposition 2.3. *Property (*) holds for S as above if $|B|$ is prime to p .*

Proof. By assumption p divides $|S|$, hence also $|G| = |G^*|$. Let $1 \neq s \in G^*$ be a (semisimple) p -element in the centre of a Sylow p -subgroup of G^* , and let $\chi \in \mathrm{Irr}(G)$ be the semisimple character in the Lusztig series $\mathcal{E}(G, s)$, unique since $\mathbf{G}$ has connected centre, see [GM20, Def. 2.6.9]. By the degree formula for Jordan decomposition [GM20, Cor. 2.6.6], $\chi(1)$ is then prime to p , and also $\chi(1) > 1$ as p does not divide $|\mathbf{Z}(G^*)|$ and so $C_{G^*}(s) < G^*$. Furthermore, by [Hi88, Cor. 3.4], the semisimple character χ lies in the same p -block of G as the semisimple character in $\mathcal{E}(G, 1)$, i.e., the trivial character, so in the principal p -block. Since p does not divide $|B|$, the permutation module 1_B^G is projective, and thus contains the projective cover of the trivial module. But all constituents of 1_B^G are unipotent, so lie in $\mathcal{E}(G, 1)$ (see [GM20, Exmp. 3.2.6]). Hence, by Brauer reciprocity, 1_G° does not occur as a constituent of χ° . Taking for θ any character of S below χ we see that $\theta \in \mathrm{Irr}_{p'}(B_0(S))$ and $d_{\theta|_S} = 0$.

Next observe that the order of any outer diagonal automorphism of S divides the order of B . Thus, H is an extension of S by a p -power order field automorphism γ . Note that any such automorphism of S extends to G and then also induces a dual automorphism on G^* , which we denote γ^* (see e.g. [Tay, 5.6]). Now choose s more precisely to be also γ^* -invariant (which is possible as γ^* is a p -element, so necessarily has non-trivial fixed points on the centre of a γ^* -stable Sylow p -subgroup of G^*). Then χ is also γ -invariant by [Tay, Prop. 7.2]. Since the index $|G : S|$ is a divisor of $|B|$, hence prime to p , there then exists a γ -invariant $\theta \in \mathrm{Irr}_{p'}(B_0(S))$ below χ , with $d_{\theta|_S} = 0$. Let $\tilde{\theta}$ be an extension of θ to $H = S\langle\gamma\rangle$ in $B_0(H)$. Then $\tilde{\theta}$ is as desired. $\square$

Proposition 2.4. *Property (*) holds for S as above if F is a Frobenius map with respect to an $\mathbb{F}_q$ -structure and p divides $q - 1$.*

Proof. Let W be the Weyl group of G , that is, the F -fixed points of the Weyl group of $\mathbf{G}$. As p divides $q - 1$, the p -decomposition matrix of the group algebra of W embeds into the p -modular decomposition matrix of G (see [Di90, 4.10]). Let $\epsilon \in \mathrm{Irr}_{p'}(W)$ be the (linear) sign character. Then $\epsilon^\circ \neq 1_W^\circ$ since $p > 2$, whence $d_{\epsilon|_W} = 0$. Now ϵ corresponds to the Steinberg character St of G . Then $d_{\mathrm{St}|_G} = d_{\epsilon|_W} = 0$, St lies in the principal p -block of G (e.g. by [En00, Thm A]) and its degree is a power of the defining characteristic, so prime to p . Now note that any p -automorphism of S is realised inside the extension of G (which induces all diagonal automorphisms) by a generator γ of the cyclic group of p -power order field automorphisms, so we may assume $H \leq \tilde{G} := G\langle\gamma\rangle$. By [GM20, Thm 4.5.11], St is invariant under γ . Let $\tilde{\mathrm{St}}$ be an extension of St to $\tilde{G}$ in $B_0(\tilde{G})$, so $d_{\tilde{\mathrm{St}}|_{\tilde{G}}} = 0$. Then $\tilde{\mathrm{St}}|_H$ is irreducible, since St restricts irreducibly to S , hence lies in $B_0(H)$ and so is as required. $\square$

Theorem 2.5. *Property (*) holds for S of Lie type when p is not the defining characteristic.*

Proof. By Proposition 2.3 we may assume that p divides the order of a Borel subgroup B of G . If F is a Frobenius map with respect to an $\mathbb{F}_q$ -structure and p divides $q - 1$, we are done by Proposition 2.4. If G is a Suzuki or Ree group and p divides $|B|$, then $p|(q^2 - 1)$ where F^2 defines an $\mathbb{F}_{q^2}$ -structure, and the exactly same arguments as in the proof of Proposition 2.4 apply.

So we are reduced to the case that F is a Frobenius map with respect to an $\mathbb{F}_q$ -structure and p divides $|B|$ but not $q - 1$. Since p is not the defining prime, this implies that G is a twisted group of Lie type ${}^2A_{n-1}$, 2D_n , or 2E_6 and p divides $q + 1$, respectively of type 3D_4 and p divides $q^2 + q + 1$. Let $d = 2, 3$ in the respective cases. Then the centraliser of a Sylow d -torus of G is a maximal torus, so has a unique d -cuspidal unipotent character. Thus, by [En00, Thm A] there is a unique unipotent block of G of maximal defect, the principal block, which hence contains the Steinberg character St of G . By [Hi90, Thm B] we have $d_{\text{St}1_G} = 0$ in our case unless $G = \text{PGU}_3(q)$. Except for that latter case, we can now argue as in the proof of Proposition 2.4 to conclude. For $G = \text{PGU}_3(q)$ let χ be the cuspidal unipotent character of degree $q(q - 1)$, prime to p . By [Ge90, Thm 4.3(a)] it lies in $B_0(G)$ and satisfies $d_{\chi 1_G} = 0$. Again, χ restricts irreducibly to S and is invariant under all automorphisms, so we can argue as before. $\square$

2.3. Groups of Lie type in defining characteristic. We do not see how to approach Property (*) for groups of Lie type in their defining characteristic in general. All characters of positive defect lie in the principal block and decomposition numbers tend to be large and little is known. The following was shown by Navarro and Tiep:

Proposition 2.6. *Property (*) holds if $S = \text{PSp}_{2n}(p^f)$, $n \geq 1$, with either $p > 3$ or $p = 3$ and n even.*

Proof. In this case the principal block of S is the only p -block of positive defect. Let H be almost simple with H/S a p -group. By [Na98, Cor. 9.6] there is just one p -block of H covering $B_0(S)$, necessarily the principal block $B_0(H)$. Now the irreducible character $\chi \in \text{Irr}_{p'}(H)$ with $d_{\chi 1_H} = 0$ constructed in [NT, Prop. 3.11] lies above a character of S of positive defect, hence in the principal p -block of H and we are done. $\square$

Observe that this does not extend to $p = 3$ and n odd: The group $H = \text{PSp}_2(3^3).3$ has no irreducible character $\chi \in \text{Irr}_{3'}(H)$ with $d_{\chi 1_H} = 0$.

The paper [NT2] also contains results for special linear and unitary groups but these are not applicable here as the considered characters are not of p' -degree. Nevertheless, we can follow their general approach.

For $G = \text{SL}_n(q)$ we let τ_j , $j = 1, \dots, q - 2$ denote the non-unipotent *Weil characters* of degree $(q^n - 1)/(q - 1)$, ordered such that τ_j is trivial on the centre $Z(\text{SL}_n(q))$ of order $z := \gcd(n, q - 1)$ if and only if $z|j$.

Proposition 2.7. *Let $S = \text{PSL}_n(q)$ with $q = p^f$, $p \neq 2$ and $n \geq 3$.*

- (a) *If either $\gcd(p - 1, (q - 1)/\gcd(n, q - 1)) > 1$ or $2^f < (q - 1)/z - 1$ then there is $\chi \in \text{Irr}_{p'}(B_0(S))$ such that $d_{\chi 1_S} = 0$.*
- (b) *Write $f = p^a f'$ with $\gcd(p, f') = 1$ and set $q' := p^{f'}$. If $a = 0$ or $2^f < (q' - 1)/(q' - 1, n) - 1$ then Property (*) holds for $S = \text{PSL}_n(q)$.*

Proof. Let $G := \mathrm{SL}_n(q)$ and set $z = \gcd(n, q - 1)$. We are interested in the characters τ_j of G that are trivial on $Z(G)$, that is, for which $j = zj'$ for some integer $1 \leq j' \leq (q - 1 - z)/z$. By [ZS, Thm 1.11] the decomposition number $d_{\tau_j 1_G}$ equals the number of solutions $x_s \in \{0, 1\}$ of the congruence

$$n(p - 1) \sum_{s=0}^{f-1} x_s p^s \equiv j \pmod{(q - 1)}.$$

Dividing by z , we need to count solutions to

$$n/z(p - 1) \sum_{s=0}^{f-1} x_s p^s \equiv j' \pmod{(q - 1)/z}.$$

If there is a prime ℓ dividing $p - 1$ and $(q - 1)/z$, then reducing modulo ℓ we see there is no solution for $j' = 1$. Also, the left hand side can take at most 2^f distinct values. Since there are $(q - 1)/z - 1$ admissible values for j' , there is j' with no solutions whenever $2^f < (q - 1)/z - 1$. Thus, under either of our assumptions we find j' with $\tau_j = \tau_{zj'} \in \mathrm{Irr}_{p'}(G)$ with $d_{\tau_j 1_G} = 0$. Since G has a single p -block of positive defect, τ_j lies in the principal block. Furthermore, by construction $Z(\mathrm{SL}_n(q))$ lies in the kernel of τ_j and hence τ_j deflates to a character of $S = \mathrm{PSL}_n(q)$. Thus we get (a).

(b) Since $p > 2$ does not divide $q - 1$, the p -power order automorphisms of S are field automorphisms, of order dividing p^a where $f = p^a f'$ is as in the statement. If $a = 0$ Property (*) follows from (a) since necessarily $H = S$. For $a > 0$ let γ be a field automorphism of G (and hence of S) of order p^a . There are exactly $q' - 2$ Weil characters of G invariant under γ , which hence extend to $G\langle\gamma\rangle$. Of these, $(q' - 1)/\gcd(q' - 1, n) - 1$ are trivial on $Z(G)$, so define characters in $\mathrm{Irr}_{p'}(S)$ invariant under γ . By the argument above, if this number is bigger than 2^f then there exists such a character χ with $d_{\chi 1_S} = 0$. Hence any character of H in $B_0(H)$ lying above it verifies Property (*). $\square$

Note that the case $n = 2$ is contained is addressed in Proposition 2.6. Observe that the condition in Proposition 2.7(a) is satisfied if $\gcd(n, q - 1) = 1$, for example. It also holds when $p > n + 1$ (since $p - 1$ always divides $q - 1$), or if $q > n(2^f + 1)$. Thus, Proposition 2.7 extends and complements [NT2, Prop. 3.3(ii)].

Corollary 2.8. *Let $S = \mathrm{PSL}_n(q)$ with $q = p^f$, $p \neq 2$ and $3 \leq n \leq 9$. Then there is $\chi \in \mathrm{Irr}_{p'}(B_0(S))$ with $d_{\chi 1_S} = 0$ unless possibly S is one of*

$$\mathrm{PSL}_4(5), \mathrm{PSL}_6(3), \mathrm{PSL}_6(7), \mathrm{PSL}_8(3), \mathrm{PSL}_8(9), \mathrm{PSL}_8(5), \mathrm{PSL}_8(25).$$

Proof. By the conditions in Proposition 2.7(a) and directly checking with the formula of Zalesski–Suprunenko if these fail one finds that the only cases with $n \leq 9$ not covered are $\mathrm{PSL}_4(3)$ and those listed in the statement. The decomposition matrix of $\mathrm{PSL}_4(3)$ is in [GAP] from which the claim can be verified. $\square$

For $G = \mathrm{SU}_n(q)$ we let τ_j , $j = 1, \dots, q$ denote the non-unipotent Weil characters, constructed by Seitz [Se75], of degree $(q^n - (-1)^n)/(q + 1)$, again ordered such that τ_j is trivial on the centre $Z(\mathrm{SU}_n(q))$ of order $z := \gcd(n, q + 1)$ if and only if $z|j$.

Proposition 2.9. *Let $S = \mathrm{PSU}_n(q)$ with $q = p^f$, $p \neq 2$ and $n \geq 3$.*

(a) *If $2^f < (q + 1)/z - 1$ then there is $\chi \in \mathrm{Irr}_{p'}(B_0(S))$ such that $d_{\chi 1_S} = 0$.*

- (b) Write $f = p^a f'$ with $\gcd(p, f') = 1$ and set $q' := p^{f'}$. If $a = 0$ or $2^f < (q' + 1)/(q' + 1, n) - 1$ then Property $(*)$ holds for $S = \text{PSU}_n(q)$.

Proof. The argument is very similar to the one for the special linear groups. Let $G = \text{SU}_n(q)$ and set $z = \gcd(n, q + 1) = |Z(G)|$. Again, we consider characters τ_j trivial on $Z(G)$, that is, for which $j = zj'$ for some integer $1 \leq j' \leq (q + 1 - z)/z$. By [Za, Main Thm] the decomposition number $d_{\tau_j 1_G}$ equals the number of solutions $x_s \in \{0, 1\}$ of the congruence

$$n((p - 1) \sum_{s=0}^{f-1} x_s p^s - 1) \equiv j \pmod{(q + 1)}.$$

(In fact, [Za] has an additional summand of $\frac{q+1}{2}$ on the right hand side, but this disappears here due to a different numbering of the τ_j , see [NT2, p. 612]; in any case, this difference will not matter for our argument here.) Since the left-hand side can take at most 2^f distinct values, while there are $(q + 1)/z - 1$ admissible values for j' the assertion in (a) follows. For (b) we can argue exactly as in the proof of Proposition 2.7. $\square$

As in [NT, NT2] we have no general results for orthogonal or exceptional type groups in their defining characteristic.

3. THE REDUCTION

In this section we prove Theorem A. We will need the following results, which we collect here for the reader's convenience.

Lemma 3.1. *Let $N \triangleleft G$ and let $\theta \in \text{Irr}_{p'}(B_0(N))$. Suppose that θ extends to PN , where $P \in \text{Syl}_p(G)$. Then there exists $\chi \in \text{Irr}_{p'}(B_0(G))$ satisfying $[\theta^G, \chi] \neq 0$.*

Proof. This is [Mur, Lemma 4.3]. $\square$

The following argument is inside the proof of [NT, Thm 2.6].

Lemma 3.2. *Let G be a finite group and suppose that $d_{\chi 1_G} \neq 0$ for every $\chi \in \text{Irr}_{p'}(B_0(G))$. Let $M \triangleleft G$ and let $P \in \text{Syl}_p(G)$. Then $d_{\psi 1_{MP}} \neq 0$ for every $\psi \in \text{Irr}_{p'}(B_0(MP))$.*

Proof. Since MP/M is a p -group, we have that $\psi_M = \tau \in \text{Irr}_{p'}(B_0(M))$. By Lemma 3.1 there exists $\chi \in \text{Irr}_{p'}(B_0(G))$ lying over τ . By hypothesis, $d_{\chi 1_G} \neq 0$, and then χ_M° contains 1_M , so $d_{\tau 1_M} \neq 0$. Since MP/M is a p -group, we have that $(MP)^\circ = M^\circ$ and then $d_{\psi 1_{MP}} \neq 0$, as wanted. $\square$

We next prove Theorem A for p -solvable groups.

Theorem 3.3. *Let G be a p -solvable group. Then the following are equivalent:*

- (i) *For every $\chi \in \text{Irr}_{p'}(B_0(G))$ we have $d_{\chi 1_G} \neq 0$.*
- (ii) *For every $\chi \in \text{Irr}_{p'}(B_0(G))$ we have $d_{\chi 1_G} = 1$.*
- (iii) *For $P \in \text{Syl}_p(G)$ we have $\mathbf{N}_G(P) = P \times K$ for some $K \leq G$.*

Proof. We first prove (iii) implies (ii). By [NTV, Thm 3.2] we have $K \subseteq \mathbf{O}_{p'}(G) =: X$. Now, let $\overline{G} = G/X$ and $\overline{P} = PX/X$, then (iii) implies that $\mathbf{N}_{\overline{G}}(\overline{P}) \cong \overline{P}$. Since $\text{Irr}_{p'}(B_0(G)) = \text{Irr}_{p'}(B_0(G/X))$ we know by [NT, Thm B] that (i) and (ii) hold.

Since (ii) implies (i) trivially, we just need to show that (i) implies (iii). We proceed by induction on $|G|$. Let $N = \mathbf{O}_{p'}(G)$ and use the bar notation. Since $\text{Irr}_{p'}(B_0(G)) = \text{Irr}_{p'}(B_0(\bar{G}))$, if $N > 1$, we have by induction that $\mathbf{N}_{\bar{G}}(\bar{P}) = \bar{P} \times \bar{K}$. By [NTV, Thm 3.2] we have that $\bar{K} \subseteq \mathbf{O}_{p'}(\bar{G}) = 1$ so $\bar{K} = 1$ and hence $\mathbf{N}_{\bar{G}}(\bar{P}) = \bar{P}$. This implies that $\mathbf{N}_G(P) = P \times \mathbf{C}_N(P)$ and we are done. So we may assume that $N = 1$. But in this case the principal p -block is the only p -block of G , and we are done by [NT, Thm B]. $\square$

We finally prove Theorem A.

Theorem 3.4. *Let G be a finite group. Assume that $p > 3$ and all non-abelian composition factors of G of order divisible by p satisfy Property (*). Then the following are equivalent:*

- (i) *For every $\chi \in \text{Irr}_{p'}(B_0(G))$ we have $d_{\chi 1_G} \neq 0$.*
- (ii) *For every $\chi \in \text{Irr}_{p'}(B_0(G))$ we have $d_{\chi 1_G} = 1$.*
- (iii) *For $P \in \text{Syl}_p(G)$ we have $\mathbf{N}_G(P) = P \times K$ for some $K \leq G$.*

Proof. As in Theorem 3.3, we just need to show that (i) implies (iii). We work to prove this implication in a series of steps, proceeding by induction on $|G|$.

Step 1. We may assume $\mathbf{O}_{p'}(G) = 1$. Indeed, let $N = \mathbf{O}_{p'}(G)$ and use the bar notation. Since $\text{Irr}_{p'}(B_0(G)) = \text{Irr}_{p'}(B_0(\bar{G}))$, if $N > 1$, we have by induction that $\mathbf{N}_{\bar{G}}(\bar{P}) = \bar{P} \times \bar{K}$. By [NTV, Thm 3.2] we have that $\bar{K} \subseteq \mathbf{O}_{p'}(\bar{G}) = 1$ so $\bar{K} = 1$ and hence $\mathbf{N}_{\bar{G}}(\bar{P}) = \bar{P}$. This implies that $\mathbf{N}_G(P) = P \times \mathbf{C}_N(P)$ and we are done.

Step 2. If $1 < M \triangleleft G$ then M is not p -solvable. In particular, if N is a minimal normal subgroup of G , then N is semisimple of order divisible by p . Let $M \triangleleft G$ be the largest p -solvable normal subgroup of G . We claim that $M = 1$. Let $\bar{G} = G/M$ and use the bar notation. Suppose $M > 1$. Since $\text{Irr}_{p'}(B_0(\bar{G})) \subseteq \text{Irr}_{p'}(B_0(G))$ we have by induction that $\mathbf{N}_{\bar{G}}(\bar{P}) = \bar{P} \times \bar{K}$. By [NT, Thm 3.2] this implies that $\bar{K} \subseteq \mathbf{O}_{p'}(\bar{G})$. Since M is the largest normal p -solvable subgroup of G , $\mathbf{O}_{p'}(\bar{G})$ is trivial and hence $\mathbf{N}_{\bar{G}}(\bar{P}) = \bar{P}$. By [GMN, Thm 1.1] this forces $\bar{G}$ to be solvable. Hence G is p -solvable and we are done by Theorem 3.3.

From now on, let $N = S_1 \times \cdots \times S_t$ be a minimal normal subgroup of G , where $S_i \cong S$ is a nonabelian simple group of order divisible by p .

Step 3. Let $X/N = \mathbf{O}_{p'}(G/N)$. Then $G = XP$. Since $\text{Irr}_{p'}(B_0(G/N)) \subseteq \text{Irr}_{p'}(B_0(G))$, by induction we have $\mathbf{N}_G(P)N/N = \mathbf{N}_{G/N}(PN/N) = PN/N \times K/N$ for some $K \leq G$. Then $K/N \subseteq X/N$ by [NTV, Thm 3.2]. Write $H = XN\mathbf{N}_G(P) = XPNK = XP$. By Lemma 3.2 applied to X in place of M , we obtain that H satisfies (i). Since the non-abelian composition factors of H are composition factors of G , if $H < G$, by induction we have $\mathbf{N}_G(P) = \mathbf{N}_H(P) = P \times L$ for some L and we are done. Hence we may assume that $H = G$.

Step 4. Let $H = \bigcap \mathbf{N}_G(S_i)$ and let X be as in Step 3. Then $G = (H \cap X)P$. Write $Y = (H \cap X)P$. We claim first that $\mathbf{O}_{p'}(Y) = 1$. Indeed, $\mathbf{O}_{p'}(Y) \cap (H \cap X) \subseteq \mathbf{O}_{p'}(H \cap X) = 1$ by Step 1. Now, $\mathbf{O}_{p'}(Y) \cong (H \cap X)\mathbf{O}_{p'}(Y)/(H \cap X)$ is a p -group, so $\mathbf{O}_{p'}(Y) = 1$ as wanted. By Lemma 3.2 applied to $H \cap X$ in place of M , we have that Y satisfies (i). Since the non-abelian composition factors of Y are composition factors of G , if $Y < G$, by induction this gives $\mathbf{N}_Y(P) = P \times K$ for some K . But then $K \subseteq \mathbf{O}_{p'}(Y)$ by [NTV,

Thm 3.2], so $K = 1$. This means that $\mathbf{N}_Y(P) = P$ and then by [GMN, Thm 1.1] the group Y is solvable. But then N is solvable, a contradiction. Hence $Y = G$ as wanted.

Step 5. Final step. Let $H = \bigcap \mathbf{N}_G(S_i)$ and X be as before. Since $G = (H \cap X)P$ and H acts trivially on $\{S_1, \dots, S_t\}$, P must act transitively on the set $\{S_1, \dots, S_t\}$. Write $S = S_1$ and, for $i = 2, \dots, t$, write $S_i = S^{x_i}$ with $x_i \in P$. We proceed now as in the proof of [NT, Thm 2.6]. Let $R = \mathbf{N}_P(S)$. If $SR = G$, then $S = N$ and $\mathbf{C}_G(S)$ is a normal p -subgroup of G . By Step 2, $\mathbf{C}_G(S) = 1$ and hence G is almost simple with socle S and $|G : S|$ a power of p . Since S satisfies Property $(*)$ by assumption, we have a contradiction. Hence we may assume that $SR < G$.

Let $Q = P \cap N$ and let $R_1 = R \cap S = Q \cap S = P \cap S \in \text{Syl}_p(S)$. Let $\gamma \in \text{Irr}_{p'}(B_0(SR))$ and notice that $\gamma_S = \psi \in \text{Irr}_{p'}(B_0(S))$ since SR/S is a p -group. For $i = 2, \dots, t$, let $\psi_i = \psi^{x_i} \in \text{Irr}_{p'}(B_0(S_i))$ and let $\eta = \psi \times \psi_2 \times \dots \times \psi_t \in \text{Irr}_{p'}(B_0(N))$, which is P -invariant by [NTT, Lemma 4.1(ii)]. Then η extends to PN . By Lemma 3.1 and hypothesis we have $d_{\eta_{1_N}} \neq 0$. By [NT, Lemma 2.3] we have that $d_{\psi_{1_S}} \neq 0$ and then, since SR/S is a p -group, we conclude that $d_{\gamma_{1_{SR}}} \neq 0$. Since $SR < G$ and S is a composition factor of G , this implies $\mathbf{N}_{SR}(R) = R \times K$ for some K . Then $K \subseteq \mathbf{O}_{p'}(SR)$. But $\mathbf{O}_{p'}(SR) = 1$ (as before, $\mathbf{O}_{p'}(SR) \cap S = 1$, so $\mathbf{O}_{p'}(SR) \cong \mathbf{SO}_{p'}(SR)/S$ is a p -group, and hence trivial), so $K = 1$ and then $\mathbf{N}_{SR}(R) = R$. Now by [GMN, Thm 1.1], SR is solvable, and hence S is solvable, a final contradiction. $\square$

We remark that, as happens in [NT], (iii) $\Rightarrow$ (ii) $\Rightarrow$ (i) is always true if $p > 3$, as is shown in the first lines of the proof of Theorem 3.3 (notice that that part does not require p -solvability, since [NTV, Thm 3.2] holds for odd primes).

REFERENCES

- [Di90] R. DIPPER, On quotients of Hom-functors and representations of finite general linear groups. I. *J. Algebra* **130** (1990), 235–259.
- [En00] M. ENGUEHARD, Sur les l -blocs unipotents des groupes réductifs finis quand l est mauvais. *J. Algebra* **230** (2000), 334–377.
- [Ge90] M. GECK, Irreducible Brauer characters of the 3-dimensional special unitary groups in nondefining characteristic. *Comm. Algebra* **18** (1990), 563–584.
- [GM20] M. GECK, G. MALLE, *The Character Theory of Finite Groups of Lie Type: A Guided Tour*. Cambridge University Press, Cambridge, 2020.
- [GMN] R. GURALNICK, G. MALLE, G. NAVARRO, Self-normalizing Sylow subgroups. *Proc. Amer. Math. Soc.* **132** (2004), 973–979.
- [Hi88] G. HISS, Regular and semisimple blocks of finite reductive groups. *J. London Math. Soc. (2)* **41** (1990), 63–68.
- [Hi90] G. HISS, The number of trivial composition factors of the Steinberg module. *Arch. Math. (Basel)* **54** (1990), 247–251.
- [Ja78] G. D. JAMES, *The Representation Theory of the Symmetric Groups*. Lecture Notes in Mathematics, 682. Springer, Berlin, 1978.
- [KM13] R. KESSAR, G. MALLE, Quasi-isolated blocks and Brauer’s height zero conjecture. *Ann. of Math. (2)* **178** (2013), 321–384.
- [Mur] M. MURAI, Block induction, normal subgroups and characters of height zero. *Osaka J. Math.* **31** (1994), 9–25.
- [Na98] G. NAVARRO, *Characters and Blocks of Finite Groups*. Volume 250 of London Mathematical Society Lecture Note Series, Cambridge University Press, Cambridge, 1998.

- [NT] G. NAVARRO, P.H. TIEP, Decomposition numbers and local properties. *J. Algebra* **558** (2020), 620–639.
- [NT2] G. NAVARRO, P.H. TIEP, Decomposition numbers and global properties. *J. Algebra* **607** (2022), 607–617.
- [NTT] G. NAVARRO, P.H. TIEP, A. TURULL, p -Rational characters and self-normalizing Sylow p -subgroups. *Represent. Theory* **11** (2007), 84–94.
- [NTV] G. NAVARRO, P.H. TIEP, C. VALLEJO, McKay natural correspondences on characters. *Algebra Number Theory* **8** (2014), 1839–1856.
- [Se75] G. M. SEITZ, Some representations of classical groups. *J. London Math. Soc. (2)* **10** (1975), 115–120.
- [Tay] J. TAYLOR, Action of automorphisms on irreducible characters of symplectic groups. *J. Algebra* **505** (2018), 211–246.
- [GAP] THE GAP GROUP, *GAP — Groups, Algorithms, and Programming, Version 4.11.0*; 2020, <http://www.gap-system.org>.
- [Za] A. E. ZALESSKIĬ, A fragment of the decomposition matrix of the special unitary group over a finite field. *Math. USSR-Izv.* **36** (1991), 23–39.
- [ZS] A. E. ZALESSKIĬ, I. D. SUPRUNENKO, Permutation representations and a fragment of the decomposition matrix of symplectic and special linear groups over a finite field. *Siberian Math. J.* **31** (1991), 744–755.

(G. Malle) FB MATHEMATIK, RPTU, POSTFACH 3049, 67653 KAISERSLAUTERN, GERMANY.
Email address: `malle@mathematik.uni-kl.de`

(N. Rizo) DEPARTAMENT DE MATEMÀTIQUES, UNIVERSITAT DE VALÈNCIA, 46100 BURJASSOT, VALÈNCIA, SPAIN
Email address: `noelia.rizo@uv.es`