THE LATTICE OF SUBMONOIDS OF THE UNIFORM BLOCK PERMUTATIONS CONTAINING THE SYMMETRIC GROUP

ROSA ORELLANA, FRANCO SALIOLA, ANNE SCHILLING, AND MIKE ZABROCKI

ABSTRACT. We study the lattice of submonoids of the uniform block permutation monoid containing the symmetric group (which is its group of units). We prove that this lattice is distributive under union and intersection by relating the submonoids containing the symmetric group to downsets in a new partial order on integer partitions. Furthermore, we show that the sizes of the \mathscr{J} -classes of the uniform block permutation monoid are sums of squares of dimensions of irreducible modules of the monoid algebra.

1. INTRODUCTION

A block permutation is a bijection between the set of blocks of a set partition A of $[k] = \{1, 2, \ldots, k\}$ and the set of blocks of a set partition B of $[\overline{k}] = \{\overline{1}, \overline{2}, \ldots, \overline{k}\}$. A block permutation is uniform if the image of each block A_i of A has the same cardinality as A_i . There are at least three approaches to the algebra of uniform block permutations that are established in the literature. The first is as a factorizable inverse semigroup [2, 4, 3]. The second is as a centralizer algebra [7, 8, 9, 10], where it is a limiting case of the Tanabe algebra [15]. The third is as a Hopf algebra [1] where the set of all uniform block permutations are graded by size and there is an external product and coproduct.

In [11], we studied the representation theory of the algebra of uniform block permutations as a monoid algebra in the pursuit of computing the restriction of an irreducible GL_n module to the symmetric group \mathfrak{S}_n of permutation matrices, where note that $\mathfrak{S}_n \subseteq GL_n$.

For a positive integer k, let \mathcal{U}_k denote the monoid of uniform block permutations of the set $[k] \cup [\overline{k}]$ with the usual diagram algebra product. The irreducible representations of \mathcal{U}_k are indexed by sequences of partitions $\vec{\lambda} = (\lambda^{(1)}, \lambda^{(2)}, \dots, \lambda^{(k)})$ such that $|\lambda^{(1)}| + 2|\lambda^{(2)}| + \dots + k|\lambda^{(k)}| = k$. Let $W_{\mathcal{U}_k}^{\vec{\lambda}}$ denote the irreducible \mathcal{U}_k -module indexed by $\vec{\lambda}$. One of the main results of [11] is a formula in terms of the operation of plethysm on symmetric functions for the multiplicity of an irreducible symmetric group module in $\operatorname{Res}_{\mathfrak{S}_k}^{\mathcal{U}_k} W_{\mathcal{U}_k}^{\vec{\lambda}}$. Since both \mathcal{U}_k and \mathfrak{S}_k are monoids, it is natural to ask for a description of the monoids

Since both \mathcal{U}_k and \mathfrak{S}_k are monoids, it is natural to ask for a description of the monoids which lie in between \mathfrak{S}_k and \mathcal{U}_k . The following theorem summarizes the results in this paper and provides a precise description.

Theorem 1.1 (Theorem 5.3). For k a positive integer, the set of submonoids of \mathcal{U}_k that contain \mathfrak{S}_k forms a distributive lattice with the operations of union and intersection.

Date: May 17, 2024.

Key words and phrases. Uniform block permutations, submonoids, distributive lattice, integer partitions.

(Corollary 5.4) Define a partial order on partitions μ, λ of k that are not equal to 1^k such that $\mu \leq \lambda$ if and only if μ is coarser than λ in refinement order and $\varsigma_{\mu} > \varsigma_{\lambda}$, where ς_{μ} denotes the smallest part of μ not equal to 1. For μ a partition of k, let J_{μ} denote a \mathscr{J} -class of \mathcal{U}_k . Every submonoid S of \mathcal{U}_k that contains \mathfrak{S}_k is of the form

$$S = \mathfrak{S}_k \cup \bigcup_{\mu \in I} J_\mu$$

for some down set I of the partial order \leq .

The group of units of a monoid M is the largest group, $G \subseteq M$, containing the identity element of the monoid. The group of units of the uniform block permutation monoid \mathcal{U}_k is the symmetric group \mathfrak{S}_k . The poset of submonoids of M which contain G [12] does not seem to have such nice properties as in the case of \mathcal{U}_k . For instance, our experiments calculating the lattice of monoids of the dual-inverse semigroup [4] indicate that this lattice of submonoids for k = 5 does not form a distributive lattice.

The paper is organized as follows. In Section 2, we define the necessary notation. Sections 3 and 4 characterize the new partial order on integer partitions used in Theorem 1.1. The connection between the partial order and the lattice of submonoids of \mathcal{U}_k containing symmetric group \mathfrak{S}_k is completed in Section 5. Section 6 concludes with some remarks about the consequences of this result. In particular, we show that the sizes of the \mathscr{J} -classes of the uniform block permutation monoid are sums of squares of dimensions of irreducibles.

Acknowledgement

We thank James East, Jinting Liang, Stuart Margolis, John Rhodes and Bruce Sagan for enlightening discussions.

This work was partially supported by NSF grants DMS–2153998, DMS–2053350 and NSERC/CNSRG.

2. NOTATION

In this section, we introduce notation related to the uniform block permutation monoid. We follow the presentation in [11] and refer the reader to this reference for additional details and references. Fix a positive integer k in this section.

We say that $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_r)$ is a *partition* of k if for all $1 \leq i < r$, λ_i is a positive integer, $\lambda_i \geq \lambda_{i+1}, \lambda_r > 0$ and $\sum_{i=1}^r \lambda_i = k$. If λ is a partition of k, we write $\lambda \vdash k$. We denote by $\mathcal{P}_k = \{\lambda \vdash k\}$ the set of partitions of k. Let $\ell(\lambda) := r$ denote the length of the partition. We also use exponential notation to represent partitions. If the partition has b occurrences of an integer i, we represent it by i^b (e.g. $\lambda = (4, 4, 4, 2, 1, 1, 1, 1)$ is also represented by $1^4 2^1 4^3$).

For a finite set X, we say that π is a *set partition* of X (indicated by $\pi \vdash X$) if $\pi = \{A_1, A_2, \ldots, A_r\}$, where $\emptyset \subsetneq A_i \subseteq X$ for $1 \leqslant i \leqslant r$, $A_i \cap A_j = \emptyset$ if $i \neq j$, and $A_1 \cup A_2 \cup \cdots \cup A_r = X$. Denote by $\ell(\pi) = r$ the length of the set partition. The *type* of a set partition π is $type(\pi) = sort(|A| : A \in \pi)$, where the operation of sort arranges the integers in the list in weakly decreasing order so that the result is a partition.

We say that π is *finer* than τ (equivalently, τ is *coarser* than π) if for every block $A \in \pi$, there is a block $B \in \tau$ such that $A \subseteq B$. The set of set partitions of X forms a lattice under refinement order, where the join operation $\pi \vee \tau$ is defined to be the finest set partition coarser than both π and τ and the meet operation $\pi \wedge \tau$ is the coarsest set partition which is finer than both π and τ .

Let $[k] := \{1, 2, ..., k\}$ and $[\overline{k}] := \{\overline{1}, \overline{2}, ..., \overline{k}\}$. We say that a set partition π of $[k] \cup [\overline{k}]$ is *uniform* if for each $A \in \pi$ the condition $|A \cap [k]| = |A \cap [\overline{k}]|$ holds. Let \mathcal{U}_k denote the set of uniform set partitions of $[k] \cup [\overline{k}]$. For reference, we mention that the number of elements in \mathcal{U}_k is sequence A023998 in the Online Encyclopedia of Integer Sequences [6] and (starting with k = 0) the first few terms of the sequence are

$1, 1, 3, 16, 131, 1496, 22482, 426833, \ldots$

For a set $S \subseteq [k] \cup [\overline{k}]$, let $\overline{S} = \{i : \overline{i} \in S \cap [\overline{k}]\} \cup \{\overline{i} : i \in S \cap [k]\}$. For $\pi \in \mathcal{U}_k$, let $\mathsf{top}(\pi)$ be the set partition of [k] consisting of the blocks $A \cap [k]$ for $A \in \pi$ and $\mathsf{bot}(\pi)$ the set partition of [k] containing the blocks $\overline{A} \cap [k]$ for $A \in \pi$. For $\pi \in \mathcal{U}_k$, we use $\mathsf{type}(\pi)$ to denote $\mathsf{type}(\mathsf{top}(\pi))$, which is a partition of the integer k.

Example 2.1. We have

$$\begin{split} \pi &= \{\{1,3,\bar{1},\bar{2}\},\{2,\bar{4}\},\{4,6,\bar{3},\bar{6}\},\{5,\bar{7}\},\{7,8,9,\bar{5},\bar{8},\bar{9}\}\} \in \mathcal{U}_9,\\ \mathrm{top}(\pi) &= \{\{1,3\},\{2\},\{4,6\},\{5\},\{7,8,9\}\},\\ \mathrm{bot}(\pi) &= \{\{1,2\},\{4\},\{3,6\},\{7\},\{5,8,9\}\},\\ \mathrm{type}(\pi) &= (3,2,2,1,1). \end{split}$$


We will also view elements of \mathcal{U}_k as a graph and use this graph structure to define a monoid product on these elements. We draw the graphs in two rows, the elements $[k] = \{1, 2, \ldots, k\}$ are arranged from left to right in a top row and the elements $[\overline{k}] = \{\overline{1}, \overline{2}, \ldots, \overline{k}\}$ from left to right in a bottom row. An element $\pi \in \mathcal{U}_k$ is represented as a graph, where two vertices are connected with a path if and only if they are in the same set of π . The *diagram* for π represents a class of labeled graphs that have the same connected components.

Consider $\pi, \gamma \in \mathcal{U}_k$ as diagrams. The monoid product $\pi \gamma \in \mathcal{U}_k$ is computed as follows: Stack the graph π on top of γ and identify the bottom vertices of π with the top vertices of γ . Compute the connected components of the three-row diagram and then eliminate the vertices of the middle row from this diagram. The element $\pi \gamma \in \mathcal{U}_k$ has the same connected components as this stacked graph.

Example 2.2. Let $\pi = \{\{1, 2, 7, \overline{2}, \overline{8}, \overline{9}\}, \{3, \overline{1}\}, \{4, 8, \overline{3}, \overline{5}\}, \{5, 9, \overline{4}, \overline{6}\}, \{6, \overline{7}\}\}$ and $\tau = \{\{1, \overline{2}\}, \{2, 3, \overline{1}, \overline{4}\}, \{4, 6, \overline{3}, \overline{5}\}, \{5, \overline{9}\}, \{7, 8, 9, \overline{6}, \overline{7}, \overline{8}\}\}$. The diagram for π is shown below on the left and the diagram for τ is on the right.



Their product $\pi \tau = \{\{1, 2, 4, 6, 7, 8, \overline{1}, \overline{4}, \overline{6}, \overline{7}, \overline{8}, \overline{9}\}, \{3, \overline{2}\}, \{5, 9, \overline{3}, \overline{5}\}\}$ is computed with the graphs:



This product is an extension of the product of permutations in the sense that an element $\pi = \{A_1, \ldots, A_r\} \in \mathcal{U}_k$ with all $|A_i| = 2$ can be identified with a permutation $\sigma \in \mathfrak{S}_k$ via $\pi = \{\{\sigma(1), \overline{1}\}, \{\sigma(2), \overline{2}\}, \ldots, \{\sigma(k), \overline{k}\}\}$. Under this identification the identity permutation is the identity of the monoid, the product for two such elements corresponds to composition of permutations, and the maximal subgroup of \mathcal{U}_k containing the identity is \mathfrak{S}_k .

The set of idempotents of \mathcal{U}_k is denoted $E(\mathcal{U}_k)$. The idempotents are all $\pi \in \mathcal{U}_k$ such that $A \cap [k] = \overline{A} \cap [k]$ for all $A \in \pi$. Each idempotent $\pi \in E(\mathcal{U}_k)$ is therefore determined by the set partition $top(\pi)$ of [k] and we use the notation $e_{top(\pi)}$ to refer to this element. We note the following properties of the idempotents (see for instance [11, Section 2.5]):

(1) The product of two idempotents is

$$e_{\pi}e_{\gamma} = e_{\pi \vee \gamma},$$

where \lor is the join operation in the lattice of set partitions.

(2) \mathfrak{S}_k acts on the set of idempotents $E(\mathcal{U}_k)$ by conjugation: for all $\tau \in \mathfrak{S}_k$ and $e_{\pi} \in E(\mathcal{U}_k)$,

$$\tau e_{\pi} \tau^{-1} = e_{\tau^{-1}(\pi)}$$

Consequently,

$$e_{\pi}\tau = \tau e_{\tau(\pi)}$$
 and $\tau e_{\pi} = e_{\tau^{-1}(\pi)}\tau$.

- (3) Every element of \mathcal{U}_k is of the form τe_{π} for $\tau \in \mathfrak{S}_k$ and $\pi \vdash [k]$.
- (4) If $\tau_1 e_{\pi} = \tau_2 e_{\gamma}$ for $\tau_1, \tau_2 \in \mathfrak{S}_k$, then $\pi = \gamma$.

Let x and y be elements of a monoid M. We say that x and y are \mathscr{J} -equivalent (one of Green's relations on semigroups [5]) if MxM = MyM. This is an equivalence relation and so partitions the elements of M into classes which are called the \mathscr{J} -classes of M. In the case that $M = \mathcal{U}_k$, the \mathscr{J} -classes are indexed by partitions μ of k and are given by $J_{\mu} = \{\pi \in \mathcal{U}_k : type(\pi) = \mu\}$ (for details, see [11, Proposition 3.5]). By the above properties, we also have $J_{\mu} = \{\sigma e_{\gamma}\tau : \sigma, \tau \in \mathfrak{S}_k\}$ for any set partition γ of [k] such that $type(\gamma) = \mu$.

3. A new partial order \leq on integer partitions

We define a new partial order on integer partitions in terms of set partitions in Section 3.1 and relate it to refinement order on integer partitions in Section 3.2.

3.1. Definition of a new partial order on integer partitions.

Definition 3.1. For $\mu, \lambda \vdash k$, define $\mu \preceq \lambda$ if there exist set partitions $\pi_0, \pi_1, \ldots, \pi_\ell \vdash [k]$ of type λ with $\pi_0 \lor \pi_1 \lor \cdots \lor \pi_\ell$ of type μ .




FIGURE 1. Hasse diagram for $(\mathcal{P}_6 \setminus \{1^6\}, \preceq)$.

Remark 3.2. Note that 1^k is incomparable to all other partitions: there is exactly one set partition π of [k] of type 1^k , so $\mu \leq 1^k$ implies $\mu = 1^k$; and if $\pi_0 \lor \pi_1 = \pi$, then $\pi_0 = \pi_1 = \pi$, so $1^k \leq \mu$ implies $\mu = 1^k$. For this reason we will mainly be concerned with the partial order on partitions of k excluding 1^k . We will make a connection between the submonoids of \mathcal{U}_k containing \mathfrak{S}_k and the downsets of the partial order $(\mathcal{P}_k \setminus \{1^k\}, \leq)$.

Proposition 3.3. The relation \leq is a partial order on integer partitions of k. Moreover, if $\mu \leq \lambda$, then μ is coarser than λ as an integer partition.

Proof. We first prove the second statement. Suppose $\mu \leq \lambda$. Then $\mu = \mathsf{type}(\pi_0 \lor \cdots \lor \pi_\ell)$ for some $\pi_0, \ldots, \pi_\ell \vdash [k]$ of type λ . Since $\pi_0 \lor \cdots \lor \pi_\ell$ is obtained by merging blocks of π_0 , its type is obtained by merging parts of $\mathsf{type}(\pi_0) = \lambda$. Hence, μ is coarser than λ .

We now prove that \leq is a partial order.

Reflexivity. $\lambda \leq \lambda$ because there exists a set partition π with $\mathsf{type}(\pi) = \lambda$.

Transitivity. Suppose $\nu \leq \mu$ and $\mu \leq \lambda$. There exist

- $\pi_0, \ldots, \pi_\ell \vdash [k]$ of type λ with type $(\pi_0 \lor \cdots \lor \pi_\ell) = \mu$; and
- $\gamma_0, \ldots, \gamma_m \vdash [k]$ of type μ with type $(\gamma_0 \lor \cdots \lor \gamma_m) = \nu$.

Since $\pi_0 \vee \cdots \vee \pi_\ell$ has type μ , there exist $u_0, \ldots, u_m \in \mathfrak{S}_k$ such that $\gamma_i = u_i(\pi_0 \vee \cdots \vee \pi_\ell) = u_i(\pi_0) \vee \cdots \vee u_i(\pi_\ell)$ for all $0 \leq i \leq m$. Thus,

$$\gamma_0 \lor \cdots \lor \gamma_m = u_0(\pi_0) \lor \cdots \lor u_0(\pi_\ell) \lor \cdots \lor u_m(\pi_0) \lor \cdots \lor u_m(\pi_\ell)$$

is a join of set partitions of type λ , showing that $\nu \leq \lambda$.

Antisymmetry. If $\mu \leq \lambda$ and $\lambda \leq \mu$, then μ is coarser than λ and λ is coarser than μ in the refinement order on integer partitions and so $\mu = \lambda$.

3.2. Comparison with refinement order. The next result describes the precise relationship between \leq and refinement order on integer partitions.

Theorem 3.4. For $\nu \vdash k$ with $\nu \neq 1^k$, let ς_{ν} be the smallest part of ν not equal to 1. Then $\mu \preceq \lambda$ if and only if μ is coarser than λ and $\varsigma_{\mu} \ge \varsigma_{\lambda}$.

The proof is given below after the presentation of a few necessary lemmas.

Recall that if $\mu \leq \lambda$ is a cover relation in the refinement order on integer partitions, then μ is obtained from λ by merging two parts λ_i and λ_j . We first prove that $\mu \prec \lambda$ is a cover relation if at least one of the two merged parts is greater than 1.

Lemma 3.5. If μ is obtained from λ by summing two parts λ_i and λ_j not both equal to 1, then $\mu \prec \lambda$.

Proof. Let $\pi = \{B_1, \ldots, B_\ell\}$ be of type λ with $|B_k| = \lambda_k$ for all $k \in [\ell]$. Fix $b_i \in B_i$ and $b_j \in B_j$ and let

$$\pi' = \left(\{B_1, \ldots, B_\ell\} \setminus \{B_i, B_j\} \right) \cup \left\{ B'_i, B'_j \right\},\$$

where

$$B'_i = (B_i \setminus \{b_i\}) \cup \{b_j\} \text{ and } B'_j = (B_j \setminus \{b_j\}) \cup \{b_i\}.$$

Assume without loss of generality that $|B_i| = \lambda_i \ge 2$. Then the intersection $B_i \cap B'_i$ is nonempty because it contains $B_i \setminus \{b_i\}$. Also, $B_j \cap B'_i \ne \emptyset$ because it contains b_j . Furthermore, $\pi \ne \pi'$. Hence,

$$\pi \vee \pi' = \left(\{B_1, \dots, B_\ell\} \setminus \{B_i, B_j\} \right) \cup \left\{ B_i \cup B_j \right\}$$

is of type μ and so $\mu \prec \lambda$ because $\mathsf{type}(\pi') = \mathsf{type}(\pi) = \lambda$.

If $\mu \prec \lambda$ is not a cover relation, then $\mu \prec \nu \prec \lambda$ for some ν , which implies μ is coarser than ν and ν is coarser than λ in refinement order. But since $\mu \lessdot \lambda$ in refinement order, this is a contradiction. Thus, $\mu \prec \lambda$.

Lemma 3.6. Suppose μ is obtained from $\lambda \neq 1^k$ by merging s ones: explicitly,

$$\lambda = (\lambda_1, \dots, \lambda_m, 1^t)$$
 and $\mu = \operatorname{sort}(\lambda_1, \dots, \lambda_m, s, 1^{t-s}),$

where $\lambda_m > 1$ and $t \ge s$. If $s \ge \lambda_m$, then $\mu \prec \lambda$.

Proof. Let $\pi = \{B_1, \ldots, B_\ell\}$ be a set partition of type λ with $|B_k| = \lambda_k$ for all $k \in [\ell]$. The block B_m has size $\lambda_m \leq s$ and B_{m+1}, \ldots, B_ℓ are singletons, so we can write

 $B_m = \{a_1, a_2, \dots, a_{\lambda_m}\}, \quad B_{m+1} = \{b_1\}, \quad \dots \quad , \quad B_{m+\lambda_m} = \{b_{\lambda_m}\}.$

Let γ be the set partition obtained from π by replacing the set partitions $B_m, B_{m+1}, \ldots, B_{m+\lambda_m}$ by

$$B'_{m} = \{b_{1}, b_{2}, \dots, b_{\lambda_{m}}\}, \quad B'_{m+1} = \{a_{1}\}, \quad \dots \quad , \quad B'_{m+\lambda_{m}} = \{a_{\lambda_{m}}\}$$

Then γ also has type λ and

$$\pi \lor \gamma = \pi \setminus \{B_{m+1}, \dots, B_{m+\lambda_m}\} \cup \{B_{m+1} \cup \dots \cup B_{m+\lambda_m}\}$$

has type $\nu := (\lambda_1, \ldots, \lambda_m, \lambda_m, 1^{t-\lambda_m})$, proving that $\nu \prec \lambda$. Note that $\mu \preceq \nu$ because μ can be obtained from ν by first merging λ_m with 1, then merging $\lambda_m + 1$ with another 1, and so on:

$$\begin{split} \mathbf{A} \succ \nu &= (\lambda_1, \dots, \lambda_m, \lambda_m, 1^{t-\lambda_m}) \\ &\succ \operatorname{sort}(\lambda_1, \dots, \lambda_m, \lambda_m + 1, 1^{t-(\lambda_m+1)}) \\ &\succ \operatorname{sort}(\lambda_1, \dots, \lambda_m, \lambda_m + 2, 1^{t-(\lambda_m+2)}) \\ &\succ \dots \succ \operatorname{sort}(\lambda_1, \dots, \lambda_m, s, 1^{t-s}) = \mu. \end{split}$$

Proof of Theorem 3.4. (\Leftarrow) Suppose that μ is coarser than λ and $\varsigma_{\mu} \geq \varsigma_{\lambda}$.

If μ can be obtained from λ by merging at each step with a part of size at least 2, then $\mu \leq \lambda$ by Lemma 3.5.

Suppose instead that there are parts of μ that are obtained from λ by merging 1s. By first merging ς_{λ} ones (Lemma 3.6), and then successively merging this new part with 1s (Lemma 3.5), it follows that $\mu \leq \lambda$.

 (\Rightarrow) Suppose $\mu \leq \lambda$ and $\varsigma_{\mu} < \varsigma_{\lambda}$. Since $\mu \leq \lambda$, there exist set partitions $\pi_0, \ldots, \pi_{\ell}$ of type λ with $\pi_0 \vee \cdots \vee \pi_{\ell}$ of type μ . This yields a chain in the lattice of set partitions

$$\underbrace{\pi_0}_{\text{type }\lambda} \leqslant \pi_0 \lor \pi_1 \leqslant \cdots \leqslant \underbrace{\pi_0 \lor \pi_1 \lor \cdots \lor \pi_\ell}_{\text{type }\mu}.$$

Since $\pi_0 \vee \cdots \vee \pi_\ell$ contains a block of size ς_μ and π_0 does not, there exists $i \in [\ell]$ such that $\gamma := \pi_0 \vee \cdots \vee \pi_{i-1}$ does not contain a block of size ς_μ and $\gamma \vee \pi_i$ does.

This block of size ς_{μ} is necessarily obtained by merging singleton blocks of π_i since all other blocks of π_i are of size greater than ς_{μ} . Hence, there exist blocks $C_1, \ldots, C_{\varsigma_{\mu}-1} \in \gamma$ and singletons $B_1, \ldots, B_{\varsigma_{\mu}} \in \pi_i$ such that $B_1 \sim_{C_1} B_2 \sim_{C_2} \cdots \sim_{C_{\varsigma_{\mu}-1}} B_{\varsigma_{\mu}}$, where $B \sim_C B'$ means C intersects both B and B'.

Let us show that each C_i equals $B_1 \cup \cdots \cup B_{\varsigma_{\mu}}$. Since B_1 and B_2 are singletons and $B_1 \sim_{C_1} B_2$, we have that $B_1 \subseteq C_1$ and $B_2 \subseteq C_1$. Similarly, $B_2, B_3 \subseteq C_2$. But since $C_1 \cap C_2 = \emptyset$ if $C_1 \neq C_2$, it follows that $C_2 = C_1$ because both contain B_2 . Continuing in this way, $C_1 = \cdots = C_{\varsigma_{\mu}-1}$, and $C_1 \supseteq B_1 \cup \cdots \cup B_{\varsigma_{\mu}}$. Since C_1 is contained in the block of $\gamma \vee \pi_i$ that contains B_1 , it follows that $C_1 \subseteq B_1 \cup \cdots \cup B_{\varsigma_{\mu}}$.

Thus, γ contains a block of size ς_{μ} , contradicting the choice of γ .

4. Cover relations for \preceq

The goal of this section is the characterization of the cover relations of \leq .

Theorem 4.1. Let μ and λ be two partitions of k. Then $\mu \prec \lambda$ if and only if either: (1) μ is obtained from λ by merging two parts λ_i and λ_j that are not both 1; (2) there exist integers $t \ge s \ge 2$ such that $\lambda_{m-1} \ge s$,

$$\lambda = (\lambda_1, \dots, \lambda_{m-1}, s, \underbrace{1, \dots, 1}_{t}) \quad and \quad \mu = (\lambda_1, \dots, \lambda_{m-1}, s, s, \underbrace{1, \dots, 1}_{t-s}).$$

4.1. Proof that the relations in Theorem 4.1 are cover relations. It follows directly

from Lemma 3.5 that the first form of the relations in Theorem 4.1 are cover relations.

Suppose there are integers $t \ge s \ge 2$ and $\lambda_{m-1} \ge s$ such that

$$\lambda = (\lambda_1, \dots, \lambda_{m-1}, s, 1^t)$$
 and $\mu = (\lambda_1, \dots, \lambda_{m-1}, s, s, 1^{t-s})$

By Lemma 3.6, it follows that $\mu \prec \lambda$. If this is not a cover relation, then there exists ν such that $\mu \prec \nu \prec \lambda$. Then $\nu = (\lambda_1, \ldots, \lambda_{m-1}, s, r, 1^{t-r})$ with 1 < r < s. By Theorem 3.4, the smallest part of ν that is not equal to 1 (this is r) is greater than or equal to the smallest part of λ that is not equal to 1 (which is s). This implies $r \ge s > r$, a contradiction.

4.2. Proof that all cover relations for \leq are of the form specified in Theorem 4.1. We begin by introducing some notation. For $\lambda = (\lambda_1, \ldots, \lambda_\ell)$ and $I \subseteq [\ell]$, define

$$\lambda_I = \sum_{i \in I} \lambda_i$$
 and $\lambda^I = (\lambda_i : i \in I) \vdash \lambda_I.$

If $\mu < \lambda$ in refinement order, then we can write

 $\mu = \operatorname{sort}(\lambda_{I_1}, \lambda_{I_2}, \dots, \lambda_{I_d}) \text{ for some } \{I_1, \dots, I_d\} \vdash [\ell].$ (4.1)

Furthermore, we order the sets I_1, \ldots, I_d so that

$$\max(\lambda^{I_1}) \ge \cdots \ge \max(\lambda^{I_d})$$

and

$$\lambda_{I_j} \ge \lambda_{I_{j+1}}$$
 if $\max(\lambda^{I_j}) = \max(\lambda^{I_{j+1}})$.

Note that this expression is not necessarily unique.

Example 4.2. For $\lambda = (6, 5, 4, 4, 3, 2, 1, 1, 1)$ and $\mu = (12, 12, 3)$, we can take

$$\lambda^{\{1,3,6\}} = (6,4,2), \quad \lambda^{\{2,4,7,8,9\}} = (5,4,1,1,1), \quad \lambda^{\{5\}} = (3),$$

or

$$\lambda^{\{1,4,6\}} = (6,4,2), \quad \lambda^{\{2,3,5\}} = (5,4,3), \quad \lambda^{\{7,8,9\}} = (1,1,1).$$

We are now ready to prove that all cover relations for \leq are of the form specified in Theorem 4.1. Suppose $\mu \prec \lambda$. Then μ is coarser than λ in refinement order, so there exists a way of merging the blocks of λ to obtain μ , as in (4.1).

Case 1: Suppose every λ^{I_j} in (4.1) contains a part greater than 1. Construct a chain by merging parts of λ as follows (*cf.* Example 4.3):

- if λ^{I_d} = (λ_{i1} ≥ · · · ≥ λ_{it}), replace λ_{i1} and λ_{i2} by λ_{i1} + λ_{i2};
 then replace λ_{i1} + λ_{i2} and λ_{i3} by λ_{i1} + λ_{i2} + λ_{i3}; and so on.

8

Proceed similarly with I_{d-1}, \ldots, I_1 . Since every step involves merging with a block of size greater than 1, we obtain a \leq -chain from μ to λ by Lemma 3.5. Since $\mu \prec \lambda$, it follows that d = 1 and μ is obtained from λ by merging two blocks that are not both 1.

Example 4.3. Let $\lambda = (6, 5, 4, 4, 3, 2, 1, 1, 1)$ and $\mu = (12, 12, 3)$. For

$$\lambda^{\{1,3,6\}} = (6,4,2), \quad \lambda^{\{2,4,7,8,9\}} = (5,4,1,1,1), \quad \lambda^{\{5\}} = (3)$$

one first merges λ_2 and λ_4 ; then λ_2, λ_4 , and λ_7 ; and so on:

$$\begin{array}{c} (\overset{1}{6},\overset{2}{5},\overset{3}{4},\overset{4}{3},\overset{5}{2},\overset{6}{1},\overset{7}{1},\overset{8}{1},\overset{9}{1}) \succ (\overset{2,4}{9},\overset{1}{6},\overset{3}{4},\overset{5}{3},\overset{6}{2},\overset{7}{1},\overset{8}{1},\overset{9}{1}) \succ (\overset{2,4,7}{10},\overset{1}{6},\overset{3}{4},\overset{5}{2},\overset{6}{1},\overset{8}{1},\overset{9}{1}) \\ \succ (\overset{2,4,7,8}{11},\overset{1}{6},\overset{3}{4},\overset{5}{2},\overset{6}{1}) \succ (\overset{2,4,7,8,9}{12},\overset{1}{6},\overset{3}{4},\overset{5}{3},\overset{6}{2},\overset{7}{1},\overset{8}{1},\overset{9}{1}) \succ (\overset{2,4,7,8,9}{11},\overset{1}{3},\overset{5}{5},\overset{6}{6},\overset{8}{4},\overset{9}{3},\overset{2}{2}) \succ (\overset{2,4,7,8,9}{11},\overset{1}{3},\overset{5}{5},\overset{6}{6},\overset{8}{4},\overset{9}{3},\overset{2}{2}) \succ (12,12,3). \end{array}$$

Case 2: Suppose $\lambda^{I_d} = 1^r$ in (4.1) with r > 1. By Theorem 3.4,

$$r = |\lambda^{I_d}| \geqslant \varsigma_\mu \geqslant \varsigma_\lambda =: s.$$

We construct a \leq -chain as follows. By hypothesis, $\lambda = (\lambda_1, \ldots, \lambda_{m-1}, s, 1^t)$ with $t \ge r$. By Lemma 3.6, since λ contains s, merging s ones yields the following relation

$$\lambda = (\lambda_1, \dots, \lambda_{m-1}, s, 1^t) \succ (\lambda_1, \dots, \lambda_{m-1}, s, s, 1^{t-s}).$$

Since $s \ge 2$, merging s with 1 is a \leq -relation by Lemma 3.5:

$$(\lambda_1,\ldots,\lambda_{m-1},s,s,1^{t-s}) \succ \operatorname{sort}(\lambda_1,\ldots,\lambda_{m-1},s+1,s,1^{t-s-1}).$$

Continuing to merge the new part with 1s, we get the chain

$$\operatorname{sort}(\lambda_1, \dots, \lambda_{m-1}, s+1, s, 1^{t-(s+1)}) \succ \operatorname{sort}(\lambda_1, \dots, \lambda_{m-1}, s+2, s, 1^{t-(s+2)})$$
$$\succ \dots \succ \operatorname{sort}(\lambda_1, \dots, \lambda_{m-1}, r, s, 1^{t-r}) =: \nu^{(d)}.$$

Example 4.4. Consider $\lambda = (6, 5, 4, 3, 1, 1, 1, 1), \mu = (10, 8, 4)$, and

$$\lambda^{I_1} = (6,4), \quad \lambda^{I_2} = (5,3), \quad \lambda^{I_3} = (1,1,1,1).$$

One obtains the chain

$$(6,5,4,3,\underline{1,1,1},1) \succ (6,5,4,3,\underline{3,1}) \succ (6,5,4,4,3) =: \nu^{(3)}$$

If $\lambda^{I_{d-1}}$ is also of the form $1^{r'}$, then $r' \ge r$ because I_1, \ldots, I_d are ordered so that $\lambda_{I_j} \ge \lambda_{I_{j+1}}$ when $\max(\lambda^{I_j}) = \max(\lambda^{I_{j+1}})$. Therefore, we can apply the above procedure to I_{d-1} . Continuing in this way, we obtain a chain

$$\nu^{(a)} \prec \nu^{(a+1)} \prec \cdots \prec \nu^{(d)} \prec (\lambda_1, \dots, \lambda_{m-1}, s, s, 1^{t-s}) \prec \lambda$$

where $\nu^{(j)}$ is obtained from $\nu^{(j+1)}$ by merging the parts in λ^{I_j} ; and a is such that $\max(\lambda^{I_a}) = 1$ and either a = 1 or $\max(\lambda^{I_{a-1}}) > 1$.

If a = 1, then $\nu^{(a)} = \mu$. In the other case, proceed as in *Case 1* to merge the parts of $\lambda^{I_{a-1}}, \ldots, \lambda^{I_1}$ (i.e., starting with the maximal element in λ^{I_j}). Thus, in both cases we obtain a chain for \leq of the form

$$\mu = \nu^{(1)} \prec \cdots \prec \nu^{(a)} \prec \cdots \prec \nu^{(d)} \prec (\lambda_1, \dots, \lambda_{m-1}, s, s, 1^{t-s}) \prec \lambda.$$

Since $\mu \prec \lambda$, this is a contradiction unless $\mu = (\lambda_1, \dots, \lambda_{m-1}, s, s, 1^{t-s})$.

5. The submonoids of \mathcal{U}_k containing \mathfrak{S}_k

In this section, we prove that the lattice of submonoids of \mathcal{U}_k containing \mathfrak{S}_k is distributive. We begin by showing that these submonoids are unions of \mathscr{J} -classes of \mathcal{U}_k .

Lemma 5.1. If S is a submonoid of \mathcal{U}_k that contains \mathfrak{S}_k , then S is a union of \mathcal{J} -classes of \mathcal{U}_k .

Proof. Suppose S is a submonoid of \mathcal{U}_k that contains \mathfrak{S}_k , and let J_λ be a \mathscr{J} -class of \mathcal{U}_k . If $S \cap J_\lambda \neq \emptyset$, then there exist $u, v \in \mathfrak{S}_k$ and $\pi \vdash [k]$ of type λ such that $ue_{\pi}v \in S$. Since $u^{-1}, v^{-1} \in \mathfrak{S}_k \subseteq S$, we have

$$e_{\pi} = u^{-1}(ue_{\pi}v)v^{-1} \in S,$$

from which it follows that $\sigma e_{\pi} \tau \in S$ for all $\sigma, \tau \in \mathfrak{S}_k$. Consequently,

$$J_{\lambda} = \{ \sigma e_{\pi} \tau : \sigma, \tau \in \mathfrak{S}_k \} \subseteq S.$$

Proposition 5.2. Let $\langle \mathfrak{S}_k, e_\pi \rangle$ be the submonoid of \mathcal{U}_k generated by \mathfrak{S}_k and e_π .

- (1) For $\gamma \vdash [k]$ with type $(\gamma) \neq 1^k$, we have $e_{\gamma} \in \langle \mathfrak{S}_k, e_{\pi} \rangle$ if and only if type $(\gamma) \preceq type(\pi)$.
- (2) The monoid $\langle \mathfrak{S}_k, e_\pi \rangle$ is the union of \mathfrak{S}_k and \mathscr{J} -classes J_μ with $\mu \leq \mathsf{type}(\pi)$:

$$\langle \mathfrak{S}_k, e_\pi \rangle = \mathfrak{S}_k \cup \bigcup_{\mu \preceq \mathsf{type}(\pi)} J_\mu.$$

Proof. Every element of $\langle \mathfrak{S}_k, e_\pi \rangle$ is either in \mathfrak{S}_k or is a product of the form

 $u_0 e_{\pi} u_1 e_{\pi} \cdots u_{\ell-2} e_{\pi} u_{\ell-1} e_{\pi} u_{\ell} e_{\pi} u_{\ell+1} \qquad \text{with } u_0, u_1, \dots, u_{\ell+1} \in \mathfrak{S}_k \text{ for some } \ell \geq 0.$

Since type $(\gamma) \neq 1^k$, then e_{γ} is of this form. Setting $v_{\ell+1} = u_{\ell+1}$ and $v_i = u_i v_{i+1}$ for $0 \leq i \leq \ell$ so that $u_i = v_i v_{i+1}^{-1}$, the above becomes

$$v_0(v_1^{-1}e_{\pi}v_1)(v_2^{-1}e_{\pi}v_2)\cdots(v_{\ell-1}^{-1}e_{\pi}v_{\ell-1})(v_{\ell}^{-1}e_{\pi}v_{\ell})(v_{\ell+1}^{-1}e_{\pi}v_{\ell+1})$$

= $v_0 e_{v_1(\pi)}e_{v_2(\pi)}\cdots e_{v_{\ell}(\pi)}e_{v_{\ell+1}(\pi)}$
= $v_0 e_{v_1(\pi)\vee v_2(\pi)\vee\cdots\vee v_{\ell}(\pi)\vee v_{\ell+1}(\pi).$

If $e_{\gamma} = v_0 e_{v_1(\pi) \vee v_2(\pi) \vee \cdots \vee v_{\ell}(\pi) \vee v_{\ell+1}(\pi)}$, then $\gamma = v_1(\pi) \vee \cdots \vee v_{\ell+1}(\pi)$ by item (4) in Section 2. Hence γ is a join of set partitions of type $\mathsf{type}(\pi)$; that is, $\mathsf{type}(\gamma) \preceq \mathsf{type}(\pi)$.

Conversely, if $\mathsf{type}(\gamma) \leq \mathsf{type}(\pi)$, then $\gamma = \pi_1 \vee \cdots \vee \pi_{\ell+1}$ for some set partitions $\pi_1, \ldots, \pi_{\ell+1}$ with $\mathsf{type}(\pi_i) = \mathsf{type}(\pi)$. Each π_i is equal to $v_i(\pi)$ for some $v_i \in \mathfrak{S}_k$, and so $\gamma = v_1(\pi) \vee \cdots \vee v_{\ell+1}(\pi)$. Thus,

$$e_{\gamma} = e_{v_1(\pi) \lor \dots \lor v_{\ell+1}(\pi)} = v_1 e_{\pi} v_1^{-1} \cdots v_{\ell+1} e_{\pi} v_{\ell+1}^{-1} \in \langle \mathfrak{S}_k, e_{\pi} \rangle.$$

Theorem 5.3. Let S_1 and S_2 be two submonoids of \mathcal{U}_k that contain \mathfrak{S}_k . Then $S_1 \cup S_2$ is also a submonoid of \mathcal{U}_k .

Proof. Let $\tau e_{\pi} \in S_1$ and $\sigma e_{\gamma} \in S_2$. Then

$$\tau e_{\pi} \cdot \sigma e_{\gamma} = \tau \sigma e_{\sigma(\pi)} e_{\gamma} = (\tau \sigma) e_{\sigma(\pi) \lor \gamma}.$$

We will show that $e_{\sigma(\pi)\vee\gamma}$ is contained in S_1 or S_2 .

Let

$$\mu:=\operatorname{type}(\sigma(\pi)\vee\gamma),\quad \lambda:=\operatorname{type}(\pi),\quad \nu:=\operatorname{type}(\gamma)$$

If $\lambda = 1^k$, then $\sigma(\pi) \vee \gamma = \gamma$ because $\sigma(\pi) = \pi$ is the unique set partition of type 1^k , from which it follows that $\tau \sigma e_{\sigma(\pi) \vee \gamma} \in S_2$. Similarly, if $\nu = 1^k$, then $\sigma(\pi) \vee \gamma = \sigma(\pi)$ and $\tau e_{\pi} \sigma e_{\sigma(\pi)} \in S_1$.

Suppose $\mu, \lambda \neq 1^k$. Since we want to show that $e_{\sigma(\pi)\vee\gamma} \in S_1 \cup S_2$, it suffices to prove $e_{\sigma(\pi)\vee\gamma} \in \langle \mathfrak{S}_k, e_\pi \rangle \cup \langle \mathfrak{S}_k, e_\gamma \rangle$ because $\langle \mathfrak{S}_k, e_\pi \rangle \subseteq S_1$ and $\langle \mathfrak{S}_k, e_\gamma \rangle \subseteq S_2$. By Proposition 5.2, $e_{\sigma(\pi)\vee\gamma} \in \langle \mathfrak{S}_k, e_\pi \rangle$ if and only if $\mu \preceq \lambda$ and $e_{\sigma(\pi)\vee\gamma} \in \langle \mathfrak{S}_k, e_\gamma \rangle$ if and only if $\mu \preceq \nu$, so it suffices to show $\mu \preceq \lambda$ or $\mu \preceq \nu$.

Suppose $\mu \not\preceq \lambda$ and $\mu \not\preceq \nu$. Note that μ is coarser than λ and μ is coarser than ν because $\sigma(\pi) \lor \gamma$ is obtained by merging blocks of $\sigma(\pi)$ and by merging blocks of γ . Then by Theorem 3.4 we have that $\varsigma_{\mu} < \varsigma_{\nu}$ and $\varsigma_{\mu} < \varsigma_{\lambda}$.

Since $\sigma(\pi) \vee \gamma$ has type μ and is obtained by merging blocks in $\sigma(\pi)$, it follows that it contains a block of size ς_{μ} that can only be obtained by merging singleton blocks in $\sigma(\pi)$. This implies that γ contains a block of size ς_{μ} (*cf.* the proof of Theorem 3.4). Since γ contains a block of size ς_{μ} , it follows that ν has a part equal to ς_{μ} , implying that $\varsigma_{\mu} \geq \varsigma_{\nu}$, a contradiction because $\varsigma_{\mu} < \varsigma_{\nu}$.

Corollary 5.4. Let S be a submonoid of \mathcal{U}_k containing \mathfrak{S}_k and let E(S) be its set of idempotents. Then $I := \{ \mathsf{type}(\pi) : e_\pi \in E(S), e_\pi \neq id \}$ is a down set for the partial order \preceq and $S = \mathfrak{S}_k \cup \bigcup_{\lambda \in I} J_{\lambda}$.

Conversely, if I is a downset of $(\mathcal{P}_k \setminus \{1^k\}, \preceq)$, then $\mathfrak{S}_k \cup \bigcup_{\lambda \in I} J_\lambda$ is a submonoid of \mathcal{U}_k .

Proof. Let us first prove that I is a down set. Suppose $\mu \leq \mathsf{type}(\pi)$ for some $\pi \in E(S)$. By Proposition 5.2, $e_{\gamma} \in \langle \mathfrak{S}_k, e_{\pi} \rangle$ for any $\gamma \vdash [k]$ of type μ . But if $e_{\gamma} \in \langle \mathfrak{S}_k, e_{\pi} \rangle$, then $e_{\gamma} \in E(S)$, and so $\mu = \mathsf{type}(\gamma) \in I$.

Every element of S is of the form σe_{π} with $\sigma \in \mathfrak{S}_k$ and $e_{\pi} \in E(S)$, so $S = \langle \mathfrak{S}_k, E(S) \rangle$. By Theorem 5.3, $\langle \mathfrak{S}_k, E(S) \rangle = \bigcup_{e_{\pi} \in E(S)} \langle \mathfrak{S}_k, e_{\pi} \rangle$, since both are the smallest submonoids containing \mathfrak{S}_k and E(S). By Proposition 5.2,

$$S = \bigcup_{e_\pi \in E(S)} \langle \mathfrak{S}_k, e_\pi \rangle = \bigcup_{e_\pi \in E(S)} \left(\mathfrak{S}_k \cup \bigcup_{\lambda \preceq \mathsf{type}(\pi)} J_\lambda \right) = \mathfrak{S}_k \cup \bigcup_{\lambda \in I} J_\lambda.$$

Now conversely, if I is a downset of $(\mathcal{P}_k \setminus \{1^k\}, \preceq)$, then by Proposition 5.2 part (2), for each e_{π} with $\mathsf{type}(\pi) \in I$, $\langle \mathfrak{S}_k, e_{\pi} \rangle \subseteq \mathfrak{S}_k \cup \bigcup_{\lambda \in I} J_{\lambda}$ and hence

$$\mathfrak{S}_k \cup \bigcup_{\lambda \in I} J_\lambda = \bigcup_{\mathsf{type}(\pi) \in I} \left< \mathfrak{S}_k, e_\pi \right> \ .$$

Then by Theorem 5.3, $\mathfrak{S}_k \cup \bigcup_{\lambda \in I} J_\lambda$ is a submonoid of \mathcal{U}_k .

Theorem 5.3 implies that the poset of submonoids of \mathcal{U}_k containing \mathfrak{S}_k under union and intersection is a distributive lattice. The lattices for k = 4, 5, 6 are presented in Figure 2.




FIGURE 2. The distributive lattices of submonoids of \mathcal{U}_k containing the symmetric group \mathfrak{S}_k for k = 4, 5, 6. Each submonoid is labeled by a box containing an antichain of the poset $(\mathcal{P}_k \setminus \{1^k\}, \preceq)$. The antichains of \preceq are in bijection with the downsets of this poset (see [13, Section 3.4]) and in turn are in bijection with the submonoids by Corollary 5.4. Below each box is the size of the corresponding submonoid.

6. Concluding Remarks

In this section, we state some interesting consequences of the characterization of the lattice of submonoids of \mathcal{U}_k that contain \mathfrak{S}_k .

6.1. The sizes of \mathscr{J} -classes are sums of squares of dimensions of irreducibles. For monoids M, N such that $\mathfrak{S}_k \subseteq M \subseteq N \subseteq \mathcal{U}_k$, the difference $N \setminus M$ is a union of \mathscr{J} -classes. If N covers M in the lattice of monoids, then $N = M \cup J_{\mu}$ for some partition μ .

We note that $J_{1^k} = \mathfrak{S}_k$. More generally, for $\lambda = 1^{a_1} 2^{a_2} \cdots k^{a_k} \vdash k$, let $\mathsf{sp}_k(\lambda)$ be the number of set partitions of [k] such that the type is equal to λ , then

(6.1)
$$\mathsf{sp}_k(\lambda) = \frac{k!}{a_1! \cdots a_k! (1!)^{a_1} (2!)^{a_2} \cdots (k!)^{a_k}}$$

Since every element of J_{λ} can be considered as a bijection from a part of size *i* in the top row to a part of size *i* in the bottom row, the number of elements of J_{λ} is equal to $sp_k(\lambda)^2a_1!a_2!\cdots a_k!$.

Recall from [11], that the irreducible representations of \mathcal{U}_k are indexed by sequences of partitions, $\vec{\lambda} = (\lambda^{(1)}, \lambda^{(2)}, \dots, \lambda^{(k)})$, such that $k = |\lambda^{(1)}| + 2|\lambda^{(2)}| + \dots + k|\lambda^{(k)}|$. Let I_k denote this set of sequences of partitions and let $W_{\mathcal{U}_k}^{\vec{\lambda}}$ represent the isomorphism class of an irreducible \mathcal{U}_k module indexed by $\vec{\lambda} \in I_k$.

Now let f^{λ} denote the number of standard tableaux of shape λ and define the $\overrightarrow{type}(\vec{\lambda}) = 1^{|\lambda^{(1)}|} 2^{|\lambda^{(2)}|} \cdots k^{|\lambda^{(k)}|}$. Then [11, Corollary 3.19] states that

(6.2)
$$\dim W_{\mathcal{U}_k}^{\vec{\lambda}} = \mathsf{sp}_k(\overrightarrow{\mathsf{type}}(\vec{\lambda})) f^{\lambda^{(1)}} f^{\lambda^{(2)}} \cdots f^{\lambda^{(k)}} .$$

Proposition 6.1. For $\mu \vdash k$, we have


$$J_{\mu}| = \sum_{\substack{\vec{\lambda} \\ \text{type}(\vec{\lambda}) = \mu}} (\dim W_{\mathcal{U}_k}^{\vec{\lambda}})^2 \ .$$

Proof. If $\mu = 1^{a_1} 2^{a_2} \cdots k^{a_k}$, then $\overrightarrow{\mathsf{type}}(\vec{\lambda}) = \mu$ implies that $|\lambda^{(i)}| = a_i$. Recall that $a! = \sum_{\lambda \vdash a} (f^{\lambda})^2$. Therefore by (6.2),

$$\sum_{\substack{\vec{\lambda} \in I_k \\ \overrightarrow{\mathsf{type}}(\vec{\lambda}) = \mu}} (\dim W_{\mathcal{U}_k}^{\vec{\lambda}})^2 = \mathsf{sp}_k(\mu)^2 \sum_{\substack{\vec{\lambda} \in I_k \\ \lambda^{(i)} \vdash a_i}} (f^{\lambda^{(1)}})^2 (f^{\lambda^{(2)}})^2 \cdots (f^{\lambda^{(k)}})^2,$$

where the sum on the right hand side is over all $\vec{\lambda} \in I_k$ such that $\lambda^{(i)}$ is a partition of a_i . The conclusion that this expression is equal to $|J_{\mu}|$ follows from the enumeration $|J_{\mu}| = \operatorname{sp}_k(\mu)^2 a_1! a_2! \cdots a_k!$ of the elements of the \mathscr{J} -class described above. \Box

6.2. The order \leq on partitions is not Cohen–Macaulay. Ziegler [16] proved that for $k \geq 19$, the poset of integers ordered by refinement is not Cohen–Macaulay (as it had been previously conjectured). His proof produces an interval of the poset at k = 19 which is an obstruction to the poset being Cohen–Macaulay and states that larger examples exist at every k.



We remark that this same interval also occurs in our poset and hence this poset will also not be Cohen–Macaulay.

Moreover, Ziegler [16] proved that for $k \ge 111$ the Möbius function is not alternating. His proof also applies to our poset.

6.3. The number of submonoids of \mathcal{U}_k that contain \mathfrak{S}_k . Corollary 5.4 can be used to compute the number of submonoids of \mathcal{U}_k containing \mathfrak{S}_k (i.e. $n_k := \#\{M : \mathfrak{S}_k \subseteq M \subseteq \mathcal{U}_k\}$) for small k by computing the number of antichains of the poset of partitions of k not equal to 1^k under the order \preceq . Using the poset functionality of SAGEMATH [14] we were able to compute the following table:

									9			12	13	14
n_k	1	2	3	6	10	31	63	287	1099	8640	62658	1546891	29789119	2525655957

The sequence has been submitted to [6] and is now sequence number A371505.

References

- Marcelo Aguiar and Rosa C. Orellana. The Hopf algebra of uniform block permutations. Journal of Algebraic Combinatorics, 28(1):115–138, 2008.
- [2] Desmond G. FitzGerald. A presentation for the monoid of uniform block permutations. Bull. Austral. Math. Soc., 68(2):317–324, 2003.
- [3] Desmond G. FitzGerald. Factorizable inverse monoids. In Semigroup forum, volume 80, pages 484–509. Springer, 2010.
- [4] Desmond G. FitzGerald and Jonathan Leech. Dual symmetric inverse monoids and representation theory. Journal of the Australian Mathematical Society, 64(3):345–367, 1998.
- [5] James A. Green. On the structure of semigroups. Annals of Mathematics, pages 163–172, 1951.
- [6] OEIS Foundation Inc. The On-Line Encyclopedia of Integer Sequences, 2019. [Online].
- [7] Masashi Kosuda. Characterization for the party algebras. Ryukyu Math. J., 13:7–22, 2000.
- [8] Masashi Kosuda. Party algebra and construction of its irreducible representations. In Formal Power Series and Algebraic Combinatorics (FPSAC01), Tempe, Arizona (USA), pages 20–26, 2001.
- [9] Masashi Kosuda. Irreducible representations of the party algebra. Osaka J. Math., 43(2):431–474, 2006.
- [10] Hiroshi Naruse. Characters of party algebras. slides of talk for Workshop on Cellular and Diagram Algebras in Mathematics and Physics at Oxford Univ., dated 04-04-2005, 2005.
- [11] Rosa Orellana, Franco Saliola, Anne Schilling, and Mike Zabrocki. Plethysm and the algebra of uniform block permutations. Algebraic Combinatorics, 5(5):1165–1203, 2022.
- [12] Lev N. Shevrin and Alexander J. Ovsyannikov. Semigroups and their subsemigroup lattices, volume 379. Springer Science & Business Media, 2013.
- [13] Richard P. Stanley. *Enumerative Combinatorics*, volume 1. Cambridge University Press, 2nd edition, 2011.
- [14] W.A. Stein et al. Sage Mathematics Software (Version 10.0). The Sage Development Team, 2023. http://www.sagemath.org.
- [15] Kenichiro Tanabe. On the centralizer algebra of the unitary reflection group G(m, p, n). Nagoya Math. J., 148:113–126, 1997.
- [16] G. Ziegler. On the poset of partitions of an integer. Journal of Combinatorial Theory, Series A, 42(2):215–222, 1986.

(R. Orellana) MATHEMATICS DEPARTMENT, DARTMOUTH COLLEGE, HANOVER, NH 03755, U.S.A. *Email address:* Rosa.C.Orellana@dartmouth.edu *URL:* https://math.dartmouth.edu/~orellana/

(F. Saliola) LACIM, DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ DU QUÉBEC À MONTRÉAL, CANADA

Email address: saliola.franco@uqam.ca

(A. Schilling) Department of Mathematics, University of California, One Shields Avenue, Davis, CA 95616-8633, U.S.A.

Email address: anne@math.ucdavis.edu
URL: http://www.math.ucdavis.edu/~anne

(M. Zabrocki) Department of Mathematics and Statistics, York University, 4700 Keele Street, Toronto, Ontario M3J 1P3, Canada

Email address: zabrocki@mathstat.yorku.ca URL: http://garsia.math.yorku.ca/~zabrocki/