

SEGRE SURFACES AND GEOMETRY OF THE PAINLEVÉ EQUATIONS

NALINI JOSHI, MARTA MAZZOCCO, AND PIETER ROFFELSEN

ABSTRACT. In this paper, we consider a six parameter family of affine Segre surfaces embedded in \mathbb{C}^6 . For generic values of the parameters, this family is associated to the q -difference sixth Painlevé equation qP_{VI} . We show that different limiting forms of this family give Segre surfaces that are isomorphic as affine varieties to the the monodromy manifolds of each Painlevé differential equation.

CONTENTS

1. Introduction	1
2. The affine Segre surface of the qP_{VI} equation	5
3. The limit of \mathcal{Z}_q as $q \uparrow 1$ and Jimbo-Fricke cubic for P_{VI}	18
4. The other Painlevé differential equations	42
5. Blow-downs of affine cubic surfaces	46
6. Poisson structure	62
7. Conclusion	64
References	65

1. INTRODUCTION

In this paper, we consider a family of affine Segre surfaces defined, for given $\mu_1, \dots, \mu_6 \in \mathbb{C}, \lambda_1, \lambda_2 \in \mathbb{C}^*$, by

$$\begin{cases} z_1 + z_2 + z_3 + z_4 + z_5 + z_6 = 0, \\ \mu_1 z_1 + \mu_2 z_2 + \mu_3 z_3 + \mu_4 z_4 + \mu_5 z_5 + \mu_6 z_6 - 1 = 0, \\ z_3 z_4 - z_1 z_2 \lambda_1 = 0, \\ z_5 z_6 - z_1 z_2 \lambda_2 = 0. \end{cases} \quad (1.1)$$

We show that limiting forms of this family give Segre surfaces as monodromy manifolds of the Painlevé equations. This result is compelling on two levels. First, the well-known monodromy manifolds of the differential Painlevé equations are cubic surfaces. Second, the geometry of each Segre surface turns out to have a deep connection to distinguished solutions of the corresponding Painlevé equation and provides a natural Poisson structure.

The generic family of surfaces (1.1), which we will denote in this paper by \mathcal{Z}_q , is associated with a *difference* equation known as the q -difference sixth Painlevé

2020 *Mathematics Subject Classification.* 34M55; 39A13 14J10 .

Key words and phrases. Segre surfaces; Painlevé equations; Monodromy manifold .

NJ's ORCID is 0000-0001-7504-4444, MM's is 0000-0001-9917-2547, PR's is 0000-0002-0461-7580.

equation qP_{VI} (2.19) [26]. The name is based on the continuum limit of this difference equation when it becomes the celebrated sixth differential Painlevé equation P_{VI} (3.1).

Like P_{VI} , qP_{VI} has been associated with conformal field theory and algebraic geometry. The latter equation has also been related to the classical study of linear q -difference equations, series expansions and connection problems; see [25], [31], [38], [26] and references therein.

Although the equations (1.1) defining \mathcal{Z}_q appear to contain 8 parameters, in Corollary 2.12 we show that, up to affine equivalence, they define a six-parameter family of embedded affine Segre surfaces in \mathbb{C}^6 . It is worth noting that \mathcal{Z}_q arises from the Riemann-Hilbert problem associated with qP_{VI} . See [26] for details of its explicit derivation. We prove that the members of the \mathcal{Z}_q family are embedded affine Segre surfaces of generic type. This in particular fills a gap stated as an open problem in [40], see Remark 2.14.

A large amount of work has been devoted to the differential Painlevé equations. For $(\omega_1, \omega_2, \omega_3, \omega_4) \in \mathbb{C}^4$, the monodromy manifold of P_{VI} is well-known to be given by the Jimbo-Fricke family of affine cubic surfaces [16], [24] defined by

$$\begin{cases} x_1 x_2 x_3 + x_1^2 + x_2^2 + x_3^2 + \omega_1 x_1 + \omega_2 x_2 + \omega_3 x_3 + \omega_4 = 0, \\ (x_1, x_2, x_3) \in \mathbb{C}^3. \end{cases} \quad (1.2)$$

Monodromy manifolds of the remaining Painlevé equations have been unified and collected in [49] and in [9] it was shown how to obtain them as limits of (1.2), corresponding to confluence limits of the Painlevé equations. All such monodromy manifolds are cubic surfaces whose projective completion has a triangle of lines at infinity.

Since the continuum limit of qP_{VI} leads to P_{VI} , and confluence limits of the latter give rise to the remaining Painlevé equations, a natural question is whether Segre surfaces also arise as monodromy manifolds of the Painlevé differential equations. Motivated by this question, we show how to obtain explicit Segre surfaces in \mathbb{C}^6 for each Painlevé equation, from appropriately parametrized coordinates and coefficients. They all have the same form:

$$\epsilon_1 z_1 + \epsilon_2 z_2 + z_3 + \epsilon_4 z_4 + z_5 + \epsilon_6 z_6 = 0 \quad (1.3a)$$

$$\rho_3 z_3 + z_4 + \rho_5 z_5 + \rho_6 z_6 - 1 = 0 \quad (1.3b)$$

$$z_3 z_4 - z_1 z_2 \lambda_1 = 0 \quad (1.3c)$$

$$z_3 z_4 - z_5 z_6 \lambda_2 = 0, \quad (1.3d)$$

with different choices of the parameters ϵ_i and ρ_j according to the chosen Painlevé equation. We call these surfaces \mathcal{Z} -Segre surfaces. They are listed in Table 1.1.

TABLE 1.1. Affine Segre surfaces for qP_{VI} and all differential Painlevé equations.

Painlevé equation	\mathcal{Z} -Segre surface
qP_{VI}	$z_1 + z_2 + z_3 + z_4 + z_5 + z_6 = 0,$ $\rho_2 z_2 + \rho_3 z_3 + z_4 + \rho_5 z_5 + \rho_6 z_6 = 1,$ $z_3 z_4 - \lambda_1 z_1 z_2 = 0, \quad z_5 z_6 - \lambda_2 z_1 z_2 = 0.$
P_{VI}	$z_1 + z_2 + z_3 + z_4 + z_5 + z_6 = 0,$ $\rho_3 z_3 + z_4 + \rho_5 z_5 + \rho_6 z_6 - 1 = 0,$ $z_3 z_4 - \lambda_1 z_1 z_2 = 0, \quad z_5 z_6 - \frac{\rho_3 \lambda_1}{\rho_5 \rho_6} z_1 z_2 = 0.$

P_V	$z_1 + z_2 + z_3 + z_4 + z_5 + z_6 = 0,$ $z_4 + \rho_5 z_5 - 1 = 0,$ $z_3 z_4 - \lambda_1 z_1 z_2 = 0, \quad z_5 z_6 - \lambda_2 z_1 z_2 = 0.$
P_V^{deg}	$z_1 + z_3 + z_4 + z_5 + z_6 = 0,$ $\rho_3 z_3 + z_4 + \rho_5 z_5 + \frac{\rho_3}{\rho_5} z_6 - 1 = 0,$ $z_3 z_4 - z_1 z_2 = 0, \quad z_5 z_6 - z_1 z_2 = 0.$
P_{IV}	$z_1 + z_2 + z_3 + z_4 + z_5 + z_6 = 0,$ $z_4 - 1 = 0,$ $z_3 z_4 - \lambda_1 z_1 z_2 = 0, \quad z_5 z_6 - \lambda_2 z_1 z_2 = 0.$
$P_{III}^{D_6}$	$z_1 + z_2 + z_3 + z_4 + z_5 = 0,$ $z_4 + \rho_5 z_5 - 1 = 0,$ $z_3 z_4 - \lambda_1 z_1 z_2 = 0, \quad z_5 z_6 - z_1 z_2 = 0.$
$P_{III}^{D_7}$	$z_1 + z_2 + z_3 + z_4 + z_5 = 0,$ $z_4 + \rho_5 z_5 - 1 = 0,$ $z_3 z_4 - z_1 z_2 = 0, \quad z_5 z_6 - z_1 z_2 = 0.$
$P_{II}^{\text{JM}}, P_{II}^{\text{FN}}$	$z_1 + z_2 + z_3 + z_4 + z_5 + z_6 = 0,$ $z_4 - 1 = 0,$ $z_3 z_4 - z_1 z_2 = 0, \quad z_5 z_6 - \lambda_2 z_1 z_2 = 0.$
P_I	$z_3 + z_4 + z_5 + z_6 = 0,$ $z_4 - 1 = 0,$ $z_3 z_4 - z_1 z_2 = 0, \quad z_3 z_4 - z_5 z_6 = 0.$

We also show that the resulting Segre surfaces are isomorphic to the corresponding cubic surfaces as affine varieties, for each Painlevé equation. To achieve this, first we calculate the continuum limit \mathcal{Z}_1 of the Segre surface \mathcal{Z}_q and show that it is given by (1.3) with generic parameters defined by equations (3.4) and (3.5). Our first main result is the following theorem.

Theorem 1.1. *The Jimbo-Fricke cubic surface for P_{VI} and the Segre surface \mathcal{Z}_1 are isomorphic as affine varieties, with parameters related by equations (3.4), (3.5), (3.23) and (3.24).*

In Remark 3.12 we show that this theorem answers certain questions asserted in [40]. The proof of Theorem 1.1 relies on two main ingredients. One is the comparison of the asymptotic behaviours of solutions of P_{VI} and qP_{VI} as $q \rightarrow 1$. The other is based on blow-downs of lines at infinity on the cubic surface.

Using an explicit form of this isomorphism (see Theorem 3.8), we show that, in all un-ramified cases, the confluence scheme of the Painlevé monodromy manifolds can be carried through to the affine transformation, therefore producing isomorphisms between each cubic surface and the \mathcal{Z} -Segre obtained by confluencing \mathcal{Z}_1 .

For the ramified cases, the confluence either produces a reducible Segre surface or a family which does not have the correct number of free parameters. This is not surprising because the confluences to ramified and to non-ramified Painlevé equations are deeply different in geometric as well as analytic terms.

For this reason, we perform an in-depth study of the lines and singularities of all cubic surfaces with a triangle of lines at infinity, as well as their blow downs, which allows us to build the isomorphic \mathcal{Z} -Segre for ramified cases as well.

Theorem 1.2. *The monodromy manifold of each differential Painlevé equation (except $P_{III}^{D_8}$) is isomorphic to the corresponding \mathcal{Z} -Segre surface reported in Table 1.1 as affine varieties.*

Remark 1.3. It is not unusual for $P_{\text{III}}^{D_8}$ to be an exceptional case here. It is often the exception in many studies of the Painlevé equations in the literature. For example, in [2], Lax pairs were proposed for all the generalised multi-particle Painlevé equations, and $P_{\text{III}}^{D_8}$ is the only one for which the isomonodromic Hamiltonian is rational rather than polynomial.

The study of lines not only helps in building the above mentioned isomorphism, but it has its own interest and a long history in algebraic geometry. The famous Cayley-Salmon theorem [6, 44] shows that there are 27 lines on a general smooth cubic surface in \mathbb{CP}^3 . For the highly transcendental solutions of the Painlevé equations, these lines give vital information about certain distinguished behaviours. For example, for the first Painlevé equation (P_{I}), lines on the cubic surface correspond to the tronquée solutions [28, Theorem 3] and in the case of P_{VI} , they correspond to truncations of the generic asymptotic expansions, see [29, Prop. 4.5] and [20, Table 1]. On the other hand, geometric properties of the Jimbo-Fricke surfaces and its confluence limits make them important objects in many areas of mathematics, including Cherednik algebras [36, 32], mirror symmetry [18], Calabi-Yau algebras [10], and moduli spaces [49].

Another perspective arises in Okamoto’s characterization of the initial-value space and symmetry group of each Painlevé equation [39, 35] in terms of divisors, intersection theory and the corresponding interpretation of such divisors as generators of affine Weyl groups. In this context, the corresponding limits of the Painlevé equations are realized as coalescences of base points.

The lattices associated with extended affine Weyl or Coxeter groups [22, 11, 19] lead naturally to the construction of discrete Painlevé equations, with each step of iteration being a translation on the lattice, mapping coordinates in one tile (or polytope) in the periodic lattice to another such tile. Multiplicative or q -difference Painlevé equations arise from such operations on a hyperbolic lattice.

In the case of the differential Painlevé equations, the monodromy manifold is “dual” to the initial value space under the Riemann–Hilbert correspondence. For q -discrete Painlevé equations, a notion of monodromy manifold was missing until recently [26, 27]. By constructing a Segre surface for each differential Painlevé equation, we provide the start of a uniform setting that naturally extends to a description of the monodromy manifold of their q -difference analogues.

The coordinate ring of a Painlevé monodromy manifold is endowed with a natural Poisson bracket [33]. By a suitable deformation quantization of this Poisson algebra, the spherical sub-algebras of certain confluent versions of the Cherednik algebra of type \check{C}_1C_1 were obtained in [32]. This led to a representation–theoretic approach to the theory of the Painlevé equations and to surprising links with the theory of basic hypergeometric polynomials. In this paper, we show that the coordinate rings of the Segre surfaces arising as blow downs of such monodromy manifolds also carry a natural Poisson structure [46, 37]. Looking beyond the current paper, it would be interesting to quantise these Segre surfaces and understand their role in the theory of basic hypergeometric polynomials as well as the relation between such quantizations and the monodromy manifolds of the corresponding multiplicative discrete Painlevé equations.

1.1. Terminology. In this paper, we always work over the field \mathbb{C} , and denote the n -dimensional complex projective space, $n \geq 1$, by $\mathbb{P}^n = \mathbb{CP}^n$. We also use the notation $\mathbb{C}^* := \mathbb{C} \setminus \{0\}$. Almost all algebraic varieties in this paper are affine varieties embedded in \mathbb{C}^n , or projective varieties embedded in \mathbb{P}^n , for a suitable positive integer n .

We use small roman letters for affine coordinates, e.g.

$$\mathbb{C}^n = \{(x_1, x_2, \dots, x_n) \in \mathbb{C}^n\},$$

and capitalised roman letters for homogeneous coordinates, e.g.

$$\mathbb{P}^n = \{[X_0 : X_1 : X_2 : \dots : X_n] \in \mathbb{P}^n\}.$$

We consider \mathbb{C}^n as a subset of \mathbb{P}^n in the natural way, through the identification

$$[X_0 : X_1 : X_2 : \dots : X_n] = [1 : x_1 : x_2 : \dots : x_n]. \quad (1.4)$$

Given an embedded affine variety $V \subseteq \mathbb{C}^n$, we define its canonical projective completion $\bar{V} \subseteq \mathbb{P}^n$ as the completion of the image of V in \mathbb{P}^n through (1.4).

Given two embedded affine varieties $V \subseteq \mathbb{C}^m$ and $W \subseteq \mathbb{C}^n$, with $m \leq n$, we say that V and W are *affinely equivalent* if there exists an affine linear mapping $\phi : \mathbb{C}^m \rightarrow \mathbb{C}^n$ of maximal rank m that restricts to an isomorphism $V \rightarrow W$. In this case, we also call W and V affinely equivalent. In particular, given an affine variety $V \subseteq \mathbb{C}^m \subseteq \mathbb{C}^{m+n}$, with $m, n \geq 1$, where \mathbb{C}^m lies linearly in \mathbb{C}^{m+n} , the two views of V as an embedded variety in \mathbb{C}^m or \mathbb{C}^{m+n} are affinely equivalent. Note that affine equivalence is generally stronger than isomorphic as affine varieties.

Projective equivalence between embedded projective varieties is defined similarly.

1.2. Acknowledgements. This work was funded by the Leverhulme Trust visiting professorship grant VP2-2018-013. Part of the work was carried out during NJ's and PR's residence at the Isaac Newton Institute for Mathematical Sciences in the program ‘‘Applicable resurgent asymptotics: towards a universal theory’’ during September-December 2022. We are grateful for the hospitality of the organizers of this program and the support of the Newton Institute. MM's research was supported by the Leverhulme Trust Research Project Grant RPG-2021-047. PR and NJ's research was supported by the Australian Research Council Discovery Project #DP210100129. The authors thank Davide Guzzetti, Oleg Lisovyy, Volodya Rubtsov for fruitful discussions.

2. THE AFFINE SEGRE SURFACE OF THE qP_{VI} EQUATION

For simplicity, we refer to the Segre surface (1.1) as

$$\mathcal{Z}_q := \{z \in \mathbb{C}^6 : h_i = 0 \text{ for } 1 \leq i \leq 4\}, \quad (2.1)$$

where for given $\mu_1, \dots, \mu_6, \lambda_1, \lambda_2 \in \mathbb{C}$, the polynomials $h_i \in \mathbb{C}[z_1, \dots, z_6]$, $1 \leq i \leq 4$, are defined by

$$h_1 = z_1 + z_2 + z_3 + z_4 + z_5 + z_6, \quad (2.2a)$$

$$h_2 = \mu_1 z_1 + \mu_2 z_2 + \mu_3 z_3 + \mu_4 z_4 + \mu_5 z_5 + \mu_6 z_6 - 1, \quad (2.2b)$$

$$h_3 = z_3 z_4 - z_1 z_2 \lambda_1, \quad (2.2c)$$

$$h_4 = z_5 z_6 - z_1 z_2 \lambda_2, \quad (2.2d)$$

where μ_k , $1 \leq k \leq 6$ and λ_1, λ_2 , are some complex numbers.

We show that a generic member of the family \mathcal{Z}_q is an affine Segre surface, with smooth canonical projective completion, and conversely, that a generic embedded affine Segre surface with smooth canonical projective completion, can be put into the form (2.1) via an affine transformation.

The family of affine surfaces \mathcal{Z}_q first appeared in [26] as the monodromy manifold of the linear problem for qP_{VI} . In that case, the coefficients μ_1, \dots, μ_6 and λ_1, λ_2 are parametrised explicitly in terms of the six parameters of the qP_{VI} equation. The fact that only six of the eight coefficients the coefficients μ_1, \dots, μ_6 and λ_1, λ_2 are independent is a consequence of the intrinsic scaling freedoms in the defining equations (2.2).

Lemma 2.1. *The family of affine surfaces \mathcal{Z}_q is equivalent to the six parameter family defined as the zero set in \mathbb{C}^6 of h_1, h_3, h_4 and h_2'' where*

$$h_2'' = \rho_2 z_2 + \rho_3 z_3 + z_4 + \rho_5 z_5 + \rho_6 z_6 - 1. \quad (2.3)$$

Proof. Using the freedom of adding arbitrary multiples of h_1 to h_2 , we may normalise h_2 so that it has no z_1 term,

$$h_2' = (\mu_2 - \mu_1)z_2 + (\mu_3 - \mu_1)z_3 + (\mu_4 - \mu_1)z_4 + (\mu_5 - \mu_1)z_5 + (\mu_6 - \mu_1)z_6 - 1.$$

Moreover, using the freedom of scaling $z_k \mapsto c z_k$, $1 \leq k \leq 6$, for a nonzero c , which leaves h_1, h_3 and h_4 invariant, but scales each coefficient in h_2 by c^{-1} , except the constant term. Choose two coefficients $\mu_i \neq \mu_j$, then one can apply this scaling with the choice $c = \mu_i - \mu_j$. Without loss of generality, assume $\mu_4 \neq \mu_1$ (if this is not the case, relabel accordingly), then we can normalise h_2' such that the coefficient of z_4 equals one, resulting in (2.3), where

$$\rho_k := \frac{\mu_k - \mu_1}{\mu_4 - \mu_1}, \quad k = 2, 3, 5, 6. \quad (2.4)$$

□

Lemma 2.1 shows that the collection of algebraic surfaces (2.1), up to affine equivalence, constitutes at most a six-dimensional family. In Section 2.1, we prove that it is in fact exactly six-dimensional. For this reason, and the fact that this family appears as the monodromy manifold of qPVI, we give the following definition:

Definition 2.2. We refer to the embedded affine surface \mathcal{Z}_q defined in (2.1), or equivalently the one defined as the zero set in \mathbb{C}^6 of h_1, h_3, h_4 and h_2'' , as the *affine Segre surface of qPVI*.

Lemma 2.1 also provides a reduced form in which redundant parameters have been eliminated. This is important in order to speak about *generic parameters*.

Definition 2.3. We say that the parameters μ_1, \dots, μ_6 and λ_1, λ_2 are generic if $\mu_4 \neq \mu_1$ and $\lambda_1, \lambda_2, \rho_2, \rho_3, \rho_5, \rho_6$, defined in (2.4), do not satisfy any non-trivial polynomial relations with rational coefficients.

This Section is organised as follows: in Section 2.1 we show that generic members of the family \mathcal{Z}_q are affine Segre surfaces and we study their projective completion. In Section 2.2, we show that the family \mathcal{Z}_q is a six parameter family. In particular, we derive a standard form for embedded affine Segre surfaces, with smooth canonical projective completion, and prove that a generic such surface can be transformed into (2.1) via an affine transformation. This is followed by Section 2.3, where we study the parametrisation of the surface in terms of the parameters of qPVI, and we further provide explicit formulas of the 16 lines on the Segre surface in Section 2.4.

2.1. Algebraic characterisation. Notice that (2.1) indeed defines an affine Segre surface as, upon eliminating two of the variables using the linear equations $\{h_1 = 0, h_2 = 0\}$, one is left with two quadratic equations $\{h_3 = 0, h_4 = 0\}$, which define an irreducible variety given by the complete intersection of two quadrics in \mathbb{C}^4 , for generic coefficients.

In order to give an algebraic characterisation of its generic members, we define the canonical projective completion $\overline{\mathcal{Z}}_q$ in \mathbb{P}^6 of \mathcal{Z}_q , replacing h_2 with h_2'' defined in (2.3) and using projective coordinates,

$$[Z_0 : Z_1 : Z_2 : Z_3 : Z_4 : Z_5 : Z_6] = [1 : z_1 : z_2 : z_3 : z_4 : z_5 : z_6],$$

by

$$Z_1 + Z_2 + Z_3 + Z_4 + Z_5 + Z_6 = 0, \quad (2.5a)$$

$$\rho_2 Z_2 + \rho_3 Z_3 + Z_4 + \rho_5 Z_5 + \rho_6 Z_6 - Z_0 = 0, \quad (2.5b)$$

$$Z_3 Z_4 - Z_1 Z_2 \lambda_1 = 0, \quad (2.5c)$$

$$Z_5 Z_6 - Z_1 Z_2 \lambda_2 = 0. \quad (2.5d)$$

Proposition 2.4. *For generic coefficients, the Segre surface $\overline{\mathcal{Z}}_q$ is smooth and the hyperplane section at infinity, $\overline{\mathcal{Z}}_q \setminus \mathcal{Z}_q$, is an irreducible smooth quartic curve, isomorphic to the intersection of two quadric surfaces in \mathbb{P}^3 , of genus 1.*

Proof. For $\overline{\mathcal{Z}}_q$ to have a singularity at some point $Z \in \mathbb{P}^6$, it is required that the the Jacobian, of the left-hand sides of equations (2.5),

$$J = \begin{bmatrix} 0 & 1 & 1 & 1 & 1 & 1 & 1 \\ -1 & 0 & \rho_2 & \rho_3 & 1 & \rho_5 & \rho_6 \\ 0 & -\lambda_1 Z_2 & -\lambda_1 Z_1 & Z_4 & Z_3 & 0 & 0 \\ 0 & -\lambda_2 Z_2 & -\lambda_2 Z_1 & 0 & 0 & Z_6 & Z_5 \end{bmatrix},$$

has rank less than four at this point. By subtracting Z_3 times the first row from the third row, it follows that this is equivalent to the following two row vectors being linearly dependent,

$$\begin{aligned} v_1 &= [-\lambda_1 Z_2 - Z_3 & -\lambda_1 Z_1 - Z_3 & Z_4 - Z_3 & -Z_3 & -Z_3], \\ v_2 &= [-\lambda_2 Z_2 & -\lambda_2 Z_1 & 0 & Z_6 & Z_5]. \end{aligned}$$

Following the proof of [42, Proposition 2.6], we find that equations (2.5) and

$$r_0 v_1 - r_1 v_2 = 0,$$

have a common solution in $Z \in \mathbb{P}^6$ and $[r_0 : r_1] \in \mathbb{P}^1$, only if $\lambda_1 = 0$, $\lambda_2 = 0$ or

$$(\lambda_1 - \lambda_2)^2 - 2(\lambda_1 + \lambda_2) + 1 = 0. \quad (2.6)$$

As a consequence, $\overline{\mathcal{Z}}_q$ is smooth for generic coefficients.

The curve at infinity is obtained by setting $Z_0 = 0$ in equations (2.5). Upon eliminating Z_3 and Z_6 using the first two resulting linear equations, we are left with two quadratic equations,

$$\begin{aligned} Q_1 &:= \lambda_1 Z_1 Z_2 - Z_4 \frac{-\rho_6 Z_1 + (\rho_2 - \rho_6) Z_2 + (1 - \rho_6) Z_4 + (\rho_5 - \rho_6) Z_5}{(\rho_6 - \rho_3)} = 0, \\ Q_2 &:= \lambda_2 Z_1 Z_2 + Z_5 \frac{-\rho_3 Z_1 + (\rho_2 - \rho_3) Z_2 + (1 - \rho_3) Z_4 + (\rho_5 - \rho_3) Z_5}{(\rho_6 - \rho_3)} = 0. \end{aligned}$$

This shows that the curve at infinity is isomorphic to the intersection of the two quadrics in $\{[Z_1 : Z_2 : Z_4 : Z_5] \in \mathbb{P}^3\}$, defined by the above two equations.

If the curve has a singularity at some point, then the gradients of Q_1 and Q_2 , with respect to Z_1, Z_2, Z_4, Z_5 , must be multiples of each other at this point, that is,

$$r_0 \nabla Q_1 - r_1 \nabla Q_2 = 0, \quad (2.7)$$

for some $[r_0 : r_1] \in \mathbb{P}^1$. Equation (2.7) constitutes four multi-linear homogeneous equations among the variables $r_0, r_1, Z_1, Z_2, Z_4, Z_5$. A direct calculation shows that these four equations, in addition to the two quadratic equations $Q_{1,2} = 0$, admit a common solution only if certain algebraic relations among the parameters $\lambda_1, \lambda_2, \rho_2, \rho_3, \rho_5, \rho_6$ are satisfied. As a consequence, the curve at infinity is smooth for generic coefficients.

Finally, we recall that a smooth intersection of two quadrics in \mathbb{P}^3 with a rational point, is isomorphic to an elliptic curve [8, §8 (iv)]. The curve at infinity is therefore irreducible and has genus 1. The proposition follows. \square

Remark 2.5. Recall that a smooth intersection of two quadrics in \mathbb{P}^3 has genus 1, so the hyperplane section at infinity, $\overline{\mathcal{Z}}_q \setminus \mathcal{Z}_q$, in Proposition 2.4, with rational point $[1 : 0 : 0 : 0]$, is generically an elliptic curve. To put it into cubic form, note that both Q_1 and Q_2 are linear in Z_1 , so their resultant with respect to Z_1 is a homogeneous cubic in Z_2, Z_4, Z_5 , defining the following cubic curve in \mathbb{P}^2 ,

$$Z_2 Z_5 \lambda_1 (Z_2 (\rho_3 - \rho_2) + Z_4 (\rho_3 - 1) + Z_5 (\rho_3 - \rho_5)) - Z_4 Z_5 (\rho_2 Z_2 + Z_4 + \rho_5 Z_5) + Z_2 Z_4 \lambda_2 (Z_2 (\rho_6 - \rho_2) + Z_4 (\rho_6 - 1) + Z_5 (\rho_6 - \rho_5)) = 0.$$

The projection of the curve at infinity onto this cubic, regularised near the point $[1 : 0 : 0 : 0]$ by sending

$$[1 : 0 : 0 : 0] \mapsto [1 : \lambda_1 (\rho_3 / \rho_6 - 1) : \lambda_2 (\rho_6 / \rho_3 - 1)],$$

is an isomorphism [8, §8 (iv)].

2.2. Smooth Segre surfaces up to affine equivalence. In this section, we derive a standard form for embedded affine Segre surfaces with smooth canonical projective completion. Because as observed at the beginning of Subsection 2.1, each generic element of \mathcal{Z}_q defines an irreducible variety given by the complete intersection of two quadrics in \mathbb{C}^4 , in this subsection we work in \mathbb{P}^4 .

Firstly, let us recall that any isomorphism between Segre surfaces, embedded in \mathbb{P}^4 , comes from a projective equivalence. We formulate this as the following lemma.

Lemma 2.6. *Any isomorphism $\phi : \mathcal{S} \rightarrow \mathcal{S}'$ between embedded smooth Segre surfaces $\mathcal{S}, \mathcal{S}' \subseteq \mathbb{P}^4$, extends to a unique linear projective transformation of \mathbb{P}^4 .*

Proof. Any smooth Segre surface $\mathcal{S} \subseteq \mathbb{P}^4$ is the image under the anti-canonical mapping of the projective plane blown up at five points in general position [13]. Explicitly, choose five skew lines L_k , $1 \leq k \leq 5$, in \mathcal{S} . Then there exist five points $u_k \in \mathbb{P}^2$, $1 \leq k \leq 5$, in general position, and a bi-rational map

$$\pi : \mathcal{S} \rightarrow \mathbb{P}^2,$$

which is the simultaneous blow down of L_k to u_k , $1 \leq k \leq 5$.

For an explicit description of π^{-1} , consider the vector space

$$\mathcal{C} = \{\text{cubic forms in } \{U_0, U_1, U_2\} \text{ that vanish at } u_k, 1 \leq k \leq 5\},$$

where we use projective coordinates $[U_0 : U_1 : U_2]$ on \mathbb{P}^2 . This vector space has dimension 5 and there exists a basis $\{C_0, \dots, C_4\}$ of \mathcal{C} such that π^{-1} is given by

$$\begin{array}{ccc} \pi^{-1} : & \mathbb{P}^2 & \dashrightarrow \mathcal{S} \\ & [U_0 : U_1 : U_2] & \mapsto [C_0 : C_1 : C_2 : C_3 : C_4]. \end{array}$$

Let $\mathcal{S}' \subseteq \mathbb{P}^4$ be another smooth Segre surface and denote correspondingly the objects introduced above by π' , u'_k , $1 \leq k \leq 5$ and \mathcal{C}' for this Segre surface. Then, an isomorphism $\phi : \mathcal{S} \rightarrow \mathcal{S}'$, induces a corresponding automorphism $\tilde{\phi}$ of \mathbb{P}^2 , making the diagram

$$\begin{array}{ccc} \mathcal{S} & \xrightarrow{\phi} & \mathcal{S}' \\ \pi \downarrow & & \downarrow \pi' \\ \mathbb{P}^2 & \xrightarrow{\tilde{\phi}} & \mathbb{P}^2 \end{array}$$

commutative. So $\tilde{\phi}$ is a projective linear transformation. Upon picking an $M \in GL_3(\mathbb{C})$ such that $\tilde{\phi}(U) = U \cdot M$, $U \in \mathbb{P}^2$, we obtain a corresponding isomorphism

$$\mathcal{L} : \mathcal{C}' \rightarrow \mathcal{C}, \mathcal{C}' \mapsto \mathcal{C}, \quad C(U) = C'(U \cdot M).$$

Denote by $N \in GL_5(\mathbb{C})$ the unique change of basis matrix

$$(\mathcal{L}C'_0, \mathcal{L}C'_1, \mathcal{L}C'_2, \mathcal{L}C'_3, \mathcal{L}C'_4) = (C_0, C_1, C_2, C_3, C_4) \cdot N, \quad (2.8)$$

then $\phi(S) = S \cdot N$ for $S \in \mathcal{S}$ and the lemma follows. \square

Remark 2.7. From the blow-up model of smooth Segre surfaces in the proof of Lemma 2.6, it follows that a Segre surface has exactly 16 lines: 5 are the exceptional lines above the base points, 10 are the total transforms of lines in \mathbb{P}^2 going through two of the base points and finally one further line is the total transform of the unique conic going through all five base points. In Section 2.4, we give explicit expressions for the 16 lines on the affine Segre surface of qP_{VI} , as well as their intersection graph.

It follows from the lemma above that, to classify smooth Segre surfaces up to isomorphism, it suffices to classify them up to projective equivalence. In turn, see [34] or [13, Theorem 8.5.1], any smooth Segre surface \mathcal{S} can, by a linear projective transformation, be put into diagonal form

$$S_0^2 + S_1^2 + S_2^2 + S_3^2 + S_4^2 = 0, \quad \gamma_0 S_0^2 + \gamma_1 S_1^2 + \gamma_2 S_2^2 + \gamma_3 S_3^2 + \gamma_4 S_4^2 = 0,$$

for some mutually distinct $\gamma_k \in \mathbb{C}$, $0 \leq k \leq 4$. Since we are interested in embedded affine Segre surfaces (with smooth canonical projective completion), we derive a corresponding standard form for them, up to affine transformations.

Given any embedded affine Segre surface $\mathcal{S} \subseteq \mathbb{C}^4$, we may choose symmetric 5×5 matrices M, N such that \mathcal{S} is described by

$$s^T M s = 0, \quad s^T N s = 0, \quad s = [1, s_1, s_2, s_3, s_4]^T. \quad (2.9)$$

The canonical projective completion $\overline{\mathcal{S}} \subseteq \mathbb{P}^4$ of \mathcal{S} is obtained through projective coordinates

$$[S_0 : S_1 : S_2 : S_3 : S_4] = [1 : s_1 : s_2 : s_3 : s_4],$$

by replacing $s \mapsto S$ in equations (2.9). The Segre surface $\overline{\mathcal{S}}$ is smooth if and only if the equation

$$|r_0 M - r_1 N| = 0, \quad (2.10)$$

has five distinct solutions $r = [r_0 : r_1] \in \mathbb{P}^1$, see [13, §8.5].

We have the freedom of applying automorphisms of \mathbb{C}^4 to equations (2.9), i.e. affine transformations

$$s \mapsto s' = G^{-1}s, \quad M \mapsto M' = G^T M G, \quad N \mapsto N' = G^T N G, \quad (2.11)$$

where $G \in GL_5(\mathbb{C})$ takes the form

$$G = \begin{bmatrix} 1 & \underline{0}^T \\ b & G_4 \end{bmatrix},$$

for some $G_4 \in GL_4(\mathbb{C})$ and vector $b \in \mathbb{C}^4$, where $\underline{0}^T$ denotes the transposed zero vector. Using such transformations, we arrive at a standard form for embedded affine Segre surfaces with smooth canonical projective completion, as detailed in the following lemma.

Lemma 2.8. *Given an embedded affine Segre surface $\mathcal{S} \subseteq \mathbb{C}^4$, whose canonical projective completion $\overline{\mathcal{S}} \subseteq \mathbb{P}^4$ is smooth, there exists an affine transformation that puts it in the standard form*

$$s_1^2 + s_2^2 + s_3^2 + s_4^2 = 0, \quad (2.12a)$$

$$\alpha_1 s_1^2 + \alpha_2 s_2^2 + \alpha_3 s_3^2 + \alpha_4 s_4^2 + \beta_1 s_1 + \beta_2 s_2 + \beta_3 s_3 + \beta_4 s_4 + 1 = 0, \quad (2.12b)$$

for some $\alpha_k, \beta_k \in \mathbb{C}$, $1 \leq k \leq 4$.

Remark 2.9. In the standard form (2.12), there is still the freedom of adding arbitrary multiples of the first equation to the second, so that we may for example normalise the α 's such that $\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 = 0$. Similarly, there is the freedom of scaling $s_k \mapsto g^{-1}s_k$, $1 \leq k \leq 4$, so that

$$\alpha_k \mapsto g^2\alpha_k, \quad \beta_k \mapsto g\beta_k, \quad (1 \leq k \leq 4),$$

In the generic setting, when the α 's are distinct, the only other freedom comes from permuting the variables $\{s_1, s_2, s_3, s_4\}$. In particular, the family of embedded affine Segre surfaces, with smooth canonical projective completion, up to affine equivalence, is six-dimensional.

Remark 2.10. We note that the canonical projective completion of the Segre surface, defined by equations (2.12), is only smooth for generic coefficients: it is smooth if and only if the discriminant $\text{Disc}_r(|N - rM|)$ is nonzero, where M and N are the symmetric matrices corresponding to equations (2.12), as in equation (2.9).

Proof of Lemma 2.8. Choose symmetric 5×5 matrices M, N such that S is described by equations (2.9). Our goal is to bring these equations into the standard form (2.12), using affine transformations (2.11), as well as by taking linear combinations of the two equations,

$$M \mapsto a_{11}M + a_{12}N, \quad N \mapsto a_{21}M + a_{22}N, \quad (a_{ij})_{1 \leq i, j \leq 2} \in GL_2(\mathbb{C}).$$

We are going to give an algorithmic method to accomplish this. We will be indexing 5×5 matrices from 0 to 4, e.g. $P = (p_{ij})_{0 \leq i, j \leq 4}$, and we further introduce the notation $P_4 = (p_{ij})_{1 \leq i, j \leq 4}$.

Let us for the moment assume that

$$|M_4 - rN_4| \neq 0. \tag{2.13}$$

This assumption is in fact moot, as it is always satisfied when \bar{S} is smooth, but we will provide an argument for this a bit later.

Due to (2.13), we know that

$$|r_0M_4 - r_1N_4| = 0, \tag{2.14}$$

has at most four solutions $r = [r_0 : r_1] \in \mathbb{P}^1$. On the other hand, since \bar{S} is smooth, equation (2.10) has five distinct solutions $r = [r_0 : r_1] \in \mathbb{P}^1$. Pick a solution $r \in \mathbb{P}^1$ to (2.10), which does not satisfy (2.14). If $r_0 \neq 0$, replace M by $r_0M_4 - r_1N_4$, otherwise simply swap M and N . Then M is not invertible, but M_4 is. Since M_4 is symmetric and invertible, we can construct an orthogonal matrix P such that $P^T M_4 P = D$, where D is a diagonal matrix with nonzero entries. By applying the affine transformation (2.11), with

$$G = \begin{bmatrix} 1 & \mathbf{0}^T \\ \mathbf{0} & PD^{-\frac{1}{2}} \end{bmatrix},$$

we normalise M such that $M_4 = I_4$, that is,

$$M = \begin{bmatrix} m_0 & m_1 & m_2 & m_3 & m_4 \\ m_1 & 1 & 0 & 0 & 0 \\ m_2 & 0 & 1 & 0 & 0 \\ m_3 & 0 & 0 & 1 & 0 \\ m_4 & 0 & 0 & 0 & 1 \end{bmatrix},$$

with $m_0 = m_1^2 + m_2^2 + m_3^2 + m_4^2$, since $|M| = 0$. Next, we apply the affine transformation (2.11), with

$$G = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ -m_1 & 1 & 0 & 0 & 0 \\ -m_2 & 0 & 1 & 0 & 0 \\ -m_3 & 0 & 0 & 1 & 0 \\ -m_4 & 0 & 0 & 0 & 1 \end{bmatrix},$$

which reduces the matrix M to

$$M = \begin{bmatrix} 0 & \underline{0}^T \\ \underline{0} & I_4 \end{bmatrix}. \quad (2.15)$$

So, the first equation in (2.9) is now given by (2.12a).

Next, we diagonalise N_4 . Let P be an orthonormal matrix such that $P^T N_4 P = D$, where D is a diagonal matrix. Applying the affine transformation (2.11), with

$$G = \begin{bmatrix} 1 & \underline{0}^T \\ \underline{0} & P \end{bmatrix}, \quad (2.16)$$

the matrix M is left invariant, whereas N takes the form

$$N = \begin{bmatrix} \beta_0 & \frac{1}{2}\beta_1 & \frac{1}{2}\beta_2 & \frac{1}{2}\beta_3 & \frac{1}{2}\beta_4 \\ \frac{1}{2}\beta_1 & \alpha_1 & 0 & 0 & 0 \\ \frac{1}{2}\beta_2 & 0 & \alpha_2 & 0 & 0 \\ \frac{1}{2}\beta_3 & 0 & 0 & \alpha_3 & 0 \\ \frac{1}{2}\beta_4 & 0 & 0 & 0 & \alpha_4 \end{bmatrix}, \quad (2.17)$$

for some $\alpha_k, \beta_k \in \mathbb{C}$, $1 \leq k \leq 4$ and $\beta_0 \in \mathbb{C}$. In fact, β_0 must be nonzero since $|M - rN|$ cannot have a double root at $r = 0$. By dividing N by β_0 , equations (2.9) have been brought into the standard form (2.12).

To finish the proof, we come back to the assumption (2.13), that we made in the beginning. Suppose that it does not hold true. Since $|M - rN| \not\equiv 0$, we may replace M by $M - rN$, for a generic r , so that M is invertible. In particular, for such a choice of M the rank of M_4 must be at least 3. On the other hand, as (2.13) does not hold true, $|M_4| = 0$ and so M_4 must have rank exactly 3. Analogous to how we brought M and N into the forms (2.15) and (2.17) above, and by additionally permuting the variables $\{s_1, s_2, s_3, s_4\}$ if necessary, we bring M and N into the form

$$M = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}, \quad N = \begin{bmatrix} \beta_0 & \frac{1}{2}\beta_1 & \frac{1}{2}\beta_2 & \frac{1}{2}\beta_3 & \frac{1}{2}\beta_4 \\ \frac{1}{2}\beta_1 & \alpha_1 & \gamma_2 & \gamma_3 & \gamma_4 \\ \frac{1}{2}\beta_2 & \gamma_2 & \alpha_2 & 0 & 0 \\ \frac{1}{2}\beta_3 & \gamma_3 & 0 & \alpha_3 & 0 \\ \frac{1}{2}\beta_4 & \gamma_4 & 0 & 0 & \alpha_4 \end{bmatrix}.$$

From the coefficient of r^1 in $|M_4 - rN_4| \equiv 0$, we immediately read off that $\alpha_1 = 0$. By similarly looking at the coefficients of r^2, r^3 and r^4 , we obtain the linear system

$$\begin{bmatrix} 1 & 1 & 1 \\ \alpha_3 + \alpha_4 & \alpha_2 + \alpha_4 & \alpha_2 + \alpha_3 \\ \alpha_3\alpha_4 & \alpha_2\alpha_4 & \alpha_2\alpha_3 \end{bmatrix} \cdot \begin{bmatrix} \gamma_2^2 \\ \gamma_3^2 \\ \gamma_4^2 \end{bmatrix} = 0. \quad (2.18)$$

If $\gamma_2 = \gamma_3 = \gamma_4 = 0$, then $|M - rN|$ has a double root at $r = 0$, so \bar{S} is not smooth and we have arrived at a contradiction. Else α_2, α_3 and α_4 are not all distinct, and by solving (2.18) for $\{\gamma_2, \gamma_3, \gamma_4\}$, one similarly sees that $|M - rN|$ always has a root with multiplicity, again yielding a contradiction. We conclude that assumption (2.13) always holds and the lemma follows. \square

In the following proposition, it is shown that a generic embedded affine Segre surface, with smooth projective completion, can be put in the form (2.1), by an affine transformation.

Proposition 2.11. *For generic $\alpha_k, \beta_k \in \mathbb{C}$, $1 \leq k \leq 4$, the Segre surface (2.12) is affinely equivalent to the Segre surface (2.1), for some values of the coefficients in (2.1).*

Proof. The proof of Lemma 2.8 gives an algorithmic procedure to put an embedded affine Segre surface, with smooth projective completion, into the form (2.12). We are going to apply this procedure to the family of Segre surfaces \mathcal{Z}_q (2.1). This will provide a set of algebraic equations among the parameters of both families of Segre surfaces. By application of the implicit function theorem and standard algebraic arguments, we obtain that this set of algebraic equations is solvable for generic parameter values $\alpha_k, \beta_k \in \mathbb{C}$, $1 \leq k \leq 4$, from which the proposition will follow.

As a first step, we eliminate two of the variables in (2.1). Assuming $\rho_2 \neq 0$, we use the linear equations $\{h_1 = 0, h_2'' = 0\}$ to eliminate z_1 and z_2 from the equations $\{h_3 = 0, h_4 = 0\}$, and, by renaming the remaining variables as $z_{k+2} = s_k$, $1 \leq k \leq 4$, we arrive at

$$\begin{aligned} s_1 s_2 - \frac{\lambda_1}{\lambda_2} s_3 s_4 &= 0, \\ s_1 s_2 - \frac{\lambda_1}{\rho_2^2} (1 + (\rho_2 - \rho_3) s_1 + (\rho_2 - 1) s_2 + (\rho_2 - \rho_5) s_3 + (\rho_2 - \rho_6) s_4) \\ &\quad (\rho_3 s_1 + s_2 + \rho_5 s_3 + \rho_6 s_4 - 1) = 0, \end{aligned}$$

where we used the ρ_k , $1 \leq k \leq 4$, defined in (2.4).

Now, construct the corresponding symmetric 5 5 matrices M, N , as in (2.9). Applying the affine transformation (2.11), with

$$G = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & -i & i & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & -\sqrt{\frac{\lambda_2}{\lambda_1}} & \sqrt{\frac{\lambda_2}{\lambda_1}} \\ 0 & 0 & 0 & i\sqrt{\frac{\lambda_2}{\lambda_1}} & i\sqrt{\frac{\lambda_2}{\lambda_1}} \end{bmatrix},$$

readily puts M into the form (2.15).

What is left, is to construct an orthonormal matrix P that diagonalizes N_4 , i.e. $P^T N_4 P = D$, where D is a diagonal matrix. Upon introducing such a P formally, applying (2.16), comparing the result with (2.17) (with $\beta_0 = 1$) and finally eliminating the entries of P by taking resultants, one is left with 7 polynomial equations, among the variables $\lambda_1, \lambda_2, \rho_2, \rho_3, \rho_5, \rho_6$ and α_k, β_k , $1 \leq k \leq 4$. By remark (2.9), we may normalise $\alpha_4 = -\alpha_1 - \alpha_2 - \alpha_3$, and by further introducing the scaling by nonzero g as specified in the remark, we end up with seven polynomial equations, with rational coefficients, among two sets of seven variables,

$$F_k = F_k(g, \lambda, \rho; \alpha, \beta) = 0 \quad (1 \leq k \leq 7),$$

where $\lambda = (\lambda_1, \lambda_2)$, $\rho = (\rho_2, \rho_3, \rho_5, \rho_6)$, $\alpha = (\alpha_k)_{1 \leq k \leq 3}$ and $\beta = (\beta_k)_{1 \leq k \leq 4}$.

Explicit formulas for these polynomials are lengthy and best obtained using mathematical software, e.g. Mathematica. One of the simplest solutions that we could find to this polynomial system, is given by

$$\begin{aligned} \lambda_1 &= -3, & \lambda_2 &= -3, & \rho_2 &= 2, \\ \rho_3 &= \frac{2}{3}, & \rho_5 &= -1, & \rho_6 &= \frac{2}{3}, \end{aligned}$$

$$\alpha_1 = \frac{1}{6}(2 + \sqrt{6})i, \quad \alpha_2 = \frac{1}{6}(-2 + \sqrt{6})i, \quad \alpha_3 = \frac{1}{6}(2 - \sqrt{6})i,$$

together with $g = 1$ and

$$\begin{aligned} \beta_1 &= \frac{1}{6}\sqrt{(-24 + 39i) - (16 - 15i)\sqrt{6}}, & \beta_2 &= \frac{1}{6}\sqrt{(-24 - 39i) + (16 + 15i)\sqrt{6}}, \\ \beta_3 &= \frac{1}{6}\sqrt{(-24 + 39i) + (16 - 15i)\sqrt{6}}, & \beta_4 &= \frac{1}{6}\sqrt{(-24 - 39i) - (16 + 15i)\sqrt{6}}. \end{aligned}$$

In particular, the system $F = (F_k)_{1 \leq k \leq 7}$ is consistent and defines a non-empty algebraic set $V \subseteq \mathbb{C}^{14}$. Furthermore, the Jacobian determinant $|J_{\lambda, \rho, g}(F)|$ of $F = (F_k)_{1 \leq k \leq 7}$ with respect to the seven variables $\{g, \lambda, \rho\}$, evaluates to a nonzero rational number at this point. Though we will not need it, we remark that the Jacobian of F with respect to the other set of seven variables, $\{\alpha, \beta\}$, is also nonzero at this point.

By the implicit function theorem, the set

$$V = \{(\alpha, \beta) \in \mathbb{C}^7 : \exists_{(g, \lambda, \rho) \in \mathbb{C}^7} : F = 0, |J_{\lambda, \rho, g}(F)| \neq 0 \text{ and } g, \lambda_1, \lambda_2 \neq 0\},$$

is open and non-empty. But V is a constructible set with respect to the Zariski topology, so either V or its complement is dense in \mathbb{C}^7 with respect to the Euclidean topology. It follows that V is dense in \mathbb{C}^7 . That is, for generic $(\alpha, \beta) \in \mathbb{C}^7$, we can find $\lambda_1, \lambda_2, \rho_2, \rho_3, \rho_5, \rho_6$, and an affine transformation (which generally depends on g), that transforms the Segre surface, defined by equations (2.12), into (2.1). The proposition follows. \square

From Lemma 2.8, Remark 2.9 and Proposition 2.11, we obtain the following result.

Corollary 2.12. *The embedded family of affine Segre surfaces \mathcal{Z}_q defined in (2.1), constitute a six-dimensional family, up to affine equivalence.*

2.3. Parametrisation. In this section, we introduce and study the parametrisation of the affine Segre surface (2.1) in terms of the parameter of $q\mathbb{P}_{\text{VI}}$, which is defined below. This parametrisation will be helpful for two separate purposes. The first is to give explicit expressions for the lines on the Segre surface. The second is to compute a continuum limit of the Segre surface.

Given $q \in \mathbb{C}$, $0 < |q| < 1$, and $\kappa = (\kappa_0, \kappa_t, \kappa_1, \kappa_\infty) \in (\mathbb{C}^*)^4$, the q -difference sixth Painlevé equation [25] is

$$q\mathbb{P}_{\text{VI}} : \begin{cases} f\bar{f} = \frac{(g - \kappa_0 t)(g - \kappa_0^{-1} t)}{(g - \kappa_\infty)(g - q^{-1}\kappa_\infty^{-1})}, \\ g\bar{g} = \frac{(f - \kappa_t t)(f - \kappa_t^{-1} t)}{q(f - \kappa_1)(f - \kappa_1^{-1})}, \\ f, g : T \rightarrow \mathbb{P}^1 \end{cases} \quad (2.19)$$

where the domain is given by a q -spiral, $T = q^{\mathbb{Z}}t_0$, and $f = f(t)$, $g = g(t)$, $\bar{f} = f(qt)$, $\bar{g} = g(qt)$, for $t \in T$. Here $q^\alpha := e^{\alpha \log q}$ and the parameters of the equation are given by $\kappa_j = q^{\vartheta_j}$ for $j = 0, t, 1, \infty$, with

$$\vartheta_0, \vartheta_t, \vartheta_1, \vartheta_\infty \in \mathbb{C}, \quad t_0 \in \mathbb{C}^*, \quad \log q \in \{x \in \mathbb{C} : \Re x < 0\}. \quad (2.20)$$

In order to define the parametrisation of the $q\mathbb{P}_{\text{VI}}$ Segre surface, we require some elementary q -special functions. Firstly, the q -Pochhammer symbol is defined by the product

$$(z; q)_\infty = \prod_{k=0}^{\infty} (1 - q^k z),$$

which is locally uniformly convergent in $(z, q) \in \mathbb{C} \times \mathbb{D}$, where \mathbb{D} denotes the open unit disc $\mathbb{D} = \{x \in \mathbb{C} : |x| < 1\}$. Secondly, the q -theta function, defined by

$$\theta_q(z) = (q; q)_\infty (z; q)_\infty (q/z; q)_\infty,$$

is analytic in $(z, q) \in \mathbb{C}^* \times \mathbb{D}$, and admits the following convergent expansion on its domain,

$$\theta_q(z) = \sum_{n=-\infty}^{\infty} (-1)^n q^{\frac{1}{2}n(n-1)} z^n. \quad (2.21)$$

This is known as the Jacobi triple product formula, which shows that $\theta_q(z)$ has essential singularities at $z = 0$ and $z = \infty$, unless $q = 0$. For $n \in \mathbb{N}^*$, we use the common abbreviation for repeated products of this function

$$\theta_q(z_1, \dots, z_n) = \theta_q(z_1) \cdot \dots \cdot \theta_q(z_n).$$

Now, the coefficients in (2.2) are explicitly parametrised by the parameters in (2.20) as follows [26],

$$\begin{aligned} \mu_1 &= \prod_{\epsilon=\pm 1} \frac{\theta_q(q^{+\vartheta_\infty} t_0)}{\theta_q(q^{\epsilon\vartheta_0+\vartheta_\infty} t_0)}, & \mu_2 &= \prod_{\epsilon=\pm 1} \frac{\theta_q(q^{-\vartheta_\infty} t_0)}{\theta_q(q^{\epsilon\vartheta_0-\vartheta_\infty} t_0)}, \\ \mu_3 &= \prod_{\epsilon=\pm 1} \frac{\theta_q(q^{\vartheta_t+\vartheta_1+\vartheta_\infty})}{\theta_q(q^{\epsilon\vartheta_0+\vartheta_t+\vartheta_1+\vartheta_\infty})}, & \mu_4 &= \prod_{\epsilon=\pm 1} \frac{\theta_q(q^{-\vartheta_t-\vartheta_1+\vartheta_\infty})}{\theta_q(q^{\epsilon\vartheta_0-\vartheta_t-\vartheta_1+\vartheta_\infty})}, \\ \mu_5 &= \prod_{\epsilon=\pm 1} \frac{\theta_q(q^{\vartheta_t-\vartheta_1+\vartheta_\infty})}{\theta_q(q^{\epsilon\vartheta_0+\vartheta_t-\vartheta_1+\vartheta_\infty})}, & \mu_6 &= \prod_{\epsilon=\pm 1} \frac{\theta_q(q^{-\vartheta_t+\vartheta_1+\vartheta_\infty})}{\theta_q(q^{\epsilon\vartheta_0-\vartheta_t+\vartheta_1+\vartheta_\infty})}, \end{aligned} \quad (2.22)$$

and

$$\lambda_1 = \frac{\eta_3^{(q)} \eta_4^{(q)}}{\eta_1^{(q)} \eta_2^{(q)}}, \quad \lambda_2 = \frac{\eta_5^{(q)} \eta_6^{(q)}}{\eta_1^{(q)} \eta_2^{(q)}}, \quad (2.23)$$

where the $\eta_k^{(q)}$ are given by

$$\begin{aligned} \eta_1^{(q)} &= +\theta_q(q^{\vartheta_0+\vartheta_\infty} t_0, q^{-\vartheta_0+\vartheta_\infty} t_0) \theta_q(q^{2\vartheta_t}, q^{2\vartheta_1}), \\ \eta_2^{(q)} &= +\theta_q(q^{\vartheta_0-\vartheta_\infty} t_0, q^{-\vartheta_0-\vartheta_\infty} t_0) \theta_q(q^{2\vartheta_t}, q^{2\vartheta_1}) q^{2\vartheta_\infty}, \\ \eta_3^{(q)} &= -\theta_q(q^{\vartheta_t-\vartheta_1} t_0, q^{-\vartheta_t+\vartheta_1} t_0) \theta_q(q^{\vartheta_0+\vartheta_t+\vartheta_1+\vartheta_\infty}, q^{-\vartheta_0+\vartheta_t+\vartheta_1+\vartheta_\infty}), \\ \eta_4^{(q)} &= -\theta_q(q^{\vartheta_t-\vartheta_1} t_0, q^{-\vartheta_t+\vartheta_1} t_0) \theta_q(q^{\vartheta_0-\vartheta_t-\vartheta_1+\vartheta_\infty}, q^{-\vartheta_0-\vartheta_t-\vartheta_1+\vartheta_\infty}) q^{2\vartheta_t+2\vartheta_1}, \\ \eta_5^{(q)} &= +\theta_q(q^{\vartheta_t+\vartheta_1} t_0, q^{-\vartheta_t-\vartheta_1} t_0) \theta_q(q^{\vartheta_0+\vartheta_t-\vartheta_1+\vartheta_\infty}, q^{-\vartheta_0+\vartheta_t-\vartheta_1+\vartheta_\infty}) q^{2\vartheta_1}, \\ \eta_6^{(q)} &= +\theta_q(q^{\vartheta_t+\vartheta_1} t_0, q^{-\vartheta_t-\vartheta_1} t_0) \theta_q(q^{\vartheta_0-\vartheta_t+\vartheta_1+\vartheta_\infty}, q^{-\vartheta_0-\vartheta_t+\vartheta_1+\vartheta_\infty}) q^{2\vartheta_t}. \end{aligned}$$

We further note that

$$\mu_k = \frac{\widehat{\eta}_k^{(q)}}{\eta_k^{(q)}}, \quad \widehat{\eta}_k^{(q)} := \eta_k^{(q)}|_{\vartheta_0=0} \quad (1 \leq k \leq 6). \quad (2.24)$$

Under the parametrisation above, the six invariant quantities, $\lambda_1, \lambda_2, \rho_2, \rho_3, \rho_5, \rho_6$, see equation (2.4), depend on six free parameters (2.20). A priori, it is conceivable that there might exist an (algebraic) relation among the coefficients in (2.1) under this parametrisation, but it follows from the following lemma that this is not the case.

Lemma 2.13. *The meromorphic mapping,*

$$(\vartheta_0, \vartheta_t, \vartheta_1, \vartheta_\infty, t_0, \log q) \mapsto (\lambda_1, \lambda_2, \rho_2, \rho_3, \rho_5, \rho_6), \quad (2.25)$$

is locally biholomorphic near almost any point in its domain (2.20).

Proof. It is helpful to write (2.25) as the composition of the mappings

$$(\vartheta_0, \vartheta_t, \vartheta_1, \vartheta_\infty, t_0, \log q) \mapsto (\kappa_0, \kappa_t, \kappa_1, \kappa_\infty, t_0, q), \quad (2.26)$$

$$(\kappa_0, \kappa_t, \kappa_1, \kappa_\infty, t_0, q) \mapsto (\lambda_1, \lambda_2, \rho_2, \rho_3, \rho_5, \rho_6), \quad (2.27)$$

where

$$(\kappa_0, \kappa_t, \kappa_1, \kappa_\infty) = (q^{\vartheta_0}, q^{\vartheta_t}, q^{\vartheta_1}, q^{-\vartheta_\infty}), \quad (2.28)$$

and the domain of (2.27) is defined by

$$\{(\kappa_0, \kappa_t, \kappa_1, \kappa_\infty, t_0, q) \in (\mathbb{C}^*)^5 \times \mathbb{D}\}. \quad (2.29)$$

Clearly, the mapping (2.26) is a local biholomorphism. To prove the lemma, it thus suffices to show that the mapping (2.27) is locally biholomorphic near almost any point in its domain. Since the domain (2.29) is connected, and the mapping (2.27) is meromorphic, it thus suffices to show that the Jacobian determinant of (2.27) is not identically zero. To prove the latter, we are going to compute an expansion of the Jacobian determinant around $q = 0$.

Recall that $\theta_q(z)$ is an analytic function with respect to $(z, q) \in \mathbb{C}^* \times \mathbb{D}$. From the Jacobi triple product formula (2.21), we obtain the following convergent expansion around $q = 0$,

$$\theta_q(z) = \sum_{n=0}^{\infty} (-1)^n (z^{-n} - z^{n+1}) q^{\frac{1}{2}n(n+1)},$$

which allows us to compute expansions around $q = 0$ for λ_1, λ_2 and $\rho_k, k = 2, 3, 5, 6$. For example,

$$\begin{aligned} \lambda_1 &= \frac{\theta_q(t_0 \kappa_t \kappa_1^{-1})^2 \theta_q(t_0 \kappa_t^{-1} \kappa_1)^2}{\theta_q(\kappa_t^2)^2 \theta_q(\kappa_1^2)^2} \prod_{\epsilon_{1,2} \in \{\pm 1\}} \frac{\theta_q(\kappa_0^{\epsilon_1} \kappa_\infty^{\epsilon_2} \kappa_t \kappa_1)}{\theta_q(\kappa_0^{\epsilon_1} \kappa_\infty^{\epsilon_2} t_0)} \\ &= \lambda_1^{(0)} + q \lambda_1^{(1)} + \mathcal{O}(q^2), \end{aligned}$$

as $q \rightarrow 0$, where

$$\begin{aligned} \lambda_1^{(0)} &= \frac{(t_0 \kappa_t \kappa_1^{-1} - 1)(t_0 \kappa_t^{-1} \kappa_1 - 1)}{(\kappa_t^2 - 1)^2 (\kappa_1^2 - 1)^2} \prod_{\epsilon_{1,2} \in \{\pm 1\}} \frac{(\kappa_0^{\epsilon_1} \kappa_\infty^{\epsilon_2} \kappa_t \kappa_1 - 1)}{(\kappa_0^{\epsilon_1} \kappa_\infty^{\epsilon_2} t_0 - 1)}, \\ \lambda_1^{(1)} &= t_0^{-1} \lambda_1^{(0)} (t_0 \kappa_t \kappa_1 - 1)(t_0 \kappa_t^{-1} \kappa_1^{-1} - 1) \\ &\quad \cdot (\kappa_0 \kappa_\infty + \kappa_0^{-1} \kappa_\infty + \kappa_0 \kappa_\infty^{-1} + \kappa_0^{-1} \kappa_\infty^{-1} - 2\kappa_t \kappa_1^{-1} - 2\kappa_t^{-1} \kappa_1). \end{aligned}$$

Similarly, we obtain expansions,

$$\rho_k = \rho_k^{(0)} + q \rho_k^{(1)} + \mathcal{O}(q^2) \quad (q \rightarrow 0),$$

for $k = 2, 3, 5, 6$, with coefficients that are easily computed explicitly.

Let J denote the Jacobian of the mapping (2.27),

$$J = \begin{bmatrix} \frac{\partial \lambda_1}{\partial \kappa_0} & \frac{\partial \lambda_1}{\partial \kappa_t} & \frac{\partial \lambda_1}{\partial \kappa_1} & \frac{\partial \lambda_1}{\partial \kappa_\infty} & \frac{\partial \lambda_1}{\partial t_0} & \frac{\partial \lambda_1}{\partial q} \\ \frac{\partial \lambda_2}{\partial \kappa_0} & \frac{\partial \lambda_2}{\partial \kappa_t} & \frac{\partial \lambda_2}{\partial \kappa_1} & \frac{\partial \lambda_2}{\partial \kappa_\infty} & \frac{\partial \lambda_2}{\partial t_0} & \frac{\partial \lambda_2}{\partial q} \\ \frac{\partial \rho_2}{\partial \kappa_0} & \frac{\partial \rho_2}{\partial \kappa_t} & \frac{\partial \rho_2}{\partial \kappa_1} & \frac{\partial \rho_2}{\partial \kappa_\infty} & \frac{\partial \rho_2}{\partial t_0} & \frac{\partial \rho_2}{\partial q} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \frac{\partial \rho_6}{\partial \kappa_0} & \frac{\partial \rho_6}{\partial \kappa_t} & \frac{\partial \rho_6}{\partial \kappa_1} & \frac{\partial \rho_6}{\partial \kappa_\infty} & \frac{\partial \rho_6}{\partial t_0} & \frac{\partial \rho_6}{\partial q} \end{bmatrix}.$$

Then J admits an expansion around $q = 0$,

$$J = \sum_{k=0}^{\infty} J_k(\kappa_0, \kappa_t, \kappa_1, \kappa_\infty, t_0) q^k,$$

which is locally uniformly convergent with respect to the remaining variables away from singularities of J . Now, a direct computation yields that the determinant of the constant term is given by

$$\begin{aligned}
|J_0| &= \begin{vmatrix} \frac{\partial \lambda_1^{(0)}}{\partial \kappa_0} & \frac{\partial \lambda_1^{(0)}}{\partial \kappa_t} & \frac{\partial \lambda_1^{(0)}}{\partial \kappa_1} & \frac{\partial \lambda_1^{(0)}}{\partial \kappa_\infty} & \frac{\partial \lambda_1^{(0)}}{\partial t_0} & \frac{\partial \lambda_1^{(0)}}{\partial q} \\ \frac{\partial \lambda_2^{(0)}}{\partial \kappa_0} & \frac{\partial \lambda_2^{(0)}}{\partial \kappa_t} & \frac{\partial \lambda_2^{(0)}}{\partial \kappa_1} & \frac{\partial \lambda_2^{(0)}}{\partial \kappa_\infty} & \frac{\partial \lambda_2^{(0)}}{\partial t_0} & \frac{\partial \lambda_2^{(0)}}{\partial q} \\ \frac{\partial \rho_2^{(0)}}{\partial \kappa_0} & \frac{\partial \rho_2^{(0)}}{\partial \kappa_t} & \frac{\partial \rho_2^{(0)}}{\partial \kappa_1} & \frac{\partial \rho_2^{(0)}}{\partial \kappa_\infty} & \frac{\partial \rho_2^{(0)}}{\partial t_0} & \frac{\partial \rho_2^{(0)}}{\partial q} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \frac{\partial \rho_6^{(0)}}{\partial \kappa_0} & \frac{\partial \rho_6^{(0)}}{\partial \kappa_t} & \frac{\partial \rho_6^{(0)}}{\partial \kappa_1} & \frac{\partial \rho_6^{(0)}}{\partial \kappa_\infty} & \frac{\partial \rho_6^{(0)}}{\partial t_0} & \frac{\partial \rho_6^{(0)}}{\partial q} \end{vmatrix} \\
&= 8 \frac{(\kappa_t^{-1} \kappa_1^{-1} t_0 - 1)^4 (\kappa_t \kappa_1^{-1} t_0 - 1)^4 (\kappa_t^{-1} \kappa_1 t_0 - 1)^4}{\kappa_0^2 \kappa_t^2 \kappa_1^2 \kappa_\infty^{-2} (\kappa_t \kappa_1 t_0 - 1) (\kappa_t^{-1} \kappa_1^{-1} \kappa_\infty^{-2} t_0 - 1)^5} \cdot \frac{(\kappa_0^2 - 1)(\kappa_\infty^{-2} - 1)^6}{(\kappa_t^2 - 1)^2 (\kappa_1^2 - 1)^2} \\
&\quad \cdot \prod_{\epsilon \in \{\pm 1\}} \frac{(\kappa_0^\epsilon \kappa_t \kappa_1 \kappa_\infty - 1)^4 (\kappa_0^\epsilon \kappa_\infty^{-1} t_0 - 1)}{(\kappa_0^\epsilon \kappa_\infty t_0 - 1)^4 (\kappa_0^\epsilon \kappa_t^{-1} \kappa_1 \kappa_\infty - 1) (\kappa_0^\epsilon \kappa_t \kappa_1^{-1} \kappa_\infty - 1) (\kappa_0^\epsilon \kappa_t \kappa_1 \kappa_\infty^{-1} - 1)}.
\end{aligned}$$

As a consequence, $|J| \neq 0$ and the lemma follows. \square

Roughly speaking, Lemma 2.8 and Remarks 2.9 and 2.10 allow us to identify generic embedded affine Segre surfaces, up to affine equivalence, with generic tuples $\{(\alpha, \beta) \in \mathbb{C}^8\}$, quotiented by the actions in Remark 2.9. As a consequence of Proposition 2.11 and Lemma 2.13, we find that the parametrisation with respect to the parameters of qP_{VI} in (2.20), maps out a subset with non-empty interior in this quotient space. This yields the following remark.

Remark 2.14. The embedded affine Segre surface for qP_{VI} is of the most generic type. In particular, its projective completion is smooth and the curve at infinity is a smooth irreducible quartic curve of genus 1.

As a consequence, we obtain negations of certain assertions or conjectures made in [40]. In particular, item 7.2.1 in the list of problems in [40, §7.2] contains an open problem and three conjectures. Our result resolves the open problem and negates these conjectures.

2.4. Lines. It is a classical fact, essentially due to Segre himself [48], that a smooth Segre surface contains 16 lines. Half of the lines in $\overline{\mathcal{Z}}_q$ admit a most simple description. Namely, take any $i \in \{1, 2\}$, $j \in \{3, 4\}$ and $k \in \{5, 6\}$, then

$$L_{i,j,k}^{(q)} := \{Z \in \overline{\mathcal{Z}}_q : Z_i = Z_j = Z_k = 0\}, \quad (2.30)$$

defines a line in $\overline{\mathcal{Z}}_q$.

For the remaining eight lines, we make use of the parametrisation in Section 2.3. To describe them, certain rational functions on $\overline{\mathcal{Z}}_q$, called Tyurin ratios, are helpful. To define these rational functions, note that equations (2.5c) and (2.5d) imply

$$\frac{Z_1}{\eta_1^{(q)}} \cdot \frac{Z_2}{\eta_2^{(q)}} = \frac{Z_3}{\eta_3^{(q)}} \cdot \frac{Z_4}{\eta_4^{(q)}} = \frac{Z_5}{\eta_5^{(q)}} \cdot \frac{Z_6}{\eta_6^{(q)}}.$$

We now consider the following rational function,

$$T_{13}^{(q)} := \frac{Z_1}{\eta_1^{(q)}} / \frac{Z_3}{\eta_3^{(q)}} = \frac{Z_4}{\eta_4^{(q)}} / \frac{Z_2}{\eta_2^{(q)}}.$$

This is a meromorphic function on the Segre surface,

$$T_{13}^{(q)} : \overline{\mathcal{Z}}_q \rightarrow \mathbb{P}^1,$$

as there is no point on the Segre surface with $Z_1 = Z_2 = Z_3 = Z_4 = 0$.




FIGURE 2.1. Clebsch graph encoding the configuration of lines and their points of intersection on the Segre surface \bar{Z}_q . The lines are explicitly described in Section 2.4 for $0 < |q| < 1$ and in Section 3.1.1 when $q = 1$. Each vertex represents a line and each edge represents an intersection point of the two lines corresponding to its endpoints. The lines are colour-coded ForestGreen or Fuchsia dependent on whether, when $q = 1$, they intersect the curve at infinity of \bar{Z}_1 respectively in conic (3.20) or conic (3.21).

Let $\alpha \in S_6$ denote the permutation,

$$\alpha = (1\ 2)(3\ 4)(5\ 6), \quad (2.31)$$

then, for any $1 \leq i, j \leq 6$ not in a same cycle of α , we similarly have a meromorphic rational function

$$T_{ij}^{(q)} := \frac{Z_i}{\eta_i^{(q)}} \Big/ \frac{Z_j}{\eta_j^{(q)}} = \frac{Z_{\alpha(j)}}{\eta_{\alpha(j)}^{(q)}} \Big/ \frac{Z_{\alpha(i)}}{\eta_{\alpha(i)}^{(q)}}. \quad (2.32)$$

We call these functions Tyurin ratios, as they can be interpreted as ratios of Tyurin parameters of some elliptic matrix functions, see [26, §2.4] or [42, §2.2]. We have the following obvious symmetries

$$T_{ij}^{(q)} = (T_{ji}^{(q)})^{-1} = T_{\alpha(j)\alpha(i)}^{(q)} = (T_{\alpha(i)\alpha(j)}^{(q)})^{-1}, \quad (2.33)$$

so that, the $6 \times 4 = 24$ choices of indices, only yield 6 Tyurin ratios which are not trivially equivalent. We further have the following multiplicative formula,

$$T_{ij}^{(q)} T_{jk}^{(q)} = T_{ik}^{(q)}, \quad (2.34)$$

for any choice of $1 \leq i, j, k \leq 6$ in mutually disjoint cycles of α .

The remaining eight lines can now be described as follows. For any $i \in \{1, 2\}$, $j \in \{3, 4\}$ and $k \in \{5, 6\}$, the following set is a line in $\overline{\mathcal{Z}}_q$,

$$\tilde{L}_{i,j,k}^{(q)} = \{Z \in \overline{\mathcal{Z}}_q : T_{ij}^{(q)}(Z) = \tau_i^{(q)}/\tau_j^{(q)}, T_{jk}^{(q)}(Z) = \tau_j^{(q)}/\tau_k^{(q)}\}, \quad (2.35)$$

where

$$\begin{aligned} \tau_1^{(q)} &= \frac{\theta_q(t_0^{+1}q^{-\vartheta_\infty - \vartheta_0})}{\theta_q(t_0^{+1}q^{+\vartheta_\infty - \vartheta_0})}, & \tau_3^{(q)} &= \frac{\theta_q(q^{+\vartheta_t + \vartheta_1 - \vartheta_\infty - \vartheta_0})}{\theta_q(q^{+\vartheta_t + \vartheta_1 + \vartheta_\infty - \vartheta_0})}, & \tau_5^{(q)} &= \frac{\theta_q(q^{+\vartheta_t - \vartheta_1 - \vartheta_\infty - \vartheta_0})}{\theta_q(q^{+\vartheta_t - \vartheta_1 + \vartheta_\infty - \vartheta_0})}, \\ \tau_2^{(q)} &= \frac{\theta_q(t_0^{-1}q^{-\vartheta_\infty - \vartheta_0})}{\theta_q(t_0^{-1}q^{+\vartheta_\infty - \vartheta_0})}, & \tau_4^{(q)} &= \frac{\theta_q(q^{-\vartheta_t - \vartheta_1 - \vartheta_\infty - \vartheta_0})}{\theta_q(q^{-\vartheta_t - \vartheta_1 + \vartheta_\infty - \vartheta_0})}, & \tau_6^{(q)} &= \frac{\theta_q(q^{-\vartheta_t + \vartheta_1 - \vartheta_\infty - \vartheta_0})}{\theta_q(q^{-\vartheta_t + \vartheta_1 + \vartheta_\infty - \vartheta_0})}. \end{aligned}$$

These descriptions of the remaining eight lines follow from the explicit expressions for the lines in [42, §3.3], where we note the following correspondence with the notation used in that paper,

$$\begin{aligned} \tilde{L}_{1,3,5}^{(q)} &= \tilde{\mathcal{L}}_1^0, & \tilde{L}_{1,4,6}^{(q)} &= \tilde{\mathcal{L}}_2^0, & \tilde{L}_{2,3,6}^{(q)} &= \tilde{\mathcal{L}}_3^0, & \tilde{L}_{2,4,5}^{(q)} &= \tilde{\mathcal{L}}_4^0, \\ \tilde{L}_{2,4,6}^{(q)} &= \tilde{\mathcal{L}}_1^\infty, & \tilde{L}_{2,3,5}^{(q)} &= \tilde{\mathcal{L}}_2^\infty, & \tilde{L}_{1,4,5}^{(q)} &= \tilde{\mathcal{L}}_3^\infty, & \tilde{L}_{1,3,6}^{(q)} &= \tilde{\mathcal{L}}_4^\infty. \end{aligned}$$

The correspondence between the two notations for the other eight lines is the same, with all the tildes removed. We further recall from [42, Theorem 2.11] the intersection graph of lines in Figure 2.1.

3. THE LIMIT OF \mathcal{Z}_q AS $q \uparrow 1$ AND JIMBO-FRICKE CUBIC FOR P_{VI}

In the singular limit $q \uparrow 1$, known as the continuum limit, qP_{VI} formally reduces to the classical sixth Painlevé equation. To see this explicitly, one substitutes formal Taylor series around $q = 1$,

$$\begin{aligned} f(t) &= f_0(t) + (q-1)f_1(t) + (q-1)^2f_2(t) + \dots, \\ g(t) &= g_0(t) + (q-1)g_1(t) + (q-1)^2g_2(t) + \dots, \end{aligned}$$

and compares terms, which leads to

$$f_0 = u, \quad g_0 = \frac{u-t}{u-1},$$

where u satisfies the sixth Painlevé equation

$$\begin{aligned} P_{VI}: \quad u_{tt} &= \left(\frac{1}{u} + \frac{1}{u-1} + \frac{1}{u-t} \right) \frac{u_t^2}{2} - \left(\frac{1}{t} + \frac{1}{t-1} + \frac{1}{u-t} \right) u_t \\ &+ \frac{u(u-1)(u-t)}{2t^2(t-1)^2} \left((2\vartheta_\infty - 1)^2 - \frac{4\vartheta_0^2 t}{u^2} + \frac{4\vartheta_1^2(t-1)}{(u-1)^2} \right. \\ &\quad \left. + \frac{(1-4\vartheta_t^2)t(t-1)}{(u-t)^2} \right). \end{aligned} \quad (3.1)$$

In this section, we correspondingly compute the continuum limit of the Segre surface \mathcal{Z}_q and compare it to the Jimbo-Fricke cubic for P_{VI} . The main result of the section, Theorem 3.8, shows that the two are isomorphic as affine varieties. More precisely, the theorem states that the two are affinely equivalent after blowing down one of the lines at infinity of the cubic.

In Section 3.1, the continuum limit is worked out, leading to an affine Segre surface \mathcal{Z}_1 . We further give an algebraic characterisation of this surface and show that the descriptions of the lines on \mathcal{Z}_q remain intact as $q \uparrow 1$.

In Section 3.2, we blow down the Jimbo-Fricke cubic surface to a Segre surface and describe the lines on both surfaces. In Section 3.3 we explain how to find an isomorphism between the resulting Segre surface and the Turyn ratios. Then, in Section 3.4, it is shown that the Jimbo-Fricke cubic is affinely isomorphic to \mathcal{Z}_1 .

3.1. The limit of \mathcal{Z}_q as $q \uparrow 1$. With regards to the Segre surface \mathcal{Z}_q , note that all its coefficients, see equations (2.22) and (2.23), are rational in terms of simple factors of the form $\theta_q(q^\alpha)$ and $\theta_q(q^\alpha t_0)$, $\alpha \in \mathbb{C}$. To compute the limits of the coefficients as $q \uparrow 1$, it is thus sufficient to understand the limiting behaviours of these simple factors. Correspondingly, we have the following lemma.

Lemma 3.1. *For any $\alpha \in \mathbb{C}$ and $t_0 \in \mathbb{C}^*$, with $\arg(-t_0) < \pi$, we have the following converging limits*

$$\lim_{q \uparrow 1} (1-q)^{-1} \frac{\theta_q(q^\alpha)}{(q; q)_\infty^3} = \frac{\sin \pi \alpha}{\pi}, \quad \lim_{q \uparrow 1} \frac{\theta_q(q^\alpha t_0)}{\theta_q(t_0)} = (-t_0)^{-\alpha}.$$

Proof. To derive the first limit, we recall that the q -gamma function, defined by

$$\Gamma_q(\alpha) = (1-q)^{1-\alpha} \frac{(q; q)_\infty}{(q^\alpha; q)_\infty},$$

converges to the usual gamma function as $q \uparrow 1$. Therefore,

$$\theta_q(q^\alpha) = \frac{(1-q)(q; q)_\infty^3}{\Gamma_q(\alpha)\Gamma_q(1-\alpha)} \sim \frac{(1-q)(q; q)_\infty^3}{\Gamma(\alpha)\Gamma(1-\alpha)} = (1-q)(q; q)_\infty^3 \frac{\sin \pi \alpha}{\pi},$$

as $q \uparrow 1$, from which the first limit in the lemma follows. The second is a direct consequence of

$$\lim_{q \uparrow 1} \frac{(q^\alpha z; q)_\infty}{(z; q)_\infty} = (1-z)^{-\alpha} \quad (z \in \mathbb{C} \setminus [1, \infty)),$$

which we take from [17, Eq. (1.3.19)]. \square

As a consequence of the above lemma, we have the following limits for ratios of q -theta functions,

$$\frac{\theta_q(q^\alpha)}{\theta_q(q^\beta)} \rightarrow \frac{\sin \pi \alpha}{\sin \pi \beta} \quad (\beta \notin \mathbb{Z}), \quad \frac{\theta_q(q^\alpha t_0)}{\theta_q(q^\beta t_0)} \rightarrow (-t_0)^{\beta-\alpha} \quad (\arg(-t_0) < \pi). \quad (3.2)$$

From these identities, we obtain the continuum limit of the Segre surface \mathcal{Z}_q for qP_{VI} , which is a family of algebraic varieties depending on four generic parameters,

$$\vartheta_0, \vartheta_t, \vartheta_1, \vartheta_\infty \in \mathbb{C}, \quad (3.3)$$

defined as the zero set of four polynomials $h_i \in \mathbb{C}[z_1, \dots, z_6]$, $i = 1, \dots, 4$ given by (2.1) with the choice of parameters defined by

$$\begin{aligned} \mu_1 &= 1, & \mu_2 &= 1, & (3.4) \\ \mu_3 &= \prod_{\epsilon=\pm 1} \frac{\sin(\pi(\vartheta_t + \vartheta_1 + \vartheta_\infty))}{\sin(\pi(\epsilon\vartheta_0 + \vartheta_t + \vartheta_1 + \vartheta_\infty))}, & \mu_4 &= \prod_{\epsilon=\pm 1} \frac{\sin(\pi(-\vartheta_t - \vartheta_1 + \vartheta_\infty))}{\sin(\pi(\epsilon\vartheta_0 - \vartheta_t - \vartheta_1 + \vartheta_\infty))}, \\ \mu_5 &= \prod_{\epsilon=\pm 1} \frac{\sin(\pi(\vartheta_t - \vartheta_1 + \vartheta_\infty))}{\sin(\pi(\epsilon\vartheta_0 + \vartheta_t - \vartheta_1 + \vartheta_\infty))}, & \mu_6 &= \prod_{\epsilon=\pm 1} \frac{\sin(\pi(-\vartheta_t + \vartheta_1 + \vartheta_\infty))}{\sin(\pi(\epsilon\vartheta_0 - \vartheta_t + \vartheta_1 + \vartheta_\infty))}, \end{aligned}$$

and

$$\begin{aligned} \lambda_1 &= \prod_{\epsilon=\pm 1} \frac{\sin(\pi(\epsilon\vartheta_0 - \vartheta_t - \vartheta_1 + \vartheta_\infty)) \sin(\pi(\epsilon\vartheta_0 + \vartheta_t + \vartheta_1 + \vartheta_\infty))}{\sin(2\pi\vartheta_t) \sin(2\pi\vartheta_1)}, & (3.5) \\ \lambda_2 &= \prod_{\epsilon=\pm 1} \frac{\sin(\pi(\epsilon\vartheta_0 - \vartheta_t + \vartheta_1 + \vartheta_\infty)) \sin(\pi(\epsilon\vartheta_0 + \vartheta_t - \vartheta_1 + \vartheta_\infty))}{\sin(2\pi\vartheta_t) \sin(2\pi\vartheta_1)}. \end{aligned}$$

Note that the t_0 dependence of the coefficients drops out in the continuum limit.

Lemma 3.2. *Equations (2.1), with coefficients given by (3.4) and (3.5), define a four-parameter family of affine Segre surfaces. Among the coefficients, there are the following non-trivial relations,*

$$\frac{\lambda_1}{\lambda_2} = \frac{(\mu_5 - 1)(\mu_6 - 1)}{(\mu_3 - 1)(\mu_4 - 1)}, \quad \mu_1 = \mu_2 = 1, \quad (3.6)$$

and these are effectively all, that is, any other relation among the coefficients is generated by them.

Similarly to the case of qPVI, we can introduce the parameters ρ defined in (2.4) which in the case of $q \uparrow 1$ become:

$$\rho_2 = 0, \quad \rho_3 = \frac{\mu_3 - 1}{\mu_4 - 1}, \quad \rho_5 = \frac{\mu_5 - 1}{\mu_4 - 1}, \quad \rho_6 = \frac{\mu_6 - 1}{\mu_4 - 1}, \quad (3.7)$$

and replace h_2 by h_2'' defined in (2.3).

Definition 3.3. We denote the family of affine surfaces defined by equations (2.1), with coefficients given by (3.4) and (3.5) by \mathcal{Z}_1 . The family \mathcal{Z}_1 is called *PVI Segre*.

Proof. Equations (3.6) follow by direct computation. Checking them by hand is easiest via equations (3.17) and (3.19) in Section 3.1.1.

In terms of the parameters λ_1, λ_2 and $\rho_2, \rho_3, \rho_5, \rho_6$ introduced in (3.7), relations (3.6) yield

$$\frac{\lambda_1}{\lambda_2} = \frac{\rho_5 \rho_6}{\rho_3}, \quad \rho_2 = 0. \quad (3.8)$$

To prove that these are effectively all the relations among the coefficients, we consider the mapping defined by (3.4), (3.5) and (3.7),

$$(\theta_0, \theta_t, \theta_1, \theta_\infty) \mapsto (\lambda_1, \lambda_2, \rho_5, \rho_6).$$

The Jacobian determinant of this mapping equals

$$-128\pi^4 \frac{(v_0 - v_0^{-1})(v_\infty - v_\infty^{-1})^3}{(v_t - v_t^{-1})^3(v_1 - v_1^{-1})^3} \prod_{\epsilon=\pm 1} \frac{(v_0^\epsilon v_\infty - v_t v_1)^2}{(v_0^\epsilon v_t - v_1 v_\infty)(v_0^\epsilon v_1 - v_t v_\infty)},$$

where we used the notation $v_k = e^{2\pi i \theta_k}$ for $k = 0, t, 1, \infty$. In particular, this mapping is locally biholomorphic near generic points in its domain, so that there are no relations among the coefficients, other than those generated by (3.8). \square

Proposition 3.4. *For generic parameter values $(\vartheta_0, \vartheta_t, \vartheta_1, \vartheta_\infty) \in \mathbb{C}^4$, the canonical projective completion $\overline{\mathcal{Z}}_1 \subseteq \mathbb{P}^6$, of the affine Segre surface \mathcal{Z}_1 , is smooth and the curve at infinity is reducible; it consists of two conics which intersect at two points.*

Proof. We use projective coordinates,

$$[Z_0 : Z_1 : Z_3 : Z_4 : Z_5 : Z_6] = [1 : z_1 : z_2 : z_3 : z_4 : z_5 : z_6],$$

so that $\overline{\mathcal{Z}}_1$ is described by equations (2.5), with coefficients satisfying (3.6).

Following the first part of the proof of Proposition 2.4, for $\overline{\mathcal{Z}}_1$ to be singular, it is necessary that $\lambda_1 = 0, \lambda_2 = 0$ or

$$(\lambda_1 - \lambda_2)^2 - 2(\lambda_1 + \lambda_2) + 1 = 0. \quad (3.9)$$

But the left-hand side reads

$$(\lambda_1 - \lambda_2)^2 - 2(\lambda_1 + \lambda_2) + 1 = \frac{\sin(2\pi\vartheta_0) \sin(2\pi\vartheta_\infty)}{\sin(2\pi\vartheta_t) \sin(2\pi\vartheta_1)},$$

which is clearly nonzero for generic parameter values. In fact, for any choice of parameters $(\vartheta_0, \vartheta_t, \vartheta_1, \vartheta_\infty) \in \mathbb{C}^4$, such that

$$2\vartheta_0, 2\vartheta_t, 2\vartheta_1, 2\vartheta_\infty \notin \mathbb{Z}, \quad \epsilon_0\vartheta_0 + \epsilon_t\vartheta_t + \epsilon_1\vartheta_1 + \epsilon_\infty\vartheta_\infty \notin \mathbb{Z}, \quad (3.10)$$

for any $\epsilon_j \in \{\pm 1\}$, $j = 0, t, 1, \infty$, the Segre surface \overline{Z}_1 is well-defined and smooth.

Next, we consider the curve at infinity, described by

$$Z_1 + Z_2 + Z_3 + Z_4 + Z_5 + Z_6 = 0, \quad (3.11a)$$

$$\rho_3 Z_3 + Z_4 + \rho_5 Z_5 + \rho_6 Z_6 = 0, \quad (3.11b)$$

$$Z_3 Z_4 - Z_1 Z_2 \lambda_1 = 0, \quad (3.11c)$$

$$Z_5 Z_6 - Z_1 Z_2 \lambda_2 = 0. \quad (3.11d)$$

Due to the special conditions (3.8) on the coefficients, this curve is reducible. To see this, note that equations (3.11c), (3.11d) imply

$$Z_3 Z_4 - Z_5 Z_6 \frac{\lambda_1}{\lambda_2} = 0. \quad (3.12)$$

Now, recall from (3.8) that $\lambda_1/\lambda_2 = \rho_5 \rho_6 / \rho_3$, and by solving equation (3.11c) for Z_3 and substituting the result into (3.12), we obtain the following factorisation

$$(Z_4 + \rho_6 Z_6)(Z_4 + \rho_5 Z_5) = 0.$$

It follows that the curve at infinity consists of two components. The first, is described by

$$\rho_3 Z_3 = -\rho_5 Z_5, \quad Z_4 = -\rho_6 Z_6, \quad (3.13)$$

and

$$\begin{aligned} Z_1 + Z_2 &= \left(\frac{\rho_5}{\rho_3} - 1\right) Z_5 + (\rho_6 - 1) Z_6, \\ Z_1 Z_2 &= \lambda_2^{-1} Z_5 Z_6. \end{aligned}$$

The second, is described by

$$\rho_3 Z_3 = -\rho_6 Z_6, \quad Z_4 = -\rho_5 Z_5, \quad (3.14)$$

and

$$\begin{aligned} Z_1 + Z_2 &= (\rho_5 - 1) Z_5 + \left(\frac{\rho_6}{\rho_3} - 1\right) Z_6, \\ Z_1 Z_2 &= \lambda_2^{-1} Z_5 Z_6. \end{aligned}$$

For generic parameter values, both of these components are smooth, and thus irreducible, conics.

Next, we consider where the conics intersect. At an intersection point, we have

$$Z_0 = 0, \quad \rho_3 Z_3 = Z_4 = -\rho_5 Z_5 = -\rho_6 Z_6.$$

It follows that the intersection points are given by

$$Z_0 = 0, \quad Z_3 = \rho_3^{-1}, \quad Z_4 = 1, \quad Z_5 = -\rho_5^{-1}, \quad Z_6 = -\rho_6^{-1},$$

with $\{Z_1, Z_2\}$ solutions of

$$\begin{aligned} Z_1 + Z_2 &= -\rho_3^{-1} - 1 + \rho_5^{-1} + \rho_6^{-1}, \\ Z_1 Z_2 &= \lambda_2^{-1} Z_5 Z_6. \end{aligned}$$

The latter system clearly has two distinct solutions for generic parameter values, so that the two conics intersect at two points. The proposition follows. \square

For an explicit description of the intersection points of the two conics in the proof of the proposition above, it is helpful to compute the continuum limits of the

$\eta_k^{(q)}$ $1, \leq k \leq 6$, in equations (2.24). These will also be helpful when describing the continuum limits of the lines. We thus compute,

$$\lim_{q \uparrow 1} \frac{\eta_k^{(q)}}{\theta_q(q^{\frac{1}{2}})\theta_q(t_0)^2} = \begin{cases} (-t_0)^{-2\theta_\infty} \eta_1^{(1)} & \text{if } k = 1, \\ (-t_0)^{+2\theta_\infty} \eta_2^{(1)} & \text{if } k = 2, \\ \eta_k^{(1)} & \text{if } 3 \leq k \leq 6, \end{cases} \quad (3.15)$$

where

$$\begin{aligned} \eta_1^{(1)} &= \eta_2^{(1)} = +\sin(2\pi\vartheta_t)\sin(2\pi\vartheta_1), \\ \eta_3^{(1)} &= -\sin(\pi(\vartheta_0 + \vartheta_t + \vartheta_1 + \vartheta_\infty))\sin(\pi(-\vartheta_0 + \vartheta_t + \vartheta_1 + \vartheta_\infty)), \\ \eta_4^{(1)} &= -\sin(\pi(\vartheta_0 - \vartheta_t - \vartheta_1 + \vartheta_\infty))\sin(\pi(-\vartheta_0 - \vartheta_t - \vartheta_1 + \vartheta_\infty)), \\ \eta_5^{(1)} &= +\sin(\pi(\vartheta_0 + \vartheta_t - \vartheta_1 + \vartheta_\infty))\sin(\pi(-\vartheta_0 + \vartheta_t - \vartheta_1 + \vartheta_\infty)), \\ \eta_6^{(1)} &= +\sin(\pi(\vartheta_0 - \vartheta_t + \vartheta_1 + \vartheta_\infty))\sin(\pi(-\vartheta_0 - \vartheta_t + \vartheta_1 + \vartheta_\infty)). \end{aligned} \quad (3.16)$$

Writing $\widehat{\eta}_k^{(1)} = \eta_k^{(1)}|_{\vartheta_0=0}$, $1 \leq k \leq 6$, we obtain the following expressions for the coefficients of \mathcal{Z}_1 ,

$$\lambda_1 = \frac{\eta_3^{(1)}\eta_4^{(1)}}{\eta_1^{(1)}\eta_2^{(1)}}, \quad \lambda_2 = \frac{\eta_5^{(1)}\eta_6^{(1)}}{\eta_1^{(1)}\eta_2^{(1)}}, \quad \mu_k = \frac{\widehat{\eta}_k^{(1)}}{\eta_k^{(1)}} \quad (1 \leq k \leq 6). \quad (3.17)$$

Moreover, we have the following special relations,

$$\eta_3^{(1)} - \widehat{\eta}_3^{(1)} = \eta_4^{(1)} - \widehat{\eta}_4^{(1)} = \sin(\pi\theta_0)^2 = \widehat{\eta}_5^{(1)} - \eta_5^{(1)} = \widehat{\eta}_6^{(1)} - \eta_6^{(1)}. \quad (3.18)$$

from which we obtain the following expressions

$$\mu_k = \begin{cases} 1 - \frac{\sin(\pi\vartheta_0)^2}{\eta_k^{(1)}} & \text{if } k = 3, 4, \\ 1 + \frac{\sin(\pi\vartheta_0)^2}{\eta_k^{(1)}} & \text{if } k = 5, 6. \end{cases} \quad (3.19)$$

Remark 3.5. Using equations (3.17) and (3.19), we can rewrite the equations for the two conics at infinity in the proof of Proposition 3.4 as follows. The first conic, see equation (3.13), is given by

$$\begin{aligned} Z_0 &= 0, \quad \frac{Z_3}{\eta_3^{(1)}} = \frac{Z_5}{\eta_5^{(1)}}, \quad \frac{Z_4}{\eta_4^{(1)}} = \frac{Z_6}{\eta_6^{(1)}}, \\ Z_1 + Z_2 + (\eta_3^{(1)} + \eta_5^{(1)})\frac{Z_5}{\eta_5^{(1)}} + (\eta_4^{(1)} + \eta_6^{(1)})\frac{Z_6}{\eta_6^{(1)}} &= 0, \quad \frac{Z_1 Z_2}{\eta_1^{(1)}\eta_2^{(1)}} = \frac{Z_5 Z_6}{\eta_5^{(1)}\eta_6^{(1)}}. \end{aligned} \quad (3.20)$$

The second, see equation (3.14), is given by

$$\begin{aligned} Z_0 &= 0, \quad \frac{Z_3}{\eta_3^{(1)}} = \frac{Z_6}{\eta_6^{(1)}}, \quad \frac{Z_4}{\eta_4^{(1)}} = \frac{Z_5}{\eta_5^{(1)}}, \\ Z_1 + Z_2 + (\eta_4^{(1)} + \eta_5^{(1)})\frac{Z_5}{\eta_5^{(1)}} + (\eta_3^{(1)} + \eta_6^{(1)})\frac{Z_6}{\eta_6^{(1)}} &= 0, \quad \frac{Z_1 Z_2}{\eta_1^{(1)}\eta_2^{(1)}} = \frac{Z_5 Z_6}{\eta_5^{(1)}\eta_6^{(1)}}. \end{aligned} \quad (3.21)$$

Their two intersection points are given by

$$Z_0 = 0, \quad Z_1 = \eta_1^{(1)}e^{+2\pi i\vartheta_\infty\epsilon}, \quad Z_2 = \eta_2^{(1)}e^{-2\pi i\vartheta_\infty\epsilon}, \quad Z_k = \eta_k^{(1)} \quad (3 \leq k \leq 6),$$

with $\epsilon = \pm 1$.

3.1.1. *Continuum limit of the lines.* As $q \uparrow 1$, the descriptions of the lines on $\overline{\mathcal{Z}}_q$, in Section 2.4, remain intact and define lines on $\overline{\mathcal{Z}}_1$.

Firstly, regarding the eight lines defined in equation (2.30), the corresponding lines as $q \uparrow 1$ are given as follows. For $i \in \{1, 2\}$, $j \in \{3, 4\}$ and $k \in \{5, 6\}$, the line $L_{i,j,k}^{(q)}$ becomes

$$L_{i,j,k}^{(1)} := \{Z \in \overline{\mathcal{Z}}_1 : Z_i = Z_j = Z_k = 0\},$$

as $q \uparrow 1$, which defines a line in $\overline{\mathcal{Z}}_1$.

To describe the limit of the remaining eight lines, we extend the definition of Tyurin ratios to $\overline{\mathcal{Z}}_1$, by setting

$$T_{ij}^{(1)} := \frac{Z_i}{\eta_i^{(1)}} \Big/ \frac{Z_j}{\eta_j^{(1)}} = \frac{Z_{\alpha(j)}}{\eta_{\alpha(j)}^{(1)}} \Big/ \frac{Z_{\alpha(i)}}{\eta_{\alpha(i)}^{(1)}},$$

when $q = 1$, for any i, j not in the same cycle of the permutation α , defined in equation (2.31).

The eight lines defined in equation (2.35) then converge to the following in the continuum limit. For any $i \in \{1, 2\}$, $j \in \{3, 4\}$ and $k \in \{5, 6\}$, the line $\tilde{L}_{i,j,k}^{(q)}$ becomes

$$\tilde{L}_{i,j,k}^{(1)} = \{Z \in \overline{\mathcal{Z}}_1 : T_{ij}^{(1)}(Z) = \tau_i^{(1)}/\tau_j^{(1)}, T_{jk}^{(1)}(Z) = \tau_j^{(1)}/\tau_k^{(1)}\},$$

as $q \uparrow 1$, where

$$\begin{aligned} \tau_1^{(1)} = 1, \quad \tau_3^{(1)} &= \frac{\sin \pi (+\vartheta_t + \vartheta_1 - \vartheta_\infty - \vartheta_0)}{\sin \pi (+\vartheta_t + \vartheta_1 + \vartheta_\infty - \vartheta_0)}, & \tau_5^{(1)} &= \frac{\sin \pi (+\vartheta_t - \vartheta_1 - \vartheta_\infty - \vartheta_0)}{\sin \pi (+\vartheta_t - \vartheta_1 + \vartheta_\infty - \vartheta_0)}, \\ \tau_2^{(1)} = 1, \quad \tau_4^{(1)} &= \frac{\sin \pi (-\vartheta_t - \vartheta_1 - \vartheta_\infty - \vartheta_0)}{\sin \pi (-\vartheta_t - \vartheta_1 + \vartheta_\infty - \vartheta_0)}, & \tau_6^{(1)} &= \frac{\sin \pi (-\vartheta_t + \vartheta_1 - \vartheta_\infty - \vartheta_0)}{\sin \pi (-\vartheta_t + \vartheta_1 + \vartheta_\infty - \vartheta_0)}. \end{aligned}$$

We further note that the intersection graph of lines in Figure 3.1 remains valid when $q = 1$. Furthermore, we have colour-coded the lines in the intersection graph in accordance to which conic at infinity in Remark 3.5 they intersect.

3.2. The Segre surface of the $\mathbf{P}_{\mathbf{VI}}$ equation. In this section, we blow down the monodromy manifold of the differential sixth Painlevé equation along a line to obtain a corresponding Segre surface. There are generally a number of ways to do this, but we are going to show that it is possible to choose a way that produces a surface affinely equivalent to \mathcal{Z}_1 .

The monodromy manifold of the sixth Painlevé equation is given by the character variety

$$\{(M_0, M_t, M_1) \in SL_2(\mathbb{C})^3\} // SL_2(\mathbb{C}),$$

which, through trace coordinates

$$\begin{aligned} x_1 &= \text{Tr } M_t M_1, & x_2 &= \text{Tr } M_0 M_1, & x_3 &= \text{Tr } M_0 M_t, \\ \nu_0 &= \text{Tr } M_0, & \nu_t &= \text{Tr } M_t, & \nu_1 &= \text{Tr } M_1, & \nu_\infty &= \text{Tr } M_1 M_t M_0, \end{aligned}$$

is identified with the hypersurface in \mathbb{C}^7 defined by

$$x_1 x_2 x_3 + x_1^2 + x_2^2 + x_3^2 + \omega_1 x_1 + \omega_2 x_2 + \omega_3 x_3 + \omega_4 = 0, \quad (3.22)$$

where

$$\begin{aligned} \omega_1 &:= -(\nu_0 \nu_\infty + \nu_t \nu_1), & \omega_2 &:= -(\nu_0 \nu_1 + \nu_t \nu_\infty), & \omega_3 &:= -(\nu_0 \nu_t + \nu_1 \nu_\infty), \\ \omega_4 &:= \nu_0^2 + \nu_t^2 + \nu_1^2 + \nu_\infty^2 + \nu_0 \nu_t \nu_1 \nu_\infty - 4. \end{aligned} \quad (3.23)$$

In the theory of Painlevé VI, the traces ν_k are typically considered fixed, and related to the parameters ϑ_k of the nonlinear ODE, by

$$\nu_k = v_k + v_k^{-1}, \quad v_k := e^{2\pi i \vartheta_k} \quad (k = 0, t, 1, \infty). \quad (3.24)$$

The algebraic equation (3.22) then defines an embedded affine cubic surface with respect to the remaining variables, known as the Jimbo-Fricke cubic surface. We remark that the variables v_k , $k = 0, t, 1, \infty$, are very useful for our purposes, since every computation that follows can essentially be done over the field $\mathbb{Q}(v_0, v_t, v_1, v_\infty)$, and thus be carried out in a computer algebra system.

Let $\mathcal{X} \subseteq \mathbb{C}^3$ denote the embedded family of affine cubic surfaces defined by equations (3.22). Using projective coordinates,

$$[X_0 : X_1 : X_2 : X_3] = [1 : x_1 : x_2 : x_3],$$

its canonical projective completion $\overline{\mathcal{X}} \subseteq \mathbb{P}^3$, is described by

$$X_1 X_2 X_3 + (X_1^2 + X_2^2 + X_3^2)X_0 + (\omega_1 X_1 + \omega_2 X_2 + \omega_3 X_3)X_0^2 + \omega_4 X_0^3 = 0. \quad (3.25)$$

The hypersurface at infinity is a tritangent plane of the cubic, so that the hyperplane section at infinity is a triangle of lines,

$$\overline{\mathcal{X}} \setminus \mathcal{X} = L_1^\infty \cup L_2^\infty \cup L_3^\infty,$$

where

$$L_k^\infty := \{[0 : X_1 : X_2 : X_3] \in \mathbb{P}^3 : X_k = 0\} \quad (k = 1, 2, 3).$$

Moreover, for any choice of parameters, there is no singularity at infinity. Conversely, any embedded affine cubic surface, with a hyperplane section at infinity given by a triangle of lines, consisting of only smooth points, can be brought into the form (3.22) by an affine transformation [36, Proposition 3.2].

3.2.1. Lines. For generic values of the parameters, the cubic surface $\overline{\mathcal{X}}$ is smooth. In fact, the precise conditions are given by (3.10), see [23]. According to the Cayley-Salmon theorem, in addition to the three lines at infinity, there must be a further 24 distinct lines on the cubic. They were first written down explicitly in [29]. Their explicit descriptions are necessary for the proof of Theorem 3.8 and so we provide them here. We give the 24 lines in three sets of eight, such that the lines in the k -th set intersect with the line L_k^∞ at infinity, for $1 \leq k \leq 3$. The first eight lines are given by

$$\begin{aligned} L_1 : \quad & x_1 = v_0 v_\infty + \frac{1}{v_0 v_\infty}, & v_0 v_\infty x_2 + x_3 &= v_0 \nu_t + v_\infty \nu_1, \\ L_2 : \quad & x_1 = v_0 v_\infty + \frac{1}{v_0 v_\infty}, & v_0 v_\infty x_3 + x_2 &= v_0 \nu_1 + v_\infty \nu_t, \\ L_3 : \quad & x_1 = \frac{v_0}{v_\infty} + \frac{v_\infty}{v_0}, & v_0 v_\infty \nu_t + \nu_1 &= v_0 x_2 + v_\infty x_3, \\ L_4 : \quad & x_1 = \frac{v_0}{v_\infty} + \frac{v_\infty}{v_0}, & v_0 v_\infty \nu_1 + \nu_t &= v_0 x_3 + v_\infty x_2, \\ L_5 : \quad & x_1 = v_t v_1 + \frac{1}{v_t v_1}, & v_t v_1 x_2 + x_3 &= v_t \nu_0 + v_1 \nu_\infty, \\ L_6 : \quad & x_1 = v_t v_1 + \frac{1}{v_t v_1}, & v_t v_1 x_3 + x_2 &= v_t \nu_\infty + v_1 \nu_0, \\ L_7 : \quad & x_1 = \frac{v_t}{v_1} + \frac{v_1}{v_t}, & v_t v_1 \nu_0 + \nu_\infty &= v_t x_2 + v_1 x_3, \\ L_8 : \quad & x_1 = \frac{v_t}{v_1} + \frac{v_1}{v_t}, & v_t v_1 \nu_\infty + \nu_0 &= v_t x_3 + v_1 x_2. \end{aligned}$$

Each of these intersects with L_1^∞ at infinity. The next eight lines are given by

$$\begin{aligned} L_9 : \quad & x_2 = v_0 v_1 + \frac{1}{v_0 v_1}, & v_0 v_1 x_1 + x_3 &= v_0 \nu_t + v_1 \nu_\infty, \\ L_{10} : \quad & x_2 = v_0 v_1 + \frac{1}{v_0 v_1}, & v_0 v_1 x_3 + x_1 &= v_0 \nu_\infty + v_1 \nu_t, \end{aligned}$$

$$\begin{array}{ll}
 L_{11} : & x_2 = \frac{v_0}{v_1} + \frac{v_1}{v_0}, & v_0 v_1 \nu_t + \nu_\infty = v_0 x_1 + v_1 x_3, \\
 L_{12} : & x_2 = \frac{v_0}{v_1} + \frac{v_1}{v_0}, & v_0 v_1 \nu_\infty + \nu_t = v_0 x_3 + v_1 x_1, \\
 L_{13} : & x_2 = v_t v_\infty + \frac{1}{v_t v_\infty}, & v_t v_\infty x_1 + x_3 = v_t \nu_0 + v_\infty \nu_1, \\
 L_{14} : & x_2 = v_t v_\infty + \frac{1}{v_t v_\infty}, & v_t v_\infty x_3 + x_1 = v_t \nu_1 + v_\infty \nu_0, \\
 L_{15} : & x_2 = \frac{v_t}{v_\infty} + \frac{v_\infty}{v_t}, & v_t v_\infty \nu_0 + \nu_1 = v_t x_1 + v_\infty x_3, \\
 L_{16} : & x_2 = \frac{v_t}{v_\infty} + \frac{v_\infty}{v_t}, & v_t v_\infty \nu_1 + \nu_0 = v_t x_3 + v_\infty x_1.
 \end{array}$$

Each of these intersects with L_2^∞ at infinity. The final eight lines are given by

$$\begin{array}{ll}
 L_{17} : & x_3 = v_0 v_t + \frac{1}{v_0 v_t}, & v_0 v_t x_1 + x_2 = v_0 \nu_1 + v_t \nu_\infty, \\
 L_{18} : & x_3 = v_0 v_t + \frac{1}{v_0 v_t}, & v_0 v_t x_2 + x_1 = v_0 \nu_\infty + v_t \nu_1, \\
 L_{19} : & x_3 = \frac{v_0}{v_t} + \frac{v_t}{v_0}, & v_0 v_t \nu_1 + \nu_\infty = v_0 x_1 + v_t x_2, \\
 L_{20} : & x_3 = \frac{v_0}{v_t} + \frac{v_t}{v_0}, & v_0 v_t \nu_\infty + \nu_1 = v_0 x_2 + v_t x_1, \\
 L_{21} : & x_3 = v_1 v_\infty + \frac{1}{v_1 v_\infty}, & v_1 v_\infty x_1 + x_2 = v_1 \nu_0 + v_\infty \nu_t, \\
 L_{22} : & x_3 = v_1 v_\infty + \frac{1}{v_1 v_\infty}, & v_1 v_\infty x_2 + x_1 = v_1 \nu_t + v_\infty \nu_0, \\
 L_{23} : & x_3 = \frac{v_1}{v_\infty} + \frac{v_\infty}{v_1}, & v_1 v_\infty \nu_0 + \nu_t = v_1 x_1 + v_\infty x_2, \\
 L_{24} : & x_3 = \frac{v_1}{v_\infty} + \frac{v_\infty}{v_1}, & v_1 v_\infty \nu_t + \nu_0 = v_1 x_2 + v_\infty x_1.
 \end{array}$$

Each of these intersects with L_3^∞ at infinity.

In Figure 3.1, the intersection graph of the above 24 lines is given. Each line L_k intersects with one line at infinity and 9 further lines. We proceed to describe these intersections in more detail. Consider one of the first eight lines, L_i , $1 \leq i \leq 8$. This line intersects with precisely one other line among the first eight lines, given by L_{i-1} if i is even and L_{i+1} if i is odd. Furthermore, for any $5 \leq j \leq 12$, the line L_i intersects either L_{2j} or L_{2j-1} . This yields a total of $1 + 8 = 9$ affine intersection points of L_i with other lines. Analogous accounting holds for any line in the remaining two sets of lines. In particular, there are a total of $\frac{1}{2} \times 24 \times 9 = 108$ affine intersection points, corresponding to the 108 edges in Figure 3.1. Adding to this the further 27 intersection points at infinity, yields a total of 135 intersection points among lines in $\overline{\mathcal{X}}$.




FIGURE 3.1. Intersection graph of the 24 affine lines on the Jimbo-Fricke surface. The vertex with index k represents the line L_k , for $1 \leq k \leq 24$, as defined in Section 3.2.1. An edge represents an intersection point of the lines corresponding to its endpoints. The subgraph in red, with vertices k , $9 \leq k \leq 24$, is the Clebsch graph which similarly encodes intersections among the lines $L_k^{\mathcal{Y}}$ on the Segre surface \mathcal{Y} , see Section 3.2.2.

3.2.2. *Blowing down the monodromy manifold.* Blowing down any of the lines on the cubic $\overline{\mathcal{X}}$ yields a corresponding Segre (quartic) surface. Generally, different choices of lines lead to non-isomorphic Segre surfaces, since the automorphism group of a generic (projective) cubic is trivial. As we will see, blowing down the line $L_1^\infty \subseteq \overline{\mathcal{X}}$, leads to a Segre surface isomorphic to $\overline{\mathcal{Z}}_1$ coming from qP_{VI} as $q \uparrow 1$. To realise this blow down, we introduce coordinates

$$y_1 = x_1, \quad y_2 = x_2, \quad y_3 = x_3, \quad y_4 = x_2x_3,$$

which satisfy

$$y_2y_3 - y_4 = 0, \tag{3.26a}$$

$$y_1y_4 + y_1^2 + y_2^2 + y_3^2 + \omega_1y_2 + \omega_2y_2 + \omega_3y_3 + \omega_4 = 0. \tag{3.26b}$$

These two equations define an affine Segre surface, which we denote by \mathcal{Y} . Using projective coordinates

$$[Y_0 : Y_1 : Y_2 : Y_3 : Y_4] = [1 : y_1 : y_2 : y_3 : y_4],$$

its canonical projective completion $\overline{\mathcal{Y}}$ in \mathbb{P}^4 , is given by

$$Y_2Y_3 - Y_0Y_4 = 0, \quad (3.27a)$$

$$Y_1Y_4 + Y_1^2 + Y_2^2 + Y_3^2 + (\omega_1Y_1 + \omega_2Y_2 + \omega_3Y_3)Y_0 + \omega_4Y_0^2 = 0. \quad (3.27b)$$

The mapping $x \mapsto y$ is bi-rational, it maps \mathcal{X} biholomorphically onto \mathcal{Y} , projects L_1^∞ onto a point, and maps L_2^∞ and L_3^∞ to two conics which make up $\overline{\mathcal{Y}} \setminus \mathcal{Y}$, explicitly,

$$\pi([0 : 0 : X_2 : X_3]) = [0 : 0 : 0 : 0 : 1], \quad (3.28)$$

$$\pi([0 : X_1 : 0 : X_3]) = [0 : X_1^2 : 0 : X_1X_3 : -(X_1^2 + X_3^2)], \quad (3.29)$$

$$\pi([0 : X_1 : X_2 : 0]) = [0 : X_1^2 : X_1X_2 : 0 : -(X_1^2 + X_2^2)]. \quad (3.30)$$

It follows that

$$\begin{aligned} \pi : \overline{\mathcal{X}} &\rightarrow \overline{\mathcal{Y}} \\ x &\mapsto y \end{aligned} \quad (3.31)$$

is indeed a blow-down of the line $L_1^\infty \subseteq \overline{\mathcal{X}}$.

Note that the two conics, traced out by (3.29) and (3.30), are described by the following equations respectively,

$$Y_0 = 0, \quad Y_2 = 0, \quad Y_1Y_4 + Y_1^2 + Y_3^2 = 0, \quad \text{and} \quad (3.32a)$$

$$Y_0 = 0, \quad Y_3 = 0, \quad Y_1Y_4 + Y_1^2 + Y_2^2 = 0. \quad (3.32b)$$

In particular, they are irreducible and intersect only at the point defined in equation (3.28) and the point $Y = [0 : -1 : 0 : 0 : 1]$.

Under the blow-down, each of the lines L_k , $1 \leq k \leq 8$, is also mapped to a conic. For example, $\pi(L_1)$ is given by

$$y_1 = v_0v_\infty + \frac{1}{v_0v_\infty}, \quad v_0v_\infty y_2 + y_3 = v_0\nu_t + v_\infty\nu_1, \quad y_4 = y_2y_3.$$

On the other hand, each of the lines L_k , $9 \leq k \leq 24$, is mapped to a line on the Segre surface. For instance, $\pi(L_9)$ reads

$$y_2 = v_0v_1 + \frac{1}{v_0v_1}, \quad v_0v_1y_1 + y_3 = v_0\nu_t + v_1\nu_\infty, \quad y_4 = \left(v_0v_1 + \frac{1}{v_0v_1}\right)y_3.$$

In other words, those lines that intersect with L_1^∞ become conics, and, those that do not, remain lines under the blow-down. Moreover, these are all the lines on the Segre surface. To see this, note that any line L on $\overline{\mathcal{Y}}$ cannot lie inside the curve at infinity, $\overline{\mathcal{Y}} \setminus \mathcal{Y}$, as both conics at infinity are irreducible. Its affine part thus admits a parametrisation,

$$L : \quad y = ta + b \quad (t \in \mathbb{C}),$$

for some $a, b \in \mathbb{C}^4$, with at least one $a_k \neq 0$, $1 \leq k \leq 3$, since $y_4 = y_2y_3$. It follows that

$$\pi^{-1}(L) = \{[t_0 : a_1t_1 + b_1t_0 : a_2t_1 + b_2t_0 : a_3t_1 + b_3t_0] : [t_0 : t_1] \in \mathbb{P}^1\},$$

is a line on the cubic.

So, the images

$$L_k^{\mathcal{Y}} := \pi(L_k) \quad (9 \leq k \leq 24), \quad (3.33)$$

are all the lines on $\overline{\mathcal{Y}}$, and their intersection graph is given by the red subgraph in Figure 3.1, which is the Clebsch graph. In particular, this showcases the classical fact that a smooth Segre surface contains exactly 16 lines [48].

3.3. Finding an isomorphism. We have constructed two affine Segre surfaces naturally associated with Painlevé VI. The first, denoted by \mathcal{Z}_1 , is the result of the continuum limit $q \uparrow 1$ of the monodromy manifold \mathcal{Z}_q for qP_{VI} . The second, denoted by \mathcal{Y} , was constructed by blowing down one of the lines at infinity of the Jimbo-Fricke cubic.

It is natural to ask whether these two Segre surfaces are affinely equivalent. We answer this question in the positive in Theorem 3.8.

A brute force way to construct such an affine equivalence explicitly, would be to use the intersections of the 16 lines in \mathcal{Z}_1 and in \mathcal{Y} . Any affine equivalence induces an isomorphism between the Clebsch graph in Figure 2.1 and the Clebsch subgraph in Figure 3.1. So, one could loop through all such isomorphisms and, for each, check whether it is compatible with an affine mapping from \mathbb{C}^6 to \mathbb{C}^4 , using explicit expressions for some of the intersection points. If so, then one can finally check whether it maps \mathcal{Z}_1 onto \mathcal{Y} . However, this approach would be computationally heavy, devoid of leveraging any existing knowledge within the theory of the Painlevé equations.

Our strategy instead, is to observe that the Tyurin ratios defined in (2.32) appear in the asymptotic expansions for solutions of qP_{VI} , see [42]. This is very interesting because, on one side the Tyurin ratios are natural globally defined rational function on the Segre surface $\overline{\mathcal{Z}}_q$, on the other side, their limit as $q \uparrow 1$ can be related to the asymptotics of P_{VI} , which are expressed in terms of the x_1, x_2, x_3 associated to the monodromy manifold.

To make the last step concrete, we will consider asymptotics of solutions to qP_{VI} and P_{VI} on the sheet $\mathbb{C} \setminus \mathbb{R}_{\geq 0}$. We thus pick a $t_0 \in \mathbb{C} \setminus \mathbb{R}_{\geq 0}$ and assume $0 < q < 1$. We start with explicit formulas for the asymptotics of $f(t)$, $t \in q^{\mathbb{Z}}t_0$, as $t \rightarrow 0$ and $t \rightarrow \infty$ in terms of $z \in \mathcal{Z}_q$ under the Riemann-Hilbert correspondence. For this purpose, the following triple of Tyurin ratios will be very helpful,

$$g^{(q)} = (g_1^{(q)}, g_2^{(q)}, g_3^{(q)}), \quad g_1^{(q)} := T_{62}^{(q)}, \quad g_2^{(q)} := T_{36}^{(q)}, \quad g_3^{(q)} := T_{64}^{(q)}. \quad (3.34)$$

We recall that these are globally defined rational functions of $z \in \mathcal{Z}_q$. All the asymptotic formulas will be expressed in terms of these three Tyurin ratios.

For generic monodromy data $z \in \mathcal{Z}_q$, the following asymptotic results were obtained in [42].

Define exponents σ_2 and σ_3 , each with $0 < \Re\sigma < \frac{1}{2}$, through

$$g_2^{(q)} = \frac{\theta_q(q^{\sigma_2 - \vartheta_t + \vartheta_\infty}, q^{\sigma_2 + \vartheta_t - \vartheta_\infty})}{\theta_q(q^{\sigma_2 + \vartheta_t + \vartheta_\infty}, q^{\sigma_2 - \vartheta_t - \vartheta_\infty})}, \quad g_3^{(q)} = \frac{\theta_q(q^{\sigma_3 - \vartheta_1 + \vartheta_\infty}, q^{\sigma_3 + \vartheta_1 - \vartheta_\infty})}{\theta_q(q^{\sigma_3 + \vartheta_1 + \vartheta_\infty}, q^{\sigma_3 - \vartheta_1 - \vartheta_\infty})}. \quad (3.35)$$

The leading order asymptotic behaviour of $f(t)$ as $t \rightarrow 0$, is given by

$$f(t) \sim q^{-\vartheta_t} \frac{(q^{\vartheta_t + \vartheta_0 + \sigma_3} - 1)(q^{\vartheta_t - \vartheta_0 + \sigma_3} - 1)(q^{\vartheta_1 + \vartheta_\infty + \sigma_3} - 1)}{(q^{\vartheta_1 + \vartheta_\infty - \sigma_3} - 1)(q^{\sigma_3} - q^{-\sigma_3})^2} \frac{1}{cs} (-t)^{1-2\sigma_3},$$

with the branch of $(-t)^{1-2\sigma_3}$ principle and c and s given by

$$c = \frac{\Gamma_q(1-2\sigma_3)^2}{\Gamma_q(1+2\sigma_3)^2} \prod_{\epsilon=\pm 1} \frac{\Gamma_q(1+\vartheta_t+\epsilon\vartheta_0+\sigma_3)\Gamma_q(1+\vartheta_1+\epsilon\vartheta_\infty+\sigma_3)}{\Gamma_q(1+\vartheta_t+\epsilon\vartheta_0-\sigma_3)\Gamma_q(1+\vartheta_1+\epsilon\vartheta_\infty-\sigma_3)},$$

$$s = -(-t_0)^{-2\sigma_3} M(g_1^{(q)}),$$

where $M(\cdot)$ is the Möbius transformation

$$M(G) = \frac{\theta_q(q^{\vartheta_1 - \vartheta_\infty + \sigma_3})\theta_q(q^{\vartheta_t + \vartheta_\infty + \sigma_3}t_0^{-1}) - G\theta_q(q^{\vartheta_1 + \vartheta_\infty + \sigma_3})\theta_q(q^{\vartheta_t - \vartheta_\infty + \sigma_3}t_0^{-1})}{\theta_q(q^{\vartheta_1 - \vartheta_\infty - \sigma_3})\theta_q(q^{\vartheta_t + \vartheta_\infty - \sigma_3}t_0^{-1}) - G\theta_q(q^{\vartheta_1 + \vartheta_\infty - \sigma_3})\theta_q(q^{\vartheta_t - \vartheta_\infty - \sigma_3}t_0^{-1})}.$$

We will refer to s as the twist parameter.

The leading order asymptotic behaviour of $f(t)$ as $t \rightarrow \infty$, is given by

$$f(t) \sim F_\infty (-1/t)^{-2\sigma_2}, \quad (3.36)$$

for a nonzero multiplier $F_\infty = F_\infty(q, \sigma_2, g_1)$, whose explicit expression is given in [42] but will not be required for our purposes, since we only need the value of the exponent σ_2 .

Next, we will consider the continuum limit of these asymptotic formulas, and compare them with Jimbo's asymptotic formulas for P_{VI} [24].

As $q \uparrow 1$, a formal computation yields that $f(t)$ converges to a solution $u(t)$, as explained at the beginning of Section 3. To take formal limits of the asymptotic formulas above, we use the limit laws in equation (3.2) as well as

$$\lim_{q \uparrow 1} g_1^{(q)} = (-t_0)^{2\vartheta_\infty} g_1^{(1)}, \quad \lim_{q \uparrow 1} g_2^{(q)} = g_2^{(1)}, \quad \lim_{q \uparrow 1} g_3^{(q)} = g_3^{(1)},$$

where the discontinuity in the first expression is a consequence of the discontinuity of $\eta_2^{(q)}$ at $q = 1$, see equation (3.15). A direct formal computation now yields

$$u(t) \sim \frac{(\vartheta_0 + \vartheta_t + \sigma_3)(-\vartheta_0 + \vartheta_t + \sigma_3)(\vartheta_\infty + \vartheta_1 + \sigma_3)}{4(\vartheta_\infty + \vartheta_1 - \sigma_3)\sigma_3^2} \frac{1}{cs} (-t)^{1-2\sigma_3},$$

as $t \rightarrow 0$, where

$$c = \frac{\Gamma(1 - 2\sigma_3)^2}{\Gamma(1 + 2\sigma_3)^2} \prod_{\epsilon = \pm 1} \frac{\Gamma(1 + \vartheta_t + \epsilon\vartheta_0 + \sigma_3)\Gamma(1 + \vartheta_1 + \epsilon\vartheta_\infty + \sigma_3)}{\Gamma(1 + \vartheta_t + \epsilon\vartheta_0 - \sigma_3)\Gamma(1 + \vartheta_1 + \epsilon\vartheta_\infty - \sigma_3)},$$

$$s = \frac{\sin(\pi(-\vartheta_\infty + \vartheta_1 + \sigma_3)) - g_1^{(1)} \sin(\pi(+\vartheta_\infty + \vartheta_1 + \sigma_3))}{\sin(\pi(+\vartheta_\infty - \vartheta_1 + \sigma_3)) - g_1^{(1)} \sin(\pi(-\vartheta_\infty - \vartheta_1 + \sigma_3))}. \quad (3.37)$$

At the same time, the asymptotic exponent at $t = \infty$ in (3.36) is unchanged under the limit $q \uparrow 1$. Furthermore, note that, as $q \uparrow 1$, equations (3.35) become

$$g_2^{(1)} = \frac{\sin(\pi(+\vartheta_\infty - \vartheta_t + \sigma_2)) \sin(\pi(-\vartheta_\infty + \vartheta_t + \sigma_2))}{\sin(\pi(+\vartheta_\infty + \vartheta_t + \sigma_2)) \sin(\pi(-\vartheta_\infty - \vartheta_t + \sigma_2))}, \quad (3.38)$$

$$g_3^{(1)} = \frac{\sin(\pi(+\vartheta_\infty - \vartheta_1 + \sigma_3)) \sin(\pi(-\vartheta_\infty + \vartheta_1 + \sigma_3))}{\sin(\pi(+\vartheta_\infty + \vartheta_1 + \sigma_3)) \sin(\pi(-\vartheta_\infty - \vartheta_1 + \sigma_3))}.$$

We now recall Jimbo's asymptotic formulas [24], which relate the exponents σ_2 and σ_3 at $t = 0$ and $t = \infty$, as well as the value of twist parameter s , to the coordinates (x_1, x_2, x_3) of the Jimbo-Fricke cubic (3.22).

Firstly, the exponents are given by

$$x_2 = 2 \cos(2\pi\sigma_2), \quad x_3 = 2 \cos(2\pi\sigma_3). \quad (3.39)$$

The explicit formula for the twist parameter s is given in [24, eq. 1.8]¹, and translates to the following in our notation,

$$s = \frac{a + b e^{2\pi i \sigma_3}}{d}, \quad (3.40)$$

$$a = \frac{1}{2} i \sin(2\pi\sigma_3) x_1 + \cos(2\pi\vartheta_0) \cos(2\pi\vartheta_1) + \cos(2\pi\vartheta_t) \cos(2\pi\vartheta_\infty),$$

$$b = \frac{1}{2} i \sin(2\pi\sigma_3) x_2 - \cos(2\pi\vartheta_t) \cos(2\pi\vartheta_1) - \cos(2\pi\vartheta_0) \cos(2\pi\vartheta_\infty),$$

$$d = 4 \prod_{\epsilon = \pm 1} \sin[\pi(\vartheta_0 + \epsilon(\vartheta_t - \sigma_3))] \sin[\pi(\vartheta_\infty + \epsilon(\vartheta_1 - \sigma_3))].$$

To obtain these formulas, we note the following correspondence between the notation in Jimbo [24] and ours,

$$s^{[J]} = -e^{-2\pi i \sigma_3} s,$$

¹We note the following typo in [24]: the last '≠' at the bottom of page 1141 should have been a \pm , as pointed out in [20].

$$\begin{aligned}\vartheta_k^{[J]} &= 2\vartheta_k \quad (k = 0, t, 1, \infty), \\ \sigma_{1t}^{[J]} &= 2\sigma_1, \quad \sigma_{01}^{[J]} = 2\sigma'_2, \quad \sigma^{[J]} = 2\sigma_3,\end{aligned}$$

with

$$2 \cos(2\pi\sigma_1) = x_1, \quad 2 \cos(2\pi\sigma'_2) = x'_2 = -x_2 - x_1x_3 + \nu_0\nu_1 + \nu_t\nu_\infty,$$

where we recall that $\nu_k = 2 \cos(2\pi\vartheta_k)$. The transformation $x_2 \mapsto x'_2$ is an element of the standard extended modular group action [23] on the Jimbo-Fricke cubic. Its application was necessary to obtain the correct result on the sheet $\mathbb{C} \setminus \mathbb{R}_{\geq 0}$, since Jimbo's original formulas are with respect to the sheet $\mathbb{C} \setminus ((-\infty, 0] \cup [1, +\infty))$.

Now, we are in a position to compare Jimbo's asymptotics formulas with those obtained from the continuum limit $q \uparrow 1$.

Proposition 3.6. *Consistency of equations (3.22), (3.37), (3.38), (3.39) and (3.40) implies*

$$g_2^{(1)} = \frac{x_2 - (v_t v_\infty^{-1} + v_t^{-1} v_\infty)}{x_2 - (v_t v_\infty + v_t^{-1} v_\infty^{-1})}, \quad g_3^{(1)} = \frac{x_3 - (v_1 v_\infty^{-1} + v_1^{-1} v_\infty)}{x_3 - (v_1 v_\infty + v_1^{-1} v_\infty^{-1})}, \quad (3.41)$$

and

$$\begin{aligned}g_1^{(1)} &= v_\infty \frac{x_1 - v_1 v_\infty^{-1} x_2 - v_t^{-1} v_\infty^{-1} x_3 + x_2 x_3 - \nu_0 v_\infty^{+1} - v_t v_1^{-1} + v_t^{-1} v_1 v_\infty^{-2}}{x_1 - v_1 v_\infty^{+1} x_2 - v_t^{-1} v_\infty^{+1} x_3 + x_2 x_3 - \nu_0 v_\infty^{-1} - v_t v_1^{-1} + v_t^{-1} v_1 v_\infty^{+2}} \\ &= v_\infty^{-1} \frac{x_1 + v_1^{-1} v_\infty^{+1} x_2 + v_t v_\infty^{+1} x_3 - \nu_0 v_\infty^{+1} - \nu_1 v_t - v_t^{-1} v_1 v_\infty^{+2}}{x_1 + v_1^{-1} v_\infty^{-1} x_2 + v_t v_\infty^{-1} x_3 - \nu_0 v_\infty^{-1} - \nu_1 v_t - v_t^{-1} v_1 v_\infty^{-2}},\end{aligned}$$

where we remind the reader of the definition of ν_k and v_k , $k = 0, t, 1, \infty$, in equation (3.24).

Proof. Let us start with deriving equations (3.41). By applying the familiar product to sum formula for the sine function to the numerator and denominator of the formula for $g_2^{(1)}$ in (3.38), we find

$$\begin{aligned}g_2^{(1)} &= \frac{\sin(\pi(+\vartheta_\infty - \vartheta_t + \sigma_2)) \sin(\pi(-\vartheta_\infty + \vartheta_t + \sigma_2))}{\sin(\pi(+\vartheta_\infty + \vartheta_t + \sigma_2)) \sin(\pi(-\vartheta_\infty - \vartheta_t + \sigma_2))} \\ &= \frac{\cos(2\pi\sigma_2) - \cos(2\pi(\vartheta_t - \vartheta_\infty))}{\cos(2\pi\sigma_2) - \cos(2\pi(\vartheta_t + \vartheta_\infty))} \\ &= \frac{x_2 - 2 \cos(2\pi(\vartheta_t - \vartheta_\infty))}{x_2 - 2 \cos(2\pi(\vartheta_t + \vartheta_\infty))} \\ &= \frac{x_2 - (v_t v_\infty^{-1} + v_t^{-1} v_\infty)}{x_2 - (v_t v_\infty + v_t^{-1} v_\infty^{-1})}.\end{aligned}$$

A similar computation leads to the formula for $g_3^{(1)}$ in (3.41).

To obtain a formula for $g_1^{(1)}$, we compare the two formulas for the twist parameter that we have, equations (3.37) and (3.40). Rather than dealing with trigonometric identities, we rewrite both formulas as rational functions in v_0, v_t, v_1, v_∞ and v_3 , where, by definition, $v_3 = e^{2\pi i \sigma_3}$. Equation (3.37) then reads

$$s = \frac{(v_3 v_1 - v_\infty) - g_1^{(1)}(v_3 v_1 v_\infty - 1)}{(v_3 v_\infty - v_1) - g_1^{(1)}(v_3 - v_1 v_\infty)}, \quad (3.42)$$

and equation (3.40) becomes

$$\begin{aligned}s &= \frac{a + b v_3}{d}, \\ a &= \frac{1}{4} ((v_3 - v_3^{-1})x_1 + (v_0 + v_0^{-1})(v_1 + v_1^{-1}) + (v_t + v_t^{-1})(v_\infty + v_\infty^{-1})),\end{aligned} \quad (3.43)$$

$$b = \frac{1}{4} \left((v_3 - v_3^{-1})x_2 + (v_0 + v_0^{-1})(v_\infty + v_\infty^{-1}) + (v_t + v_t^{-1})(v_1 + v_1^{-1}) \right),$$

$$d = \frac{(v_t - v_0v_3)(v_3 - v_0v_t)(v_1 - v_3v_\infty)(v_3 - v_1v_\infty)}{4v_0v_tv_1v_\infty v_3^2}.$$

Now, we want to point out a symmetry that will be important in what follows. If we take either formula for the twist parameter s , and substitute v_3 for its reciprocal, then this is equivalent to taking the reciprocal of s , in other words,

$$s|_{v_3 \mapsto v_3^{-1}} = s^{-1}. \quad (3.44)$$

In formula (3.42), this fact is obvious. In formula (3.43) it is not that transparent and a direct computation shows that it is in fact equivalent to the cubic relation (3.22) among x_1, x_2 and $x_3 = v_3 + v_3^{-1}$. We come back to this symmetry in a moment.

By equating the right-hand sides of equations (3.42) and (3.43) and multiplying out the denominators, we obtain a polynomial equation, of degree 1 in the variables $g_1^{(1)}$, x_2 and x_3 , and of degree 3 in v_3 . Considering this as a polynomial equation just in v_3 , we can eliminate the constant term and highest order term, by replacing respectively $1 \mapsto v_3x_3 - v_3^2$ and $v_3^3 \rightarrow v_3^2x_3 - v_3$. Dividing the result by v_3 , we obtain a polynomial equation in v_3 of degree one,

$$v_1v_\infty^{-1}(A_+ - v_\infty g_1^{(1)}A_-) - v_3(B_+ - v_\infty^{-1}g_1^{(1)}B_-) = 0, \quad (3.45)$$

where

$$A_\pm = x_1 - v_1v_\infty^{\mp 1}x_2 - v_t^{-1}v_\infty^{\mp 1}x_3 + x_2x_3 - \nu_0v_\infty^{\pm 1} - v_tv_1^{-1} + v_t^{-1}v_1v_\infty^{\mp 2},$$

$$B_\pm = x_1 + v_1^{-1}v_\infty^{\pm 1}x_2 + v_tv_\infty^{\pm 1}x_3 - \nu_0v_\infty^{\pm 1} - \nu_1v_t - v_t^{-1}v_1v_\infty^{\pm 2}.$$

By symmetry (3.44), both equation (3.45) and the same equation with v_3 replaced by v_3^{-1} hold true simultaneously. This implies

$$(A_+ - v_\infty g_1^{(1)}A_-) = 0, \quad (B_+ - v_\infty^{-1}g_1^{(1)}B_-) = 0,$$

and therefore

$$g_1^{(1)} = v_\infty \frac{A_+}{A_-} = v_\infty^{-1} \frac{B_+}{B_-},$$

which are the two formulas for $g_1^{(1)}$ in the proposition, and concludes the proof. \square

Remark 3.7. In Proposition 3.6, two different formulas are given for $g_1^{(1)}$. Their equality is equivalent to the Jimbo-Fricke cubic equation.

3.4. An isomorphism. In Section 3.3, we found explicit formulas relating the z -variables on the Segre surface \mathcal{Z}_1 , with the x -variables on the Jimbo-Fricke cubic \mathcal{X} , through the intermediate Tyurin ratios $g^{(1)}$, see in particular Proposition 3.6. These formulas lead to the isomorphism in the following theorem, which is the focus of this section.

Theorem 3.8. *The Jimbo-Fricke cubic surface \mathcal{X} and the Segre surface \mathcal{Z}_1 , defined in Definition 3.3, are isomorphic as affine varieties. An explicit isomorphism is given by the polynomial mapping*

$$\Phi_{\mathcal{X}} : \begin{array}{ccc} \mathcal{X} & \rightarrow & \mathcal{Z}_1 \\ x & \mapsto & z \end{array}$$

where

$$z_1 = +\gamma^{-1} \left(v_\infty \left(x_2 - v_\infty^{-1}\nu_t \right) \left(x_3 - v_\infty^{-1}\nu_1 \right) + (v_\infty - v_\infty^{-1}) \left(x_1 - v_\infty\nu_0 \right) \right),$$

$$z_2 = +\gamma^{-1} \left(v_\infty^{-1} \left(x_2 - v_\infty\nu_t \right) \left(x_3 - v_\infty\nu_1 \right) - (v_\infty - v_\infty^{-1}) \left(x_1 - v_\infty^{-1}\nu_0 \right) \right),$$

$$\begin{aligned}
z_3 &= -\delta^{-1} \left(v_0 - v_t v_1 v_\infty \right) \left(v_0 - \frac{1}{v_t v_1 v_\infty} \right) \left(x_2 - \frac{v_t}{v_\infty} - \frac{v_\infty}{v_t} \right) \left(x_3 - \frac{v_1}{v_\infty} - \frac{v_\infty}{v_1} \right), \\
z_4 &= -\delta^{-1} \left(v_0 - \frac{v_t v_1}{v_\infty} \right) \left(v_0 - \frac{v_\infty}{v_t v_1} \right) \left(x_2 - v_t v_\infty - \frac{1}{v_t v_\infty} \right) \left(x_3 - v_1 v_\infty - \frac{1}{v_1 v_\infty} \right), \\
z_5 &= +\delta^{-1} \left(v_0 - \frac{v_t v_\infty}{v_1} \right) \left(v_0 - \frac{v_1}{v_t v_\infty} \right) \left(x_2 - \frac{v_t}{v_\infty} - \frac{v_\infty}{v_t} \right) \left(x_3 - v_1 v_\infty - \frac{1}{v_1 v_\infty} \right), \\
z_6 &= +\delta^{-1} \left(v_0 - \frac{v_1 v_\infty}{v_t} \right) \left(v_0 - \frac{v_t}{v_1 v_\infty} \right) \left(x_2 - v_t v_\infty - \frac{1}{v_t v_\infty} \right) \left(x_3 - \frac{v_1}{v_\infty} - \frac{v_\infty}{v_1} \right),
\end{aligned}$$

with the constants γ and δ defined by

$$\begin{aligned}
\gamma &= (v_0 - 1)(v_0^{-1} - 1)(v_\infty - v_\infty^{-1})^2, \\
\delta &= (v_0 - 1)^2(v_t - v_t^{-1})(v_1 - v_1^{-1})(v_\infty - v_\infty^{-1})^2.
\end{aligned}$$

Composition of this isomorphism with the inverse of the blow-down mapping π restricted to \mathcal{Y} ,

$$\Phi_{\mathcal{Y}} := \Phi_{\mathcal{X}} \circ (\pi^{-1}|_{\mathcal{Y}}),$$

defines an affine equivalence

$$\begin{aligned}
\Phi_{\mathcal{Y}} : \mathcal{Y} &\rightarrow \mathcal{Z}_1 \\
y &\mapsto z
\end{aligned}$$

between the Segre surfaces \mathcal{Y} and \mathcal{Z}_1 and, in particular, extends to a projective equivalence between their completions $\overline{\mathcal{Y}}$ and $\overline{\mathcal{Z}}_1$.

Remark 3.9. Note that the expressions for the polynomial mapping $\Phi_{\mathcal{X}}$ in Theorem 3.8 are affine linear in $\{x_1, x_2, x_3, x_2 x_3\}$. Indeed, they can be written as

$$z_k = \xi_{0k} + \xi_{1k} x_1 + \xi_{2k} x_2 + \xi_{3k} x_3 + \xi_{4k} x_2 x_3, \quad 1 \leq k \leq 6,$$

for some coefficients that can be read off directly from the formulas in the theorem. Consequently, $\Phi_{\mathcal{Y}}$ can be written as

$$z_k = \xi_{0k} + \xi_{1k} y_1 + \xi_{2k} y_2 + \xi_{3k} y_3 + \xi_{4k} y_4, \quad 1 \leq k \leq 6, \quad (3.46)$$

and extends to an affine linear map from \mathbb{C}^4 to \mathbb{C}^6 that maps \mathcal{Y} to \mathcal{Z}_1 , as stated in the theorem.

Remark 3.10. The affine equivalence $\Phi_{\mathcal{Y}}$ in Theorem 3.8 maps the 16 lines in \mathcal{Y} , see equation (3.33), to the 16 lines in \mathcal{Z}_1 , see Section 3.1.1, as follows

$$\begin{aligned}
L_9^{\mathcal{Y}} &\mapsto \tilde{L}_{2,4,5}^{(1)}, & L_{10}^{\mathcal{Y}} &\mapsto \tilde{L}_{1,4,5}^{(1)}, & L_{11}^{\mathcal{Y}} &\mapsto \tilde{L}_{2,3,6}^{(1)}, & L_{12}^{\mathcal{Y}} &\mapsto \tilde{L}_{1,3,6}^{(1)}, \\
L_{13}^{\mathcal{Y}} &\mapsto L_{2,4,6}^{(1)}, & L_{14}^{\mathcal{Y}} &\mapsto L_{1,4,6}^{(1)}, & L_{15}^{\mathcal{Y}} &\mapsto L_{1,3,5}^{(1)}, & L_{16}^{\mathcal{Y}} &\mapsto L_{2,3,5}^{(1)}, \\
L_{17}^{\mathcal{Y}} &\mapsto \tilde{L}_{2,4,6}^{(1)}, & L_{18}^{\mathcal{Y}} &\mapsto \tilde{L}_{1,4,6}^{(1)}, & L_{19}^{\mathcal{Y}} &\mapsto \tilde{L}_{2,3,5}^{(1)}, & L_{20}^{\mathcal{Y}} &\mapsto \tilde{L}_{1,3,5}^{(1)}, \\
L_{21}^{\mathcal{Y}} &\mapsto L_{2,4,5}^{(1)}, & L_{22}^{\mathcal{Y}} &\mapsto L_{1,4,5}^{(1)}, & L_{23}^{\mathcal{Y}} &\mapsto L_{1,3,6}^{(1)}, & L_{24}^{\mathcal{Y}} &\mapsto L_{2,3,6}^{(1)}.
\end{aligned} \quad (3.47)$$

In particular, it induces an isomorphism between the Clebsch graph in Figure 2.1 and the Clebsch subgraph in Figure 3.1.

Remark 3.11. Recall that the curve at infinity $\overline{\mathcal{Y}} \setminus \mathcal{Y}$ factorises into two conics (3.32a) and (3.32b) and, similarly, the curve at infinity $\overline{\mathcal{Z}}_1 \setminus \mathcal{Z}_1$ factorises into two conics (3.20) and (3.21). The extension of $\Phi_{\mathcal{Y}}$ to a projective equivalence between $\overline{\mathcal{Y}}$ and $\overline{\mathcal{Z}}_1$ in Theorem 3.8, maps conics (3.32a) and (3.32b) respectively to conics (3.20) and (3.21).

This is consistent with Remark 3.10 in the following way. In Figure 2.1, lines in \mathcal{Z}_1 are coloured respectively ForestGreen or Fuchsia dependent on whether they intersect conic (3.20) or conic (3.21) at infinity. The lines $L_k^{\mathcal{Y}}$, $9 \leq k \leq 16$, which intersect with conic (3.32a) at infinity, are mapped to the ForestGreen lines, whilst

the lines $L_k^{\mathcal{Y}}$, $17 \leq k \leq 24$, which intersect with conic (3.32b) at infinity, are mapped to the Fuchsia ones.

Remark 3.12. A question is raised in Item 7.2.10 (b) of [40, §7.2], about the correspondence between the monodromy manifold of q - P_{VI} and that of P_{VI} based on the number of lines each contains. It is stated that “*It seems dubious that it can be translated into a birational map: the number of lines increases by confluence.*” (Note that the term confluence in this quote refers to the continuum limit.) But it is well known that one can increase the number of lines on an algebraic surface by allowing blow-ups. Theorem 3.8, shows that there is a birational map of a most simple kind involving a blow-up between these two monodromy manifolds.

We prove Theorem 3.8 in several steps and in a reverse order compared to the way the theorem is stated. Namely, we use the results in Section 3.3, to construct a bi-rational mapping between $\overline{\mathcal{Y}}$ and $\overline{\mathcal{Z}}_1$. We show that this mapping is a projective equivalence with the right properties, so that it induces an affine equivalence between \mathcal{Y} and \mathcal{Z}_1 and in the end leads to the isomorphism $\Phi_{\mathcal{X}}$. We further prove Remarks 3.10 and 3.11 on the way.

Our starting point for the proof of Theorem 3.8, is the set of explicit formulas obtained in Section 3.3 that relate the z -variables on \mathcal{Z}_1 with the x -variables on \mathcal{X} , through the intermediate Tyurin ratios $g_k^{(1)}$, $1 \leq k \leq 3$. We recall them here, but written with respect to homogeneous coordinates Z and Y on $\overline{\mathcal{Z}}_1$ and $\overline{\mathcal{Y}}$ respectively.

On the one hand, with respect to homogeneous coordinates for $\overline{\mathcal{Z}}_1$, we have

$$\begin{aligned} g_1^{(1)} &= \frac{Z_6/\eta_6^{(1)}}{Z_2/\eta_2^{(1)}} = \frac{Z_1/\eta_1^{(1)}}{Z_5/\eta_5^{(1)}}, \\ g_2^{(1)} &= \frac{Z_3/\eta_3^{(1)}}{Z_6/\eta_6^{(1)}} = \frac{Z_5/\eta_5^{(1)}}{Z_4/\eta_4^{(1)}}, \\ g_3^{(1)} &= \frac{Z_6/\eta_6^{(1)}}{Z_4/\eta_4^{(1)}} = \frac{Z_3/\eta_3^{(1)}}{Z_5/\eta_5^{(1)}}. \end{aligned} \quad (3.48)$$

On the other hand, with respect to homogeneous coordinates for $\overline{\mathcal{Y}}$, we have, see Proposition 3.6,

$$\begin{aligned} g_1^{(1)} &= v_{\infty} \frac{Y_1 - v_1 v_{\infty}^{-1} Y_2 - v_t^{-1} v_{\infty}^{-1} Y_3 + Y_4 - Y_0(\nu_0 v_{\infty}^{+1} + v_t v_1^{-1} - v_t^{-1} v_1 v_{\infty}^{-2})}{Y_1 - v_1 v_{\infty}^{+1} Y_2 - v_t^{-1} v_{\infty}^{+1} Y_3 + Y_4 - Y_0(\nu_0 v_{\infty}^{-1} + v_t v_1^{-1} - v_t^{-1} v_1 v_{\infty}^{+2})} \\ &= v_{\infty}^{-1} \frac{Y_1 + v_1^{-1} v_{\infty}^{+1} Y_2 + v_t v_{\infty}^{+1} Y_3 - Y_0(\nu_0 v_{\infty}^{+1} + \nu_1 v_t + v_t^{-1} v_1 v_{\infty}^{+2})}{Y_1 + v_1^{-1} v_{\infty}^{-1} Y_2 + v_t v_{\infty}^{-1} Y_3 - Y_0(\nu_0 v_{\infty}^{-1} + \nu_1 v_t + v_t^{-1} v_1 v_{\infty}^{-2})}. \\ g_2^{(1)} &= \frac{Y_2 - Y_0(v_t v_{\infty}^{-1} + v_t^{-1} v_{\infty})}{Y_2 - Y_0(v_t v_{\infty} + v_t^{-1} v_{\infty}^{-1})}, \\ g_3^{(1)} &= \frac{Y_3 - Y_0(v_1 v_{\infty}^{-1} + v_1^{-1} v_{\infty})}{Y_3 - Y_0(v_1 v_{\infty} + v_1^{-1} v_{\infty}^{-1})}. \end{aligned} \quad (3.49)$$

We correspondingly define the following mappings

$$g_{\mathcal{Z}} : \overline{\mathcal{Z}}_1 \rightarrow (\mathbb{P}^1)^3, \quad Z \mapsto g^{(1)}, \quad (3.50)$$

$$g_{\mathcal{Y}} : \overline{\mathcal{Y}} \rightarrow (\mathbb{P}^1)^3, \quad Y \mapsto g^{(1)}, \quad (3.51)$$

which we prove to be analytic mappings in the following lemma.

Lemma 3.13. *The mappings $g_{\mathcal{Z}}$ and $g_{\mathcal{Y}}$ are analytic, that is, each of their components defines a meromorphic function on the respective complex varieties $\overline{\mathcal{Z}}_1$ and $\overline{\mathcal{Y}}$.*

Proof. For the mapping g_Z this assertion is trivial, since its components are Tyurin ratios on \overline{Z}_1 , which we know to be meromorphic functions (see the argument in the beginning of Section 2.4).

Regarding g_Y , the components $g_2^{(1)}$ and $g_3^{(1)}$ are clearly meromorphic as they can be written as compositions of Möbius transforms with the respective projections $y \mapsto y_2$ and $y \mapsto y_3$ from \overline{Y} to \mathbb{P}^1 .

Regarding $g_1^{(1)}$, we have two formulas (3.49), which are equivalent on \overline{Y} . Now, if $g_1^{(1)}$ is not meromorphic at some point, then this point must be a common root of the numerators and denominators in both formulas. Each of those numerators and denominators is projective linear, and thus defines a hyperplane in \mathbb{P}^4 . Their intersection is a single point, given by

$$\begin{aligned} Y_0 &= v_t v_1 - v_t^{-1} v_1^{-1}, \\ Y_1 &= v_t^2 v_1^2 - v_t^{-2} v_1^{-2}, \\ Y_2 &= \nu_0(v_t - v_t^{-1}) + \nu_\infty(v_1 - v_1^{-1}), \\ Y_3 &= \nu_0(v_1 - v_1^{-1}) + \nu_\infty(v_t - v_t^{-1}), \\ Y_4 &= (v_t v_1 - v_t^{-1} v_1^{-1}) (\nu_0 \nu_\infty - (v_t - v_t^{-1})(v_1 - v_1^{-1})). \end{aligned}$$

This point does not lie on \overline{Y} for generic parameter values, as it for example does not satisfy $Y_0 Y_4 - Y_2 Y_3 = 0$. It follows that $g_1^{(1)}$ is also a meromorphic function on \overline{Y} , and the lemma follows. \square

Since \overline{Z}_1 is two-dimensional, the meromorphic functions $g_k^{(1)}$, $1 \leq k \leq 3$, cannot be independent. They are explicitly related by the following identity,

$$\eta_1^{(1)} g_2^{(1)} g_1^{(1)} + \eta_2^{(1)} g_3^{(1)} / g_1^{(1)} + \eta_3^{(1)} g_2^{(1)} g_3^{(1)} + \eta_4^{(1)} + \eta_5^{(1)} g_2^{(1)} + \eta_6^{(1)} g_3^{(1)} = 0. \quad (3.52)$$

To verify this identity, we take the left-hand side and first apply the multiplication rule (2.34), and then apply the symmetry $T_{ij}^{(1)} = T_{\alpha(j)\alpha(i)}^{(1)}$ a couple of times:

$$\begin{aligned} &\eta_1^{(1)} g_2^{(1)} g_1^{(1)} + \eta_2^{(1)} g_3^{(1)} / g_1^{(1)} + \eta_3^{(1)} g_2^{(1)} g_3^{(1)} + \eta_4^{(1)} + \eta_5^{(1)} g_2^{(1)} + \eta_6^{(1)} g_3^{(1)} = \\ &\eta_1^{(1)} T_{36}^{(1)} T_{62}^{(1)} + \eta_2^{(1)} T_{64}^{(1)} / T_{62}^{(1)} + \eta_3^{(1)} T_{36}^{(1)} T_{64}^{(1)} + \eta_4^{(1)} + \eta_5^{(1)} T_{36}^{(1)} + \eta_6^{(1)} T_{64}^{(1)} = \\ &\eta_1^{(1)} T_{32}^{(1)} + \eta_2^{(1)} T_{24}^{(1)} + \eta_3^{(1)} T_{34}^{(1)} + \eta_4^{(1)} + \eta_5^{(1)} T_{36}^{(1)} + \eta_6^{(1)} T_{64}^{(1)} = \\ &\eta_1^{(1)} T_{14}^{(1)} + \eta_2^{(1)} T_{24}^{(1)} + \eta_3^{(1)} T_{34}^{(1)} + \eta_4^{(1)} + \eta_5^{(1)} T_{54}^{(1)} + \eta_6^{(1)} T_{64}^{(1)} = \\ &\frac{\eta_4^{(1)}}{Z_4} (Z_1 + Z_2 + Z_3 + Z_4 + Z_5 + Z_6) = 0, \end{aligned}$$

where the last equality follows from (2.5a).

Equation (3.52) is an equality among meromorphic functions on \overline{Z}_1 . However, it also defines a surface.

Definition 3.14. Define $\overline{\mathcal{G}} \subseteq \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ as the topological closure of the surface defined by equation (3.52) in $\{(g_1, g_2, g_3) \in (\mathbb{C}^*)^3\}$. Also, set

$$\mathcal{G} := \overline{\mathcal{G}} \setminus \{g \in \overline{\mathcal{G}} : g_2 = 1 \text{ or } g_3 = 1\},$$

so that $\overline{\mathcal{G}}$ is the topological closure of \mathcal{G} .

Remark 3.15. Even though we defined the surface $\overline{\mathcal{G}}$ through topological closure, we note that it makes algebro-geometric sense. Namely, under the generalised Segre embedding

$$\begin{aligned} \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 &\rightarrow \mathbb{P}^7, \\ (g_1, g_2, g_3) &\mapsto U, \end{aligned}$$

where, in homogeneous coordinates $g_k = [g_k^a : g_k^b]$, $1 \leq k \leq 3$,

$$U = [U_{aaa} : U_{aab} : U_{aba} : U_{baa} : U_{abb} : U_{bab} : U_{bba} : U_{bbb}],$$

$$U_{s_1 s_2 s_3} = g_1^{s_1} g_2^{s_2} g_3^{s_3}, \quad (s_1, s_2, s_3 \in \{a, b\}),$$

the surface $\overline{\mathcal{G}}$ becomes a projective variety in \mathbb{P}^7 . This projective variety admits a cumbersome description involving 13 quadratic equations, 9 of which describe the image of $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ itself under the embedding. For our purposes, it is much more convenient to work with the analytic definition of $\overline{\mathcal{G}}$ in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$.

By a similar token, since $\overline{\mathcal{Y}}$ is two-dimensional, the meromorphic functions $g_k^{(1)}$, $1 \leq k \leq 3$, considered as functions of Y through equations (3.49), cannot be independent. Remarkably, they are related by the exact same formula, equation (3.52), as can be verified by direct computation.

For the next important preparatory step, for the proof of Theorem 3.8, we define three dense open subsets

$$U_{\mathcal{Z}} \subseteq \overline{\mathcal{Z}}_1, \quad U_{\mathcal{G}} \subseteq \overline{\mathcal{G}}, \quad U_{\mathcal{Y}} \subseteq \overline{\mathcal{Y}}.$$

The set $U_{\mathcal{Z}}$ is defined by the Segre surface $\overline{\mathcal{Z}}_1$, minus eight lines and the hyperplane section at infinity,

$$U_{\mathcal{Z}} = \mathcal{Z}_1 \setminus \bigcup_{(i,j,k) \in I} L_{i,j,k}^{(1)}, \quad I := \{1, 2\} \times \{3, 4\} \times \{5, 6\}, \quad (3.53)$$

where we recall the notation for lines on \mathcal{Z}_1 in Section 3.1.1. Similarly, $U_{\mathcal{Y}}$ is defined by the Segre surface $\overline{\mathcal{Y}}$, minus eight lines and the hyperplane section at infinity,

$$U_{\mathcal{Y}} = \mathcal{Y} \setminus \bigcup_{j \in J} L_j^{\mathcal{Y}}, \quad J := \{13, 14, 15, 16, 21, 22, 23, 24\},$$

where we recall the notation for lines on $\overline{\mathcal{Y}}$ introduced in equation (3.33).

Finally, $U_{\mathcal{G}}$ is simply defined as the intersection

$$U_{\mathcal{G}} = \mathcal{G} \cap (\mathbb{C}^* \times \mathbb{C}^* \times \mathbb{C}^*),$$

where we recall that \mathcal{G} is defined in Definition 3.14.

These dense open subsets are chosen exactly such that $g_1^{(1)}, g_2^{(1)}, g_3^{(1)}$ take finite, nonzero values on them. Regarding $U_{\mathcal{G}}$, this is tautological; regarding the two other open subsets, it follows from the following lemma.

Lemma 3.16. *The mapping $g_{\mathcal{Z}}$ maps $U_{\mathcal{Z}}$ biholomorphically onto $U_{\mathcal{G}}$. Similarly, $g_{\mathcal{Y}}$ maps $U_{\mathcal{Y}}$ biholomorphically onto $U_{\mathcal{G}}$.*

Proof. We start with the, due to Lemma 3.13, analytic mapping $g_{\mathcal{Z}}$. Note that, for any Tyurin ratio $T_{ij}^{(1)}$,

$$T_{ij}^{(1)} = 0 \iff Z_i = Z_{\alpha(j)} = 0, \text{ and}$$

$$T_{ij}^{(1)} = \infty \iff Z_j = Z_{\alpha(i)} = 0.$$

In particular, if $T^{(1)}(Z) \in \{0, \infty\}$, for some Tyurin ratio $T^{(1)}$, then Z necessarily lies on one of the eight lines $L_{i,j,k}^{(1)}$, $(i, j, k) \in I$. By construction, those lines do not intersect with $U_{\mathcal{Z}}$. Since all three components of $g_{\mathcal{Z}}$ are Tyurin ratios, it follows that $g_{\mathcal{Z}}(U_{\mathcal{Z}}) \subseteq (\mathbb{C}^*)^3$.

Now, take any point $z \in U_{\mathcal{Z}}$ and let $g^{(1)} = g_{\mathcal{Z}}(z) \in (\mathbb{C}^*)^3$, then the cubic equation (3.52) is satisfied by $g^{(1)}$. To infer that $g^{(1)} \in U_{\mathcal{G}}$, it remains to be checked that neither $g_2^{(1)} = 1$ nor $g_3^{(1)} = 1$.

Suppose, on the contrary, that $g_2^{(1)} = 1$. Then

$$Z_6 = \frac{\eta_6^{(1)}}{\eta_3^{(1)}} Z_3, \quad Z_5 = \frac{\eta_5^{(1)}}{\eta_4^{(1)}} Z_4. \quad (3.54)$$

We now consider the difference of the first two defining equations, (2.5a) and (2.5b), of the Segre surface \overline{Z}_1 , given by

$$(\mu_3 - 1)Z_3 + (\mu_4 - 1)Z_4 + (\mu_5 - 1)Z_5 + (\mu_6 - 1)Z_6 = Z_0.$$

Direct substitution of (3.54) and the equation for μ_k , $3 \leq k \leq 6$, in (3.17) leads to

$$1/\eta_3^{(1)}(\widehat{\eta}_3^{(1)} - \eta_3^{(1)} + \widehat{\eta}_6^{(1)} - \eta_6^{(1)})Z_3 + 1/\eta_4^{(1)}(\widehat{\eta}_4^{(1)} - \eta_4^{(1)} + \widehat{\eta}_5^{(1)} - \eta_5^{(1)})Z_4 = Z_0.$$

Now, both the coefficient of Z_3 and of Z_4 in the above expression are zero, due to equation (3.18), and it follows that $Z_0 = 0$. So z lies in the hyperplane section at infinity of \overline{Z}_1 and in particular $z \notin U_Z$. Similarly, it is shown that $g_3^{(1)} \neq 1$ on U_Z . It follows that

$$g_Z(U_Z) \subseteq U_G.$$

Now, g_Z has a rational inverse on $\overline{\mathcal{G}}$, given by

$$\begin{aligned} Z_1 &= \eta_1^{(1)} g_2^{(1)} g_1^{(1)}, & Z_3 &= \eta_3^{(1)} g_2^{(1)} g_3^{(1)}, & Z_5 &= \eta_5^{(1)} g_2^{(1)}, \\ Z_2 &= \eta_2^{(1)} g_3^{(1)} / g_1^{(1)}, & Z_4 &= \eta_4^{(1)}, & Z_6 &= \eta_6^{(1)} g_3^{(1)}, \end{aligned} \quad (3.55)$$

and

$$Z_0 = \widehat{\eta}_1^{(1)} g_2^{(1)} g_1^{(1)} + \widehat{\eta}_2^{(1)} g_3^{(1)} / g_1^{(1)} + \widehat{\eta}_3^{(1)} g_2^{(1)} g_3^{(1)} + \widehat{\eta}_4^{(1)} + \widehat{\eta}_5^{(1)} g_2^{(1)} + \widehat{\eta}_6^{(1)} g_3^{(1)}. \quad (3.56)$$

Note that this mapping is analytic on U_G and the image of U_G is a subset of \overline{Z}_1 . Furthermore, note that none of the Z_k , $1 \leq k \leq 6$, can equal zero on U_G , since the coordinates $g_k^{(1)}$, $1 \leq k \leq 3$, are by definition finite and nonzero. In particular, the image of this mapping is disjoint from any of the lines $L_{i,j,k}^{(1)}$, $(i, j, k) \in I$.

It remains to be checked that the image of the mapping is disjoint with the hyperplane section at infinity of \overline{Z}_1 , that is, the right-hand side of equation (3.56) does not vanish on U_G . Take a $g^{(1)} \in U_G$, then $g_2^{(1)}, g_3^{(1)} \neq 1$. The resultant, with respect to $g_1^{(1)}$, of equation (3.52) and the right-hand side of equation (3.56), each after multiplication by $g_1^{(1)}$, is given by

$$\frac{1}{256} g_2^{(1)} g_3^{(1)} (g_2^{(1)} - 1)^2 (g_3^{(1)} - 1)^2 (v_0 - 1)^2 (v_0^{-1} - 1)^2 (v_t - v_t^{-1})^2 (v_1 - v_1^{-1})^2.$$

In particular, it is nonzero and thus the right-hand side of equation (3.56) cannot vanish on U_G . It follows that the rational inverse of g_Z , given above, is analytic and maps U_G into U_Z . We conclude that g_Z maps U_Z biholomorphically onto U_G .

Next, we consider the analytic mapping g_Y on $\overline{\mathcal{Y}}$. Its three components satisfy the cubic relation (3.52). From the explicit formula for $g_2^{(1)}$, it follows that $g_2^{(1)} = 0$ if and only if

$$Y_2 - Y_0 \left(\frac{v_t}{v_\infty} + \frac{v_\infty}{v_t} \right) = 0. \quad (3.57)$$

On the other hand, the intersection of the Segre surface $\overline{\mathcal{Y}}$ with the hyperplane (3.57), is given by the union of the two lines L_{15} and L_{16} . In this way, we obtain the following equivalences,

$$g_2^{(1)} = 0 \iff Y \in L_{15} \cup L_{16}, \quad g_2^{(1)} = \infty \iff Y \in L_{13} \cup L_{14}, \quad (3.58a)$$

$$g_3^{(1)} = 0 \iff Y \in L_{23} \cup L_{24}, \quad g_3^{(1)} = \infty \iff Y \in L_{21} \cup L_{22}. \quad (3.58b)$$

Since, by construction, the open set U_Y is disjoint with the eight lines on the right-hand sides of the above equivalences, $g_2^{(1)}$ and $g_3^{(1)}$ are finite and nonzero on U_Y .

Consequently, by the cubic relation (3.52), $g_1^{(1)}$ is also finite and nonzero on $U_{\mathcal{Y}}$. Furthermore, observe that $g_2^{(1)}$ and $g_3^{(1)}$ cannot attain the value 1 in the affine part of the Segre surface, since this requires $y_2 = \infty$ or $y_3 = \infty$. All in all, it follows that

$$g_{\mathcal{Y}}(U_{\mathcal{Y}}) \subseteq U_{\mathcal{G}}.$$

Now, $g_{\mathcal{Y}}$ has a rational inverse on $\overline{\mathcal{G}}$, given by

$$\begin{aligned} y_2 &= \frac{(v_t v_{\infty} + v_t^{-1} v_{\infty}^{-1})g_2^{(1)} - (v_t v_{\infty}^{-1} + v_t^{-1} v_{\infty})}{g_2^{(1)} - 1}, \\ y_3 &= \frac{(v_1 v_{\infty} + v_1^{-1} v_{\infty}^{-1})g_3^{(1)} - (v_1 v_{\infty}^{-1} + v_1^{-1} v_{\infty})}{g_3^{(1)} - 1}, \\ y_4 &= y_2^{(1)} y_3^{(1)}, \end{aligned}$$

and

$$\begin{aligned} y_1 &= (v_t - v_t^{-1})(v_1 - v_1^{-1})(v_{\infty} - v_{\infty}^{-1}) \frac{g_1^{(1)} g_2^{(1)}}{(g_2^{(1)} - 1)(g_3^{(1)} - 1)} \\ &\quad + v_{\infty}(v_0 + v_0^{-1}) - v_{\infty}(v_{\infty} - v_{\infty}^{-1}) \frac{(v_t g_2^{(1)} - v_t^{-1})(v_1 g_3^{(1)} - v_1^{-1})}{(g_2^{(1)} - 1)(g_3^{(1)} - 1)}. \end{aligned}$$

Note that this mapping is clearly analytic on $U_{\mathcal{G}}$ and the corresponding image lies in $U_{\mathcal{Y}}$. We conclude that $g_{\mathcal{Y}}$ maps $U_{\mathcal{Y}}$ biholomorphically onto $U_{\mathcal{G}}$ and the lemma follows. \square

We now have all the ingredients to prove Theorem 3.8.

Proof of Theorem 3.8. Consider the bi-rational mapping

$$\begin{array}{ccc} \Phi : \overline{\mathcal{Y}} & \dashrightarrow & \overline{\mathcal{Z}}_1 \\ & \mapsto & \mathcal{Z} \end{array} \quad (3.59)$$

obtained by composing $g_{\mathcal{Y}}$, defined in equation (3.51), with the rational inverse of $g_{\mathcal{Z}}$, given in equation (3.55). Due to Lemma 3.16, Φ maps the dense open subset $U_{\mathcal{Y}} \subseteq \overline{\mathcal{Y}}$ biholomorphically onto the dense open subset $U_{\mathcal{Z}} \subseteq \overline{\mathcal{Z}}_1$. We proceed to check that Φ is a biholomorphism between the entire domain and co-domain.

Now, the complements $\overline{\mathcal{Y}} \setminus U_{\mathcal{Y}}$ and $\overline{\mathcal{Z}}_1 \setminus U_{\mathcal{Z}}$ each consist of 8 lines and 2 conics. We first consider the conics on either side. These conics lie in the hyperplane sections $\overline{\mathcal{Y}} \setminus \mathcal{Y}$ and $\overline{\mathcal{Z}}_1 \setminus \mathcal{Z}_1$ of the domain and co-domain respectively. Recall that $\overline{\mathcal{Y}} \setminus \mathcal{Y}$ factors into the two conics described in equations (3.32). On the first conic, equation (3.32a), we have $Y_0 = Y_2 = 0$ and the components of $g_{\mathcal{Y}}$ read

$$\begin{aligned} g_1 &= v_{\infty} \frac{Y_4 + (1 + v_1 v_{\infty}^{-1})Y_1 + (v_t v_1 - v_t^{-1} v_{\infty}^{-1})Y_3}{Y_4 + (1 + v_1 v_{\infty})Y_1 + (v_t v_1 - v_t^{-1} v_{\infty})Y_3}, \\ g_2 &= \frac{Y_4 - (v_t v_{\infty}^{-1} + v_t^{-1} v_{\infty})Y_3}{Y_4 - (v_t v_{\infty} + v_t^{-1} v_{\infty}^{-1})Y_3}, \\ g_3 &= 1. \end{aligned}$$

Applying the rational inverse of $g_{\mathcal{Z}}$ to this, we obtain

$$\begin{aligned} Z_0 &= 0, \\ Z_1/\eta_1^{(1)} &= v_{\infty} Y_4 + (v_{\infty} - v_{\infty}^{-1})Y_1 - (v_t + v_t^{-1})Y_3, \\ Z_2/\eta_2^{(1)} &= v_{\infty}^{-1} Y_4 + (v_{\infty}^{-1} - v_{\infty})Y_1 - (v_t + v_t^{-1})Y_3, \\ Z_3/\eta_3^{(1)} &= Z_5/\eta_5^{(1)} = Y_4 - (v_t v_{\infty}^{-1} + v_t^{-1} v_{\infty})Y_3, \end{aligned}$$

$$Z_4/\eta_4^{(1)} = Z_6/\eta_6^{(1)} = Y_4 - (v_t v_\infty + v_t^{-1} v_\infty^{-1})Y_3.$$

This defines an isomorphism between the conics (3.32a) and (3.20) in $\overline{\mathcal{Y}} \setminus \mathcal{Y}$ and $\overline{\mathcal{Z}}_1 \setminus \mathcal{Z}_1$ respectively. In particular, Φ maps the conic (3.32a) onto the conic (3.20).

Similarly, on the second conic, defined by equation (3.32b), we have $Y_0 = Y_3 = 0$ and the components of $g_{\mathcal{Y}}$ read

$$\begin{aligned} g_1 &= v_\infty \frac{Y_4 + (1 + v_1 v_\infty^{-1})Y_1 + (1 - v_1 v_\infty^{-1})Y_2}{Y_4 + (1 + v_1 v_\infty)Y_1 + (1 - v_1 v_\infty)Y_2}, \\ g_2 &= 1, \\ g_3 &= \frac{Y_4 - (v_1 v_\infty^{-1} + v_1^{-1} v_\infty)Y_2}{Y_4 - (v_1 v_\infty + v_1^{-1} v_\infty^{-1})Y_2}, \end{aligned}$$

Applying the rational inverse of $g_{\mathcal{Z}}$ to this, we obtain

$$\begin{aligned} Z_0 &= 0, \\ Z_1/\eta_1^{(1)} &= v_\infty Y_4 + (v_\infty - v_\infty^{-1})Y_1 - (v_1 + v_1^{-1})Y_2, \\ Z_2/\eta_2^{(1)} &= v_\infty^{-1} Y_4 + (v_\infty^{-1} - v_\infty)Y_1 - (v_1 + v_1^{-1})Y_2, \\ Z_3/\eta_3^{(1)} &= Z_6/\eta_6^{(1)} = Y_4 - (v_1 v_\infty^{-1} + v_1^{-1} v_\infty)Y_2, \\ Z_4/\eta_4^{(1)} &= Z_5/\eta_5^{(1)} = Y_4 - (v_1 v_\infty + v_1^{-1} v_\infty^{-1})Y_2. \end{aligned}$$

This defines an isomorphism between the conics (3.32a) and (3.21) in $\overline{\mathcal{Y}} \setminus \mathcal{Y}$ and $\overline{\mathcal{Z}}_1 \setminus \mathcal{Z}_1$ respectively. In particular, Φ maps the conic (3.32b) onto the conic (3.21).

Next, we consider the remaining lines on either side. By the explicit equation for $g_1^{(1)}$ on $\overline{\mathcal{Z}}_1$ in equations (3.48), we have

$$\begin{aligned} g_1^{(1)} = 0 &\iff Z_1 = Z_6 = 0 \iff Z \in L_{1,3,6}^{(1)} \cup L_{1,4,6}^{(1)}, \text{ and} \\ g_1^{(1)} = \infty &\iff Z_2 = Z_5 = 0 \iff Z \in L_{2,3,5}^{(1)} \cup L_{2,4,5}^{(1)}. \end{aligned}$$

Similarly, for $g_2^{(1)}$ and $g_3^{(1)}$, we find

$$\begin{aligned} g_2^{(1)} = 0 &\iff Z_3 = Z_5 = 0 \iff Z \in L_{1,3,5}^{(1)} \cup L_{2,3,5}^{(1)}, \text{ and} \\ g_2^{(1)} = \infty &\iff Z_4 = Z_6 = 0 \iff Z \in L_{1,4,6}^{(1)} \cup L_{2,4,6}^{(1)}, \text{ and} \\ g_3^{(1)} = 0 &\iff Z_3 = Z_6 = 0 \iff Z \in L_{1,3,6}^{(1)} \cup L_{2,3,6}^{(1)}, \text{ and} \\ g_3^{(1)} = \infty &\iff Z_4 = Z_5 = 0 \iff Z \in L_{1,4,5}^{(1)} \cup L_{2,4,5}^{(1)}. \end{aligned}$$

Comparing these equivalences with those in equation (3.58), we find that

$$\begin{aligned} \Phi(L_{15}^{\mathcal{Y}} \cup L_{16}^{\mathcal{Y}}) &= L_{1,3,5}^{(1)} \cup L_{2,3,5}^{(1)}, & \Phi(L_{13}^{\mathcal{Y}} \cup L_{14}^{\mathcal{Y}}) &= L_{1,4,6}^{(1)} \cup L_{2,4,6}^{(1)}, \\ \Phi(L_{23}^{\mathcal{Y}} \cup L_{24}^{\mathcal{Y}}) &= L_{1,3,6}^{(1)} \cup L_{2,3,6}^{(1)}, & \Phi(L_{21}^{\mathcal{Y}} \cup L_{22}^{\mathcal{Y}}) &= L_{1,4,5}^{(1)} \cup L_{2,4,5}^{(1)}. \end{aligned}$$

To disentangle which line gets sent to which, see Remark 3.10, and really check that Φ maps these lines isomorphically to one another, we require some workable formulas for the mapping Φ . To obtain these, it is convenient to first compute some formulas for the affine z -variables in terms of the affine x -variables on \mathcal{X} .

Upon substituting the formulas for $g_k^{(1)}$, $1 \leq k \leq 3$, in terms of the x -variables in Proposition 3.6, into equation (3.56) for Z_0 , and simplifying modulo the Jimbo-Fricke cubic, we get

$$Z_0 = \frac{\delta}{4w_0(x_2 - (v_t v_\infty + \frac{1}{v_t v_\infty}))(x_3 - (v_1 v_\infty + \frac{1}{v_1 v_\infty}))},$$

where the constant δ is as defined in Theorem 3.8.

As a consequence, substituting the formulas for $g_k^{(1)}$, $1 \leq k \leq 3$, in terms of the x -variables in Proposition 3.6, into equations (3.55) for Z_k , $1 \leq k \leq 6$, and computing the quotient $z_k = Z_k/Z_0$, immediately gives us the formulas for the z_k , $3 \leq k \leq 6$, given in Theorem 3.8. On the other hand, the resulting formulas for z_1 and z_2 are not (yet) in polynomial form, like in Theorem 3.8. Regardless, we obtain the following corresponding formulas with regards to the homogeneous Y -variables on $\overline{\mathcal{Y}}$ and Z -variables on $\overline{\mathcal{Z}}_1$, for the bi-rational mapping Φ ,

$$\begin{aligned}
 Z_0 &= Y_0, \\
 Z_1 &= \gamma^{-1} \left[Y_4 - \left(v_1 v_\infty + \frac{1}{v_1 v_\infty} \right) Y_2 - \left(\frac{v_t}{v_\infty} + \frac{v_\infty}{v_t} \right) Y_3 \right. \\
 &\quad \left. + \left(\frac{v_t}{v_\infty} + \frac{v_\infty}{v_t} \right) \left(v_1 v_\infty + \frac{1}{v_1 v_\infty} \right) Y_0 \right] g_1^{(1)}, \\
 Z_2 &= \gamma^{-1} \left[Y_4 - \left(\frac{v_1}{v_\infty} + \frac{v_\infty}{v_1} \right) Y_2 - \left(v_t v_\infty + \frac{1}{v_t v_\infty} \right) Y_3 \right. \\
 &\quad \left. + \left(v_t v_\infty + \frac{1}{v_t v_\infty} \right) \left(\frac{v_1}{v_\infty} + \frac{v_\infty}{v_1} \right) Y_0 \right] 1/g_1^{(1)}, \\
 Z_3 &= -\delta^{-1} \left(v_0 - v_t v_1 v_\infty \right) \left(v_0 - \frac{1}{v_t v_1 v_\infty} \right) \left[Y_4 - \left(\frac{v_1}{v_\infty} + \frac{v_\infty}{v_1} \right) Y_2 \right. \\
 &\quad \left. - \left(\frac{v_t}{v_\infty} + \frac{v_\infty}{v_t} \right) Y_3 + \left(\frac{v_t}{v_\infty} + \frac{v_\infty}{v_t} \right) \left(\frac{v_1}{v_\infty} + \frac{v_\infty}{v_1} \right) Y_0 \right], \\
 Z_4 &= -\delta^{-1} \left(v_0 - \frac{v_t v_1}{v_\infty} \right) \left(v_0 - \frac{v_\infty}{v_t v_1} \right) \left[Y_4 - \left(v_1 v_\infty + \frac{1}{v_1 v_\infty} \right) Y_2 \right. \\
 &\quad \left. - \left(v_t v_\infty + \frac{1}{v_t v_\infty} \right) Y_3 + \left(v_t v_\infty + \frac{1}{v_t v_\infty} \right) \left(v_1 v_\infty + \frac{1}{v_1 v_\infty} \right) Y_0 \right], \\
 Z_5 &= +\delta^{-1} \left(v_0 - \frac{v_t v_\infty}{v_1} \right) \left(v_0 - \frac{v_1}{v_t v_\infty} \right) \left[Y_4 - \left(v_1 v_\infty + \frac{1}{v_1 v_\infty} \right) Y_2 \right. \\
 &\quad \left. - \left(\frac{v_t}{v_\infty} + \frac{v_\infty}{v_t} \right) Y_3 + \left(\frac{v_t}{v_\infty} + \frac{v_\infty}{v_t} \right) \left(v_1 v_\infty + \frac{1}{v_1 v_\infty} \right) Y_0 \right], \\
 Z_6 &= +\delta^{-1} \left(v_0 - \frac{v_1 v_\infty}{v_t} \right) \left(v_0 - \frac{v_t}{v_1 v_\infty} \right) \left[Y_4 - \left(\frac{v_1}{v_\infty} + \frac{v_\infty}{v_1} \right) Y_2 \right. \\
 &\quad \left. - \left(v_t v_\infty + \frac{1}{v_t v_\infty} \right) Y_3 + \left(v_t v_\infty + \frac{1}{v_t v_\infty} \right) \left(\frac{v_1}{v_\infty} + \frac{v_\infty}{v_1} \right) Y_0 \right].
 \end{aligned} \tag{3.60}$$

where γ is as defined in Theorem 3.8.

Recalling the explicit formulas for $g_1^{(1)}$ with respect to Y in equations (3.49), these equations, as written, are not projective linear in Y . Nonetheless, they are simple enough for explicit computations. In particular, by a local analysis around each line on $\overline{\mathcal{Y}}$, we find that Φ maps it projectively linearly onto a corresponding line in $\overline{\mathcal{Z}}_1$, as detailed in Remark 3.10.

For example, on the line L_{15}^Y , parametrised by

$$Y_1 = \left(v_0 v_\infty + \frac{v_1}{v_t} \right) Y_0 - \frac{v_\infty}{v_t} Y_3, \quad Y_2 = \left(\frac{v_t}{v_\infty} - \frac{v_\infty}{v_t} \right) Y_0, \quad Y_4 = \left(\frac{v_t}{v_\infty} - \frac{v_\infty}{v_t} \right) Y_3,$$

with $[Y_0 : Y_3] \in \mathbb{CP}^1$, we find

$$\begin{aligned}
 Z_0 &= Y_0, \\
 Z_1 &= Z_3 = Z_5 = 0, \\
 Z_2 &= +\gamma^{-1} \left[\left(\frac{v_\infty}{v_t} - \frac{v_t}{v_\infty} \right) Y_3 - Y_0 \left((v_0 + v_0^{-1})(v_\infty - v_\infty^{-1}) - (v_t - v_t)(v_1 + 1/v_1) \right) \right], \\
 Z_4 &= -\delta^{-1} \left(v_0 - \frac{v_t v_1}{v_\infty} \right) \left(v_0 - \frac{v_\infty}{v_t v_1} \right) (v_t - v_t^{-1})(v_\infty - v_\infty^{-1}) \left(Y_3 - Y_0 \left(v_1 v_\infty + \frac{1}{v_1 v_\infty} \right) \right),
 \end{aligned}$$

$$Z_6 = +\delta^{-1}\left(v_0 - \frac{v_1 v_\infty}{v_t}\right)\left(v_0 - \frac{v_t}{v_1 v_\infty}\right)(v_t - v_t^{-1})(v_\infty - v_\infty^{-1})\left(Y_3 - Y_0\left(\frac{v_1}{v_\infty} + \frac{v_\infty}{v_1}\right)\right),$$

so that Φ maps L_{15}^Y projectively linearly onto $L_{1,3,5}^{(1)}$.

All in all, we know that Φ maps the dense open subset $U_{\mathcal{Y}}$ biholomorphically onto $U_{\mathcal{Z}}$ and it maps each irreducible component of $\overline{\mathcal{Y}} \setminus U_{\mathcal{Y}}$ onto a corresponding irreducible component of $\overline{\mathcal{Z}}_1 \setminus U_{\mathcal{Z}}$. To conclude that Φ is an isomorphism, we may proceed in two ways.

The first, is to check that Φ is locally biholomorphic around each point in the complement of $U_{\mathcal{Z}}$. From this it follows that Φ is a locally biholomorphic bi-rational mapping and thus an isomorphism.

Alternatively, one can check that Φ respects the intersections between the different irreducible components in $\overline{\mathcal{Y}} \setminus U_{\mathcal{Y}}$ and $\overline{\mathcal{Z}}_1 \setminus U_{\mathcal{Z}}$, as follows by comparing the Clebsch graphs in Figures 2.1 and 3.1 and equations (3.47), noting in particular that all the ForestGreen coloured lines in Figure 2.1 are mapped to the lines $L_k^{\mathcal{Y}}$, $9 \leq k \leq 16$, and all the purple coloured lines are mapped to the lines $L_k^{\mathcal{Y}}$, $17 \leq k \leq 24$, see Remark 3.11. This way, we can check that Φ is a bi-rational bijection between $\overline{\mathcal{Y}}$ and $\overline{\mathcal{Z}}_1$. Since $\overline{\mathcal{Y}}$ and $\overline{\mathcal{Z}}_1$ are smooth and thus normal and irreducible, it follows from Zariski's main theorem that Φ is an isomorphism.

Either way yields the required result. We now apply Lemma 2.6, which shows that Φ must extend to a unique (rank four) projective linear map between the ambient spaces \mathbb{P}^4 and \mathbb{P}^6 . Furthermore, since Φ maps the hyperplane sections at infinity to one another, it follows that Φ also restricts to an affine equivalence $\Phi_{\mathcal{Y}} := \Phi|_{\mathcal{Y}}$ between the affine Segre surfaces \mathcal{Y} and \mathcal{Z}_1 ,

$$\Phi_{\mathcal{Y}} : \mathcal{Y} \rightarrow \mathcal{Z}_1.$$

Furthermore, since the blow-down mapping π , defined in (3.31), is an isomorphism between \mathcal{X} and \mathcal{Y} as affine varieties when restricted to \mathcal{X} , the map

$$\Phi_{\mathcal{X}} = \Phi_{\mathcal{Y}} \circ \pi|_{\mathcal{X}},$$

is an isomorphism of affine varieties,

$$\Phi_{\mathcal{X}} : \mathcal{X} \rightarrow \mathcal{Z}_1.$$

What is left, is to derive the explicit formulas for $\Phi_{\mathcal{X}}$ in the theorem. Since $\Phi_{\mathcal{Y}}$ is an affine equivalence, we know that it can be written as an affine linear map as in equation (3.46). Therefore, $\Phi_{\mathcal{X}}$ can be written as

$$z_k = \xi_{0k} + \xi_{1k} x_1 + \xi_{2k} x_2 + \xi_{3k} x_3 + \xi_{4k} x_2 x_3, \quad 1 \leq k \leq 6, \quad (3.61)$$

for some coefficients ξ_{jk} , $0 \leq j \leq 4$, $1 \leq k \leq 6$.

The formulas for Z_k , $k = 0, 3, 4, 5, 6$, in equations (3.60), immediately give us the formulas for z_k , $3 \leq k \leq 6$, in terms of the affine x -variables in the theorem, using

$$Y_1 = x_1 Y_0, \quad Y_2 = x_2 Y_0, \quad Y_3 = x_3 Y_0, \quad Y_4 = x_2 x_3 Y_0.$$

Note, in particular, that these formulas are of the form (3.61).

Finally, we compute the coefficients in (3.61) for $k = 1, 2$ to obtain the formulas for z_1 and z_2 in Theorem 3.8. We do this by computing the z_1 and z_2 entries of the images under $\Phi_{\mathcal{X}}$ of five explicit points on the Jimbo-Fricke cubic,

$$\begin{aligned} p_1 : & \quad x_1 = v_0 v_\infty + v_0^{-1} v_\infty^{-1}, & x_2 = 2, & x_3 = -2v_0 v_\infty + v_0 v_t + v_\infty v_1, \\ p_2 : & \quad x_1 = v_0 v_\infty^{-1} + v_0^{-1} v_\infty, & x_2 = 2, & x_3 = -2v_0 v_\infty^{-1} + v_0 v_t + v_\infty^{-1} v_1, \\ p_3 : & \quad x_2 = v_0 v_1 + v_0^{-1} v_1^{-1}, & x_1 = 2, & x_3 = -2v_0 v_1 + v_0 v_t + v_1 v_\infty, \\ p_4 : & \quad x_3 = v_0 v_t^{-1} + v_0^{-1} v_t, & x_1 = 2, & x_2 = -2v_0^{-1} v_t + v_0^{-1} v_1 + v_t v_\infty, \\ p_5 : & \quad x_3 = v_0 v_t + v_0^{-1} v_t^{-1}, & x_1 = 2, & x_2 = -2v_0 v_t + v_0 v_1 + v_t v_\infty. \end{aligned}$$

The corresponding values of z_1 and z_2 under $\Phi_{\mathcal{X}}$ of these points are as follows,

$$\begin{aligned}
 z_1(p_1) &= -\frac{(2v_0v_\infty - v_0\nu_t - v_1(v_\infty - v_\infty^{-1}))(2v_0v_\infty - v_0\nu_t - v_1^{-1}(v_\infty - v_\infty^{-1}))}{v_0\gamma}, \\
 z_2(p_1) &= -\frac{v_0(v_t - v_\infty)^2(v_tv_\infty - 1)^2}{v_t^2v_\infty^2\gamma}, \\
 z_1(p_2) &= -\frac{(2v_0v_\infty - v_1\nu_t - v_0(v_\infty - v_\infty^{-1}))(2v_1v_\infty^{-1} - v_1^{-1}\nu_t - v_0(v_\infty - v_\infty^{-1}))}{v_0\gamma}, \\
 z_2(p_2) &= -\frac{(v_t - v_\infty)^2(v_tv_\infty - 1)^2}{v_0v_t^2v_\infty^2\gamma}, \\
 z_1(p_3) &= -\frac{(v_0v_tv_1v_\infty - 1)(v_0v_1v_\infty - v_t)(2v_0v_1 - v_0\nu_t - v_\infty(v_1 - v_1^{-1}))}{v_0v_tv_1v_\infty\gamma}, \\
 z_2(p_3) &= -\frac{(v_0v_tv_1 - v_\infty)(v_0v_1 - v_tv_\infty)(2v_0v_1 - v_0\nu_t - v_\infty^{-1}(v_1 - v_1^{-1}))}{v_0v_tv_1v_\infty\gamma}, \\
 z_1(p_4) &= -\frac{(v_0v_1 - v_tv_\infty)(v_0 - v_tv_1v_\infty)(2v_0^{-1}v_t - v_0^{-1}\nu_1 - v_\infty(v_t - v_t^{-1}))}{v_0v_tv_1v_\infty\gamma}, \\
 z_2(p_4) &= -\frac{(v_0v_1v_\infty - v_t)(v_0v_\infty - v_tv_1)(2v_0^{-1}v_t - v_0^{-1}\nu_1 - v_\infty^{-1}(v_t - v_t^{-1}))}{v_0v_tv_1v_\infty\gamma}, \\
 z_1(p_5) &= -\frac{(v_0v_tv_1v_\infty - 1)(v_0v_tv_\infty - v_1)(2v_0v_t - v_0\nu_1 - v_\infty(v_t - v_t^{-1}))}{v_0v_tv_1v_\infty\gamma}, \\
 z_2(p_5) &= -\frac{(v_0v_tv_1 - v_\infty)(v_0v_t - v_1v_\infty)(2v_0v_t - v_0\nu_1 - v_\infty^{-1}(v_t - v_t^{-1}))}{v_0v_tv_1v_\infty\gamma}.
 \end{aligned}$$

By plugging these values into equations (3.61), we get two sets of five equations among five unknown coefficients, which each have a unique solution. The result yields the two explicit equations for z_1 and z_2 in Theorem 3.8, thus concluding the proof the theorem. \square

Remark 3.17. We note that the involutive automorphism of the Segre surface $\overline{\mathcal{Z}}_1$, given by swapping z_1 and z_2 , corresponds to the following automorphism of $\overline{\mathcal{Y}}$,

$$y_1 \mapsto -\omega_1 - y_1 - y_4, \quad y_{2,3,4} \mapsto y_{2,3,4},$$

via the isomorphism in Theorem 3.8. In turn, via the blow-down of the Jimbo-Fricke cubic, this defines an automorphism of the affine cubic \mathcal{X} ,

$$x_1 \mapsto -\omega_1 - x_1 - x_2x_3, \quad x_{2,3} \mapsto x_{2,3},$$

which is one of the generators of the extended modular group action on the Jimbo-Fricke cubic [23].

Regarding the three Tyurin ratios $g^{(q)} = (g_1^{(q)}, g_2^{(q)}, g_3^{(q)})$, the relation (3.52) when $q = 1$, generalises to

$$\eta_1^{(q)} g_2^{(q)} g_1^{(q)} + \eta_2^{(q)} g_3^{(q)} / g_1^{(q)} + \eta_3^{(q)} g_2^{(q)} g_3^{(q)} + \eta_4^{(q)} + \eta_5^{(q)} g_2^{(q)} + \eta_6^{(q)} g_3^{(q)} = 0. \quad (3.62)$$


This equation is quadratic in $g_1^{(q)}$ and the involutive automorphism above is equivalent to sending $g_1^{(q)}$ to its other root of this quadratic, with $q = 1$, explicitly

$$g_1^{(q)} \mapsto -1/\eta_1^{(q)}(\eta_3^{(q)} g_3^{(q)} + \eta_4^{(q)} / g_2^{(q)} + \eta_5^{(q)} + \eta_6^{(q)} g_3^{(q)} / g_2^{(q)}), \quad g_{2,3}^{(q)} \mapsto g_{2,3}^{(q)}.$$

The last formula gives a natural generalisation of the above automorphism to \mathcal{Z}_q for $q \neq 1$, though its action on the z -variables seems quite involved.

4. THE OTHER PAINLEVÉ DIFFERENTIAL EQUATIONS

In [9] it was proved that the confluence scheme of the Painlevé differential equations



corresponds to appropriate limits on the associated monodromy manifolds. In this section, we show that, in all un-ramified cases (in blue), the confluence scheme of the Painlevé monodromy manifolds can be carried through to the affine transformation constructed in Theorem 3.8, therefore producing isomorphisms between each \mathcal{Y} -Segre, i.e. those ones resulting from blowing down the monodromy manifold at a line at infinity, and the \mathcal{Z} -Segre obtained by confluing \mathcal{Z}_1 .

For the ramified cases (in red), the confluence either produces a reducible Segre surface or a family which does not have the correct number of free parameters. This is not surprising because the confluences to ramified and non-ramified Painlevé equations are deeply different in geometric as well as analytic terms. In Section 6, we perform an in depth study of the singularity structure of all cubic surfaces with a triangle of lines at infinity, of their blow downs to the corresponding \mathcal{Y} -Segre surfaces and the expected singularity structure of the \mathcal{Z} -Segre ones. This will allow us to build the isomorphic \mathcal{Z} -Segre for all ramified cases.

This section is organised as follows: In Subsection 4.1 we summarise the confluence of monodromy manifolds obtained in [9] for the non-ramified cases and deduce the confluence on the corresponding \mathcal{Y} -Segre surfaces. Then, in 4.2, we apply the confluence of monodromy manifolds to the affine transformation constructed in Theorem 3.8 and provide the Segre surfaces for all non-ramified cases.

4.1. Confluence of monodromy manifolds. Following [49] and [9] the monodromy manifolds of all Painlevé equations are given by

$$\mathcal{X}^{(d)} := \text{Spec}(\mathbb{C}[x_1, x_2, x_3]/\langle \phi^{(d)} = 0 \rangle) \quad (4.1)$$

where the polynomial $\phi^{(d)}$ has always the same form in the variables x_1, x_2, x_3 , but different coefficients for different Painlevé equations:

$$\phi^{(d)} = x_1 x_2 x_3 + \epsilon_1^{(d)} x_1^2 + \epsilon_2^{(d)} x_2^2 + \epsilon_3^{(d)} x_3^2 + \omega_1^{(d)} x_1 + \omega_2^{(d)} x_2 + \omega_3^{(d)} x_3 + \omega_4^{(d)} = 0. \quad (4.2)$$

Here d is an index running on the list of all the Painlevé auxiliary linear problems $P_{VI}, P_V, P_V^{\text{deg}}, P_{IV}, P_{III}^{D_6}, P_{III}^{D_7}, P_{III}^{D_8}, P_{II}^{\text{JM}}, P_{II}^{\text{FN}}, P_I$, the parameters $\epsilon_i^{(d)}$ may take value 0 or 1 according to the chosen d , and the parameters $\omega_i^{(d)}$, $i = 1, 2, 3, 4$ are related to the Painlevé equations constants as given in section 2 of [9]. For convenience, we summarise the monodromy manifolds of the non-ramified Painlevé equations in Table 4.1.

	monodromy manifold	parameters
P_{VI}	$x_1x_2x_3 + \sum_{k=1}^3(x_k^2 + \omega_kx_k) + \omega_4 = 0$	$\omega_1 = -(\nu_0\nu_\infty + \nu_t\nu_1),$ $\omega_2 = -(\nu_0\nu_1 + \nu_t\nu_\infty),$ $\omega_3 = -(\nu_0\nu_t + \nu_1\nu_\infty),$ $\omega_4 = \nu_0^2 + \nu_t^2 + \nu_1^2 + \nu_\infty^2 + \nu_0\nu_t\nu_1\nu_\infty - 4.$
P_V	$x_1x_2x_3 + x_1^2 + x_2^2 + \sum_{k=1}^3\omega_kx_k + \omega_4 = 0$	$\omega_1 = -(\nu_0\nu_\infty + \nu_t),$ $\omega_2 = -(\nu_0 + \nu_t\nu_\infty),$ $\omega_3 = -(\nu_0\nu_t + \nu_\infty),$ $\omega_4 = 1 + \nu_\infty^2 + \nu_0\nu_t\nu_\infty.$
P_V^{deg}	$x_1x_2x_3 + x_1^2 + x_2^2 + \sum_{k=1}^3\omega_kx_k + \omega_4 = 0$	$\omega_1 = -\nu_0$ $\omega_2 = -\nu_t,$ $\omega_3 = 0,$ $\omega_4 = 1.$
P_{IV}	$x_1x_2x_3 + x_1^2 + \sum_{k=1}^3\omega_kx_k + \omega_4 = 0$	$\omega_1 = -(\nu_0\nu_\infty + 1),$ $\omega_2 = \omega_3 = -\nu_\infty,$ $\omega_4 = \nu_\infty^2 + \nu_0\nu_\infty.$
$P_{III}^{D_6}$	$x_1x_2x_3 + x_1^2 + x_2^2 + \sum_{k=1}^3\omega_kx_k + \omega_4 = 0$	$\omega_1 = -\nu_0\nu_\infty - 1$ $\omega_2 = -(\nu_0 + \nu_\infty),$ $\omega_3 = 0,$ $\omega_4 = \nu_0\nu_\infty.$
$P_{III}^{D_7}$	$x_1x_2x_3 + x_1^2 + x_2^2 + \sum_{k=1}^3\omega_kx_k + \omega_4 = 0$	$\omega_1 = -1$ $\omega_2 = -\nu_\infty,$ $\omega_3 = 0,$ $\omega_4 = 0.$
$P_{III}^{D_8}$	$x_1x_2x_3 + x_1^2 + x_2^2 + \sum_{k=1}^3\omega_kx_k + \omega_4 = 0$	$\omega_1 = \omega_3 = \omega_4 = 0,$ $\omega_2 = -1.$
P_{II}^{FN}	$x_1x_2x_3 + x_1^2 + \sum_{k=1}^3\omega_kx_k + \omega_4 = 0$	$\omega_1 = -(\nu_0 + \nu_0^{-1}),$ $\omega_2 = -1 \quad \omega_3 = 0,$ $\omega_4 = 1.$
P_{II}^{JM}	$x_1x_2x_3 + \sum_{k=1}^3\omega_kx_k + \omega_4 = 0$	$\omega_1 = \omega_2 = \omega_3 = -\nu_\infty,$ $\omega_4 = \nu_\infty(1 + \nu_\infty).$
P_I	$x_1x_2x_3 + \sum_{k=1}^3\omega_kx_k + \omega_4 = 0$	$\omega_1 = -1, \quad \omega_2 = -1,$ $\omega_3 = 0, \quad \omega_4 = 1.$

TABLE 4.1. The monodromy manifolds of the Painlevé differential equations - red denotes ramified cases and blue un-ramified. The parameters $\nu_i, v_i, i = 0, t, 1, \infty$ are defined in (3.24).

The confluence procedure always involves a re-scaling of two variables $x_i, x_j, i \neq j$ and of some of the parameters in ϵ followed by taking the limit for $\epsilon \rightarrow 0$. We blow down each monodromy manifold to a \mathcal{Y} -Segre surface always in the same way, namely we set

$$y_1 := x_1, \quad y_2 := x_2, \quad y_3 := x_3, \quad y_4 := x_2x_3.$$

We summarise these confluences in table 4.2.

Confluence	Re-scaling on x_1, x_2, x_3	Re-scaling on \mathcal{Y} -Segre	Re-scaling on parameters
$P_{VI} \mapsto P_V$	$x_1 \mapsto \frac{x_1}{\epsilon}, x_2 \mapsto \frac{x_2}{\epsilon},$	$y_1 \mapsto \frac{y_1}{\epsilon}, y_2 \mapsto \frac{y_2}{\epsilon}, y_4 \mapsto \frac{y_4}{\epsilon}$	$\nu_\infty \mapsto \frac{\nu_\infty}{\epsilon}, \nu_1 \mapsto \frac{1}{\epsilon}$
$P_V \mapsto P_{IV}$	$x_1 \mapsto \frac{x_1}{\epsilon}, x_3 \mapsto \frac{x_3}{\epsilon},$	$y_1 \mapsto \frac{y_1}{\epsilon}, y_3 \mapsto \frac{y_3}{\epsilon}, y_4 \mapsto \frac{y_4}{\epsilon}$	$\nu_\infty \mapsto \frac{\nu_\infty}{\epsilon}, \nu_t \mapsto \frac{1}{\epsilon}$
$P_V \mapsto P_V^{\text{deg}}$	$x_1 \mapsto \frac{x_1}{\epsilon}, x_2 \mapsto \frac{x_2}{\epsilon},$	$y_1 \mapsto \frac{y_1}{\epsilon}, y_2 \mapsto \frac{y_2}{\epsilon}, y_4 \mapsto \frac{y_4}{\epsilon}$	$\nu_\infty \mapsto \frac{1}{\epsilon}, \nu_1 \mapsto \frac{\nu_1}{\epsilon}$
$P_V \mapsto P_{III}^{D_6}$	$x_1 \mapsto \frac{x_1}{\epsilon}, x_2 \mapsto \frac{x_2}{\epsilon},$	$y_1 \mapsto \frac{y_1}{\epsilon}, y_2 \mapsto \frac{y_2}{\epsilon}, y_4 \mapsto \frac{y_4}{\epsilon}$	$\nu_0 \mapsto \frac{\nu_0}{\epsilon}, \nu_t \mapsto \frac{1}{\epsilon}$
$P_{III}^{D_6} \mapsto P_{III}^{D_7}$	$x_1 \mapsto \frac{x_1}{\epsilon}, x_2 \mapsto \frac{x_2}{\epsilon},$	$y_1 \mapsto \frac{y_1}{\epsilon}, y_2 \mapsto \frac{y_2}{\epsilon}, y_4 \mapsto \frac{y_4}{\epsilon}$	$\nu_0 \mapsto \frac{1}{\epsilon}, \nu_\infty \mapsto \epsilon \nu_\infty$
$P_{IV} \mapsto P_{II}^{\text{JM}}$	$x_2 \mapsto \frac{x_2}{\epsilon}, x_3 \mapsto \frac{x_3}{\epsilon},$	$y_2 \mapsto \frac{y_2}{\epsilon}, y_3 \mapsto \frac{y_3}{\epsilon}, y_4 \mapsto \frac{y_4}{\epsilon^2}$	$\nu_0 \mapsto \frac{1}{\epsilon}, \nu_\infty \mapsto \frac{\nu_\infty}{\epsilon}$

TABLE 4.2. Re-scalings giving rise to the confluence of the non-ramified Painlevé monodromy manifolds and \mathcal{Y} -Segre surfaces.

4.2. Confluence of the affine transformation between \mathcal{Y} -Segre and \mathcal{Z} -Segre. The affine transformation between \mathcal{Y} -Segre and \mathcal{Z} -Segre has always the same form:

$$z_k = \xi_{0k} + \xi_{1k}y_1 + \xi_{2k}y_2 + \xi_{3k}y_3 + \xi_{4k}y_4, \quad k = 1, \dots, 6$$

where the coefficients ξ_{jk} , $k = 1, \dots, 6$, $j = 0, \dots, 4$, depend on ν_i , $i = 0, t, 1, \infty$ as in (3.24).

4.2.1. *Segre surface of P_V .* The confluence limit from P_{VI} to P_V , we see that the re-scaling of the parameters in the last column of Table 4.2 produces the following re-scalings on the coefficients ξ_{kj} :

$$\xi_{0k} \mapsto \xi_{0k}, \quad \xi_{1k} \mapsto \frac{\xi_{1k}}{\epsilon}, \quad \xi_{2k} \mapsto \frac{\xi_{2k}}{\epsilon}, \quad \xi_{3k} \mapsto \xi_{3k}, \quad \xi_{4k} \mapsto \frac{\xi_{4k}}{\epsilon},$$

with the limiting ξ_{jk} not all zero for some fixed k . This means that all z_k remain finite because the re-scalings for the coefficients ξ_{jk} are compensated by the re-scaling in the y_1, \dots, y_4 variables. Therefore the \mathcal{Z} -Segre equations maintain the same form. The parameters are re-scaled as follows

$$\rho_3 \rightarrow \epsilon \rho_3, \quad \rho_4 \rightarrow \rho_4, \quad \rho_4 \rightarrow \rho_4, \quad \rho_6 \rightarrow \epsilon \rho_6, \quad \rho_5 \rho_6 \rho_3 \rightarrow \lambda_2, \quad \lambda_1 \rightarrow \lambda_1,$$

therefore the parameters ρ_3, ρ_6 become 0.

$$z_1 + z_2 + z_3 + z_4 + z_5 + z_6 = 0, \quad (4.3a)$$

$$z_4 + \rho_5 z_5 - 1 = 0, \quad (4.3b)$$

$$z_3 z_4 - \lambda_1 z_1 z_2 = 0, \quad (4.3c)$$

$$z_5 z_6 - \lambda_2 z_1 z_2 = 0. \quad (4.3d)$$

Explicitly, setting

$$\gamma = (v_0 - 1) \left(\frac{1}{v_0} \right) v_\infty^2, \quad \delta = (v_0 - 1)^2 \nu_t v_\infty^2,$$

the formulae relating the \mathcal{Z} -Segre to the monodromy manifold in the P_V case are:

$$z_1 = \gamma^{-1} (x_2 (v_\infty x_3 - 1) + v_\infty (x_1 - v_\infty \nu_0)),$$

$$z_2 = -\gamma^{-1} (x_2 - \nu_t v_\infty) + v_\infty x_1,$$

$$\begin{aligned}
 z_3 &= \delta^{-1}v_0(v_t x_2 - v_\infty)(v_\infty x_3 - v_\infty^2 - 1), \\
 z_4 &= \delta^{-1}v_t^{-1}(v_\infty v_0 - v_t)(v_t v_0 - v_\infty)(x_2 - v_t v_\infty), \\
 z_5 &= -\delta^{-1}v_t^{-2}(v_\infty v_0 v_t - 1)(v_0 - v_t v_\infty)(v_t x_2 - v_\infty), \\
 z_6 &= -\delta^{-1}v_t^{-1}v_0(v_\infty x_3 - v_\infty^2 - 1)(x_2 - v_\infty v_t).
 \end{aligned}$$

4.2.2. *Segre surface of P_{IV} .* The confluence limit from P_V to P_{IV} behaves in a similar way to the one from P_{VI} to P_V , so we omit the discussion.

$$z_1 + z_2 + z_3 + z_4 + z_5 + z_6 = 0, \quad (4.4a)$$

$$z_4 - 1 = 0, \quad (4.4b)$$

$$z_3 z_4 - \lambda_1 z_1 z_2 = 0, \quad (4.4c)$$

$$z_5 z_6 - \lambda_2 z_1 z_2 = 0. \quad (4.4d)$$

Explicitly, the formulae relating the \mathcal{Z} -Segre to the monodromy manifold in the P_{IV} case are:

$$\begin{aligned}
 z_1 &= \frac{1 + v_0^2 - \frac{v_0}{v_\infty}x_1 - \frac{v_\infty}{v_1}x_2 x_3}{(v_0 - 1)^2}, & z_2 &= \frac{v_0(x_1 - 1)}{v_\infty(v_0 - 1)^2}, \\
 z_3 &= \frac{v_0(x_2 - v_\infty)(x_3 - v_\infty)}{v_\infty(v_0 - 1)^2}, & z_4 &= \frac{(v_0 - v_\infty)(v_\infty v_0 - 1)}{v_\infty(v_0 - 1)^2}, \\
 z_5 &= \frac{v_0(x_2 - v_\infty)}{(v_0 - 1)^2}, & z_6 &= \frac{v_0(x_3 - v_\infty)}{(v_0 - 1)^2}.
 \end{aligned}$$

4.2.3. *Segre surface of $P_{III}^{D_6}$.* The re-scaling of the parameters in the last column of Table 4.2 corresponding to the confluence limit from P_V to $P_{III}^{D_6}$ produces the following re-scalings on the coefficients ξ_{kj} :

$$\xi_{0k} \mapsto \xi_{0k}, \quad \xi_{1k} \mapsto \frac{\xi_{1k}}{\epsilon}, \quad \xi_{2k} \mapsto \frac{\xi_{2k}}{\epsilon}, \quad \xi_{3k} \mapsto \xi_{3k}, \quad \xi_{4k} \mapsto \frac{\xi_{4k}}{\epsilon},$$

with $\xi_{j6} = \mathcal{O}(\epsilon^2)$ for $j = 0, 3$ and $\xi_{j6} = \mathcal{O}(\epsilon^3)$ for $j = 1, 2, 4$. This means that the variable z_6 is of order ϵ^2 and drops out of the first two equations defining the \mathcal{Z} -Segre. The parameters are rescaled as follows

$$\rho_5 \rightarrow \rho_5, \quad \lambda_1 \rightarrow \lambda_1, \quad \lambda_2 \rightarrow \epsilon^2 \lambda_2.$$

These re-scalings imply that the last two equations defining the \mathcal{Z} -Segre preserve their form. By rescaling z_6 , we can set $\lambda_2 = 1$:

$$z_1 + z_2 + z_3 + z_4 + z_5 = 0, \quad (4.5a)$$

$$z_4 + \rho_5 z_5 - 1 = 0, \quad (4.5b)$$

$$z_3 z_4 - \lambda_1 z_1 z_2 = 0, \quad (4.5c)$$

$$z_5 z_6 - z_1 z_2 = 0. \quad (4.5d)$$

Explicitly, the formulae relating the \mathcal{Z} -Segre to the monodromy manifold in the P_{III} case are:

$$\begin{aligned}
 z_1 &= \frac{v_0 v_\infty^2 - v_\infty x_1 + x_2 - v_\infty x_2 x_3}{v_0 v_\infty^2}, & z_2 &= \frac{v_\infty x_1 + x_2 - v_\infty}{v_0 v_\infty^2}, \\
 z_3 &= \frac{x_2(v_\infty x_3 - v_\infty^2 - 1)}{v_0 v_\infty^2}, & z_4 &= \frac{(v_0 v_\infty - 1)(x_2 - v_\infty)}{v_0 v_\infty^2}, \\
 z_5 &= \frac{x_2(v_\infty - v_0)}{v_0 v_\infty^2}, & z_6 &= \frac{(x_2 - v_\infty)(1 + v_\infty^2 - v_\infty x_3)}{v_0 v_\infty^2}.
 \end{aligned}$$

4.2.4. *Segre surface of P_{II}^{JM}* . The re-scaling of the parameters in the last column of Table 4.2 corresponding to the confluence limit from P_{IV} to P_{II}^{JM} produces the following re-scaling on the coefficients ξ_{kj} :

$$\xi_{0k} \mapsto \xi_{0k}, \quad \xi_{1k} \mapsto \xi_{1k}, \quad \xi_{2k} \mapsto \frac{\xi_{2k}}{\epsilon}, \quad \xi_{3k} \mapsto \frac{\xi_{3k}}{\epsilon}, \quad \xi_{4k} \mapsto \frac{\xi_{4k}}{\epsilon^2},$$

with $\xi_{j2} = \mathcal{O}(\epsilon^2)$ for $j = 0, \dots, 4$. This means that the variable z_2 is of order ϵ^2 and drops out of the first equation defining the \mathcal{Z} -Segre. The parameters are re-scaled as follows

$$\rho_4 \rightarrow \rho_4, \quad \lambda_1 \rightarrow \frac{\lambda_1}{\epsilon^2}, \quad \lambda_2 \rightarrow \frac{\lambda_2}{\epsilon^2},$$

so that the last two equations defining the \mathcal{Z} -Segre preserve their form.

$$z_1 + z_3 + z_4 + z_5 + z_6 = 0 \tag{4.6a}$$

$$\rho_4 z_4 - 1 = 0 \tag{4.6b}$$

$$z_3 z_4 - \lambda_1 z_1 z_2 = 0 \tag{4.6c}$$

$$z_3 z_4 - \lambda_2 z_5 z_6 = 0. \tag{4.6d}$$

Explicitly

$$z_1 = 1 - \frac{x_2 x_3}{v_\infty}, \quad z_2 = \frac{x_1 - 1}{v_\infty}, \quad z_3 = v_\infty - x_2 - x_3 + \frac{x_2 x_3}{v_\infty},$$

$$z_4 = v_\infty - 1, \quad z_5 = x_2 - v_\infty, \quad z_6 = x_3 - v_\infty.$$

Note that we can re-scale all z_i variables to set $\rho_4 = 1$, then we can absorb the parameter λ_1 in z_2 , hence the final \mathcal{Z} -Segre for P_{II}^{JM} is a one parameter family as expected.

5. BLOW-DOWNS OF AFFINE CUBIC SURFACES

Consider an embedded affine cubic surface $\mathcal{X} \subseteq \mathbb{C}^3$ together with its canonical projective completion $\bar{\mathcal{X}} \subseteq \mathbb{P}^3$. We say that \mathcal{X} has a triangle (of lines) at infinity, if the hyperplane section at infinity, $\bar{\mathcal{X}} \setminus \mathcal{X}$, is a cubic curve which is the product of three lines that intersect pairwise at distinct points.

The embedded affine cubic surfaces corresponding to differential Painlevé equations [9, 49] are all smooth for generic parameter values with a triangle of lines at infinity. They are, however, geometrically distinguished by their singularity structures on this triangle. This is important in order to characterise the lines on the \mathcal{Y} -Segre obtained as blow down of the corresponding affine cubics. Indeed, as discussed in Subsections 3.3 and 4, understanding the lines in the \mathcal{Y} -Segre is key to construct the isomorphism to the \mathcal{Z} -Segre in the ramified cases.

In this section, we study smooth embedded affine cubic surfaces, with a triangle at infinity, and affine Segre surfaces naturally associated to them.

In Section 5.1, we classify all embedded smooth cubic surfaces with a triangle at infinity. Then, in Section 5.2, we give a natural construction of three associated Segre surfaces, in the regular case corresponding to blowing down any of the three lines at infinity. In Section 5.3, we use this construction to derive an explicit isomorphism between the two (decorated) character varieties for P_{II} known in the literature, one coming from the Jimbo-Miwa linear problem, the other from the Flaschka-Newell linear problem. Similarly, in sections 5.4, 5.5 and 5.6 we build the remaining \mathcal{Z} -Segre surfaces for the ramified Painlevé equations.

5.1. A classification. In this section, we classify embedded smooth affine cubic surfaces with a triangle at infinity. We start by deriving a normal form.

Lemma 5.1. *Any embedded affine cubic surface in \mathbb{C}^3 , with a triangle of lines at infinity, is affinely equivalent to*

$$x_1x_2x_3 + \epsilon_1x_1^2 + \epsilon_2x_2^2 + \epsilon_3x_3^2 + \omega_1x_1 + \omega_2x_2 + \omega_3x_3 + \omega_4 = 0, \quad (5.1)$$

for some $\epsilon_{1,2,3} \in \{0, 1\}$ and $\omega_k \in \mathbb{C}$, $1 \leq k \leq 4$.

Proof. Let \mathcal{X} be an affine cubic surface in $\{(x_1, x_2, x_3) \in \mathbb{C}^3\}$. Denote its canonical projective completion by $\overline{\mathcal{X}}$ in \mathbb{P}^3 , so that the curve at infinity $\overline{\mathcal{X}} \setminus \mathcal{X}$ is a triangle of lines. Using projective coordinates,

$$[X_0 : X_1 : X_2 : X_3] = [1 : x_1 : x_2 : x_3], \quad (5.2)$$

the curve at infinity is thus described by

$$L_1L_2L_3 = 0, \quad X_0 = 0,$$

where each $L_k = L_k(X_1, X_2, X_3)$ is homogenous and linear. Since the three lines $L_k = 0$, $1 \leq k \leq 3$, pairwise intersect in distinct points, the affine map

$$x_k \mapsto L_k(x_1, x_2, x_3) \quad (1 \leq k \leq 3),$$

has full rank and application of its inverse puts the cubic surface into the form

$$x_1x_2x_3 + a_1x_2x_3 + a_2x_1x_3 + a_3x_1x_2 + Q(x_1, x_2, x_3) = 0, \quad (5.3)$$

for some $a_{1,2,3} \in \mathbb{C}$ and a quadratic polynomial Q without mixed terms. Applying the affine scaling $x_k \mapsto x_k - a_k$, we may eliminate all mixed terms from (5.3), so that we are left with equation (5.1), for some $\epsilon_{1,2,3} \in \mathbb{C}$ and $\omega_k \in \mathbb{C}$, $1 \leq k \leq 4$. By finally rescaling $x_k \mapsto c_kx_k$, for some $c_k \in \mathbb{C}^*$, $1 \leq k \leq 3$, we can ensure that each $\epsilon_k \in \{0, 1\}$ and the lemma follows. \square

Next, we are going to have a look at singularities at infinity. Take any (irreducible) affine cubic surface $\mathcal{X} \subseteq \mathbb{C}^3$, with a triangle of lines at infinity, in normal form (5.1). Using homogeneous coordinates (5.2), its canonical projective completion is given by the zero locus of the homogeneous polynomial

$$F := X_1X_2X_3 + (\epsilon_1X_1^2 + \epsilon_2X_2^2 + \epsilon_3X_3^2)X_0 + (\omega_1X_1 + \omega_2X_2 + \omega_3X_3)X_0^2 + \omega_4X_0^3.$$

The curve at infinity $\overline{\mathcal{X}} \setminus \mathcal{X}$ is the triangle composed of the three lines

$$L_k^\infty = \{X \in \mathbb{P}^3 : X_k = X_0 = 0\}, \quad (k = 1, 2, 3). \quad (5.4)$$

The gradient of F at $X_0 = 0$ is given by

$$\nabla F|_{X_0=0} = (\epsilon_1X_1^2 + \epsilon_2X_2^2 + \epsilon_3X_3^2, X_2X_3, X_1X_3, X_1X_2),$$

from which it immediately follows that the triangle of lines at infinity contains no singularities if and only if $\epsilon_1 = \epsilon_2 = \epsilon_3 = 1$. Furthermore, singularities can only be located at the three intersection points among the lines and, for any labelling $\{i, j, k\} = \{1, 2, 3\}$, the intersection point of L_i^∞ and L_j^∞ is a singularity if and only if $\epsilon_k = 0$.

Let us now focus on one of the intersection points of the three lines at infinity, say the point $S = [0 : 0 : 0 : 1]$. If $\epsilon_3 = 0$, then this is a singular point. We determine its type, in the notation by Arnol'd [1], following [5]. We start by introducing some inhomogeneous coordinates around S ,

$$[X_0 : X_1 : X_2 : X_3] = [u_1 : u_2 : u_3 : 1],$$

so that S corresponds to $u = (0, 0, 0)$. The equation for the cubic in these local coordinates reads

$$f(u) := u_2u_3 + \epsilon_1u_1u_2^2 + \epsilon_2u_1u_3^2 + \omega_1u_1^2u_2 + \omega_2u_1^2u_3 + \omega_3u_1^2 + \omega_4u_1^3 = 0.$$

Next, we apply a weighted scaling

$$u_k = r^{j_k} v_k \quad (1 \leq k \leq 3),$$

where r is a free scalar, and we look for triples $(j_1, j_2, j_3) \in \mathbb{Q}_{>0}^3$, such that

$$f(u) = r f_0(v) + o(r) \quad (r \rightarrow 0),$$

where the leading order coefficient $f_0(v)$ is such that $\{f_0(v) = 0\}$ has an isolated singularity at $v = (0, 0, 0)$. In such case, [5, Lemma 1] shows that the type of the singularity is given by

$$\text{type}(S) = \begin{cases} A_n & \text{if } (j_1, j_2, j_3) = (\frac{1}{n+1}, \frac{1}{2}, \frac{1}{2}), \quad (n \geq 1), \\ D_n & \text{if } (j_1, j_2, j_3) = (\frac{1}{n-1}, \frac{n-2}{2(n-1)}, \frac{1}{2}), \quad (n \geq 4), \\ E_6 & \text{if } (j_1, j_2, j_3) = (\frac{1}{3}, \frac{1}{4}, \frac{1}{2}), \\ E_7 & \text{if } (j_1, j_2, j_3) = (\frac{1}{3}, \frac{2}{9}, \frac{1}{2}), \\ E_8 & \text{if } (j_1, j_2, j_3) = (\frac{1}{3}, \frac{1}{5}, \frac{1}{2}). \end{cases}$$

We first consider the following values for the weights $(j_1, j_2, j_3) = (\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$, in which case

$$f(u) = r(v_2 v_3 + \omega_3 v_1^2) + o(r) \quad (r \rightarrow 0).$$

Therefore, as long as $\omega_3 \neq 0$, the leading order term has an isolated singularity at $v = (0, 0, 0)$, and S is a singularity of type A_1 .

Suppose now that $\omega_3 = 0$. Then we take the weights $(j_1, j_2, j_3) = (\frac{1}{3}, \frac{1}{2}, \frac{1}{2})$, so that

$$f(u) = r(v_2 v_3 + \omega_4 v_1^3) + o(r) \quad (r \rightarrow 0).$$

As long as $\omega_4 \neq 0$, the leading order term has an isolated singularity at $v = (0, 0, 0)$, and S is a singularity of type A_2 .

Next, suppose that also $\omega_4 = 0$. Then we take the weights $(j_1, j_2, j_3) = (\frac{1}{4}, \frac{1}{2}, \frac{1}{2})$, so that

$$f(u) = r(v_2 v_3 + \omega_1 v_1^2 v_2 + \omega_2 v_1^2 v_3) + o(r) \quad (r \rightarrow 0).$$

As long as $\omega_1 \omega_2 \neq 0$, the leading order term has an isolated singularity at $v = (0, 0, 0)$, and S is a singularity of type A_3 .

Next, suppose that also $\omega_1 \omega_2 = 0$. Without loss of generality, we consider the case $\omega_1 = 0$. The only admissible choice of weights is $(j_1, j_2, j_3) = (\frac{1}{5}, \frac{2}{5}, \frac{3}{5})$, and these weights do not allow us to read of the singularity type.

We therefore first apply a locally invertible polynomial mapping, $u \mapsto (u_1, u_2 - \omega_2 u_1^2, u_3)$, before scaling with the weights $(j_1, j_2, j_3) = (\frac{1}{5}, \frac{1}{2}, \frac{1}{2})$, yielding

$$f(u_1, u_2 - \omega_2 u_1^2, u_3) = r(v_2 v_3 + \epsilon_1 \omega_2^2 v_1^5) + o(r) \quad (r \rightarrow 0).$$

Now, necessarily $\epsilon_1 = 1$ since else the cubic is reducible. Therefore, as long as $\omega_2 \neq 0$, the leading order term has an isolated singularity at $v = (0, 0, 0)$, and S is a singularity of type A_4 . Similarly, if $\omega_2 = 0$ but $\omega_1 \neq 0$, then S is a singularity of type A_4 .

All in all, we have

$$\text{type}(S) = \begin{cases} - & \text{if } \epsilon_3 = 1, \\ A_1 & \text{if } \epsilon_3 = 0, \omega_3 \neq 0, \\ A_2 & \text{if } \epsilon_3 = 0, \omega_3 = 0, \omega_4 \neq 0, \\ A_3 & \text{if } \epsilon_3 = 0, \omega_3 = 0, \omega_4 = 0, \omega_1 \omega_2 \neq 0, \\ A_4 & \text{if } \epsilon_3 = 0, \omega_3 = 0, \omega_4 = 0, \omega_1 \omega_2 = 0, \omega_1 + \omega_2 \neq 0, \end{cases} \quad (5.5)$$

where $-$ signifies that S is a regular point. In the final case when $\omega_k = 0$, $1 \leq k \leq 4$, the singularity is non-isolated. Indeed, in that case, necessarily $\epsilon_1 = \epsilon_2 = 1$ (as otherwise the cubic surface is reducible) and its defining equation reads

$$X_1 X_2 X_3 + X_0 X_1^2 + X_0 X_2^2 = 0.$$

This surface is singular on the line $\{X_1 = X_2 = 0\}$ and has a further infinite number of lines lying in hyperplanes of the form $\{X_3 = t X_0\}$, $t \in \mathbb{C}$, which all intersect this line.

Returning to the general discussion, we note, in particular, that singularities of types A_5 , D_4 , D_5 , E_6 , \tilde{E}_6 cannot be realised as an intersection point in a triangle of lines on a cubic surface. These singularity types, however, do appear in cubic surfaces [5]. For example, the cubic surface

$$X_1 X_2 X_3 - X_1^3 - X_2^3 + X_0^2 X_1 = 0,$$

has an A_5 singularity at $[0 : 0 : 0 : 1]$. It contains only three lines,

$$\{X \in \mathbb{P}^3 : X_1 = X_2 = 0\},$$

$$\{X \in \mathbb{P}^3 : X_2 = 0, X_0 + X_1 = 0\},$$

$$\{X \in \mathbb{P}^3 : X_2 = 0, X_0 - X_1 = 0\},$$

which all intersect at this singularity. In particular, this singularity is not the corner of a triangle.

It follows from the above considerations, that a smooth affine cubic surface, with a triangle of lines at infinity, can only have singularities (in its projective completion) at the three intersection points of lines at infinity, and they can only be of type A_k , $1 \leq k \leq 4$.

By comparison with the classification of singular cubic surfaces, and their number of lines, in [5], we obtain a complete list of smooth affine cubic surfaces, with a triangle of lines at infinity, classified according to the singularity types of the corners. The result is given in Table 5.1. We discuss a few examples in this table.

Example 5.2 (A_1, A_1, A_3). By (5.5), the general form of a cubic surface with these singularities at infinity is

$$x_1 x_2 x_3 + \omega_1 x_1 + \omega_2 x_2 = 0, \quad \omega_1, \omega_2 \neq 0.$$

It has A_1 singularities at $[0 : 1 : 0 : 0]$ and $[0 : 0 : 1 : 0]$, an A_3 singularity at $[0 : 0 : 0 : 1]$, and no further singularities in its canonical projective completion. Apart from the three lines at infinity, there are two further lines, given by $\{x_1 = x_2 = 0\}$ and $\{x_1 + x_2 = x_3 = 0\}$. By scaling $x_1 \mapsto -\omega_2 x_1$ and $x_2 \mapsto -\omega_1 x_2$, we may normalise the cubic such that $\omega_1 = \omega_2 = -1$.

Example 5.3 ($-$, A_2, A_2). By (5.5), the general form of a cubic surface with these singularities at infinity is

$$x_1 x_2 x_3 + x_1^2 + \omega_1 x_1 + \omega_4 = 0, \quad \omega_4 \neq 0.$$

It has A_2 singularities at $[0 : 0 : 1 : 0]$ and $[0 : 0 : 0 : 1]$ in its projective completion, and no further singularities at infinity. Only when $\omega_1^2 = 4\omega_4$, the surface has a further finite singularity at $(x_1, x_2, x_3) = (-\frac{1}{2}\omega_1, 0, 0)$, of type A_1 . Apart from the three lines at infinity, there are generically four further lines. By scaling $x_{1,2} \mapsto \sqrt{\omega_4} x_{1,2}$, we may normalise the cubic such that $\omega_4 = 1$, yielding a one-parameter family of affine cubic surfaces.

Example 5.4 (A_1, A_1, A_2). By (5.5), the general form of the cubic surface is

$$x_1 x_2 x_3 + \omega_1 x_1 + \omega_2 x_2 + \omega_4 = 0, \quad \omega_1, \omega_2, \omega_4 \neq 0.$$

singularities	P-eqn	#lines	cubic
$-, -, -$	P_{VI}	24	$x_1x_2x_3 + x_1^2 + x_2^2 + x_3^2 + \omega_1x_1 + \omega_2x_2 + \omega_3x_3 + \omega_4$
$-, -, A_1$	P_V	18	$x_1x_2x_3 + x_1^2 + x_2^2 + \omega_1x_1 + \omega_2x_2 + \omega_3x_3 + R(\omega_{1,2,3})$
$-, -, A_2$	$P_{III}^{D_6}, P_V^{\text{deg}}$	12	$x_1x_2x_3 + x_1^2 + x_2^2 + \omega_1x_1 + \omega_2x_2 + \omega_1 - 1$
$-, -, A_3$	$P_{III}^{D_7}$	7	$x_1x_2x_3 + x_1^2 + x_2^2 + \omega_1x_1 - x_2$
$-, -, A_4$	$P_{III}^{D_8}$	3	$x_1x_2x_3 + x_1^2 + x_2^2 - x_2$
$-, A_1, A_1$	P_{IV}	13	$x_1x_2x_3 + x_1^2 + \omega_1x_1 + \omega_2(x_2 + x_3) + \omega_2(1 + \omega_1 - \omega_2)$
A_1, A_1, A_1	P_{II}^{JM}	9	$x_1x_2x_3 - x_1 + \omega_2x_2 - x_3 - \omega_2 + 1$
$-, A_1, A_2$	P_{II}^{FN}	8	$x_1x_2x_3 + x_1^2 + \omega_1x_1 - x_2 + 1$
A_1, A_1, A_2	P_I	5	$x_1x_2x_3 - x_1 - x_2 + 1$
$-, A_2, A_2$	-	4	$x_1x_2x_3 + x_1^2 + \omega_1x_1 + 1$
$-, A_1, A_3$	-	4	$x_1x_2x_3 + x_1^2 - x_1 - x_2$
A_1, A_1, A_3	-	2	$x_1x_2x_3 - x_1 - x_2$
A_1, A_2, A_2	-	2	$x_1x_2x_3 - x_1 + 1$
$-, A_1, A_4$	-	1	$x_1x_2x_3 + x_1^2 - x_2$
A_2, A_2, A_2	-	0	$x_1x_2x_3 + 1$

TABLE 5.1. Table of smooth embedded affine cubic surfaces, with a triangle of lines at infinity, listed according to the types of singularities (in their canonical projective completions) at the three intersection points of lines at infinity. In the first column the types of singularities respectively at $[0 : 1 : 0 : 0]$, $[0 : 0 : 1 : 0]$, $[0 : 0 : 0 : 1]$, where the symbol ‘-’ stands for a regular point. In the second column, the corresponding Painlevé equation(s), in the third column the number of affine lines and in the fourth column normal forms for the cubics. All the ω ’s are considered generic and in the second row, the rational function R is given by $R = 1 + \omega_3^2 - \frac{\omega_3(\omega_2 + \omega_1\omega_3)(\omega_1 + \omega_2\omega_3)}{(\omega_3^2 - 1)^2}$.

It has A_1 singularities at $[0 : 1 : 0 : 0]$ and $[0 : 0 : 1 : 0]$, an A_2 singularity at $[0 : 0 : 0 : 1]$, and no further singularities in its projective completion. Apart from the three lines at infinity, there are five further lines. By scaling

$$x_1 \mapsto -\frac{\omega_1}{\omega_4}x_1, \quad x_2 \mapsto -\frac{\omega_2}{\omega_4}x_2, \quad x_3 \mapsto \frac{\omega_1\omega_2}{\omega_4}x_3,$$

we may normalise the cubic such that $-\omega_1 = -\omega_2 = \omega_4 = 1$. This is the decorated character variety of P_I .

Example 5.5 $(-, -, A_2)$. By (5.5), the general form of the cubic surface is

$$x_1x_2x_3 + x_1^2 + x_2^2 + \omega_1x_1 + \omega_2x_2 + \omega_4 = 0, \quad \omega_4 \neq 0.$$

It has an A_2 singularity at $[0 : 0 : 0 : 1]$, and no further singularities in its projective completion, unless

$$(\omega_1^2 - 4\omega_4^2)(\omega_2^2 - 4\omega_4^2) = 0.$$

Apart from the three lines at infinity, there are generically 12 further lines. By scaling $x_{1,2} \mapsto u x_{1,2}$, where u is a root of $u^2 - 4\omega_1 + \omega_4 = 0$, we may normalise the surface such that $\omega_4 = \omega_1 - 1$, leading to the two-parameter family of cubic surfaces given in Table 5.1. This is the decorated character variety of $P_{III}^{D_6}$.

Example 5.6 $(-, A_1, A_1)$. By (5.5), the general form of the cubic surface is

$$x_1x_2x_3 + x_1^2 + \omega_1x_1 + \omega_2x_2 + \omega_3x_3 + \omega_4 = 0, \quad \omega_2, \omega_3, \omega_4 \neq 0.$$

It has A_1 singularities at $[0 : 0 : 1 : 0]$ and $[0 : 0 : 0 : 1]$, and no further singularities in its projective completion, unless

$$18 \omega_1 \omega_2 \omega_3 \omega_4 + 27 \omega_2^2 \omega_3^2 - \omega_1^2 \omega_4^2 + 4 \omega_4^3 - 4 \omega_1^3 \omega_2 \omega_3 = 0.$$

Apart from the three lines at infinity, there are generically 13 further lines. By scaling, we can ensure that $\omega_2 = \omega_3$ and $\omega_4 = \omega_2(1 + \omega_1 - \omega_2)$, leading to the two-parameter family of cubic surfaces given in Table 5.1. This is the decorated character variety of P_{IV} .

5.2. Constructing Segre surfaces. In Section 3.2.2, we showed how to blow down one of the lines at infinity of the cubic surface for P_{VI} , leading to an associated affine Segre surface. In this section, we consider this construction for general affine cubic surfaces with a triangle of lines at infinity.

Let us return to the general affine cubic surface $\mathcal{X} \subseteq \mathbb{C}^3$, with a triangle of lines at infinity, in normal form (5.1). For simplicity, let us further assume that \mathcal{X} has no finite singularities.

We focus on the construction of an affine Segre surface which, in the regular case, comes from a blow-down of the line L_1^∞ at infinity. Correspondingly introducing the variables

$$y_1 = x_1, \quad y_2 = x_2, \quad y_3 = x_3, \quad y_4 = x_2 x_3,$$

we obtain an affine Segre surface $\mathcal{Y} \subseteq \mathbb{C}^4$, given by

$$y_2 y_3 - y_4 = 0, \tag{5.6}$$

$$y_1 y_4 + \epsilon_1 y_1^2 + \epsilon_2 y_2^2 + \epsilon_3 y_3^2 + \omega_1 y_1 + \omega_2 y_2 + \omega_3 y_3 + \omega_4 = 0. \tag{5.7}$$

The polynomial mapping

$$\pi : \mathcal{X} \rightarrow \mathcal{Y}, x \mapsto y,$$

is an isomorphism between the affine varieties \mathcal{X} and \mathcal{Y} . Using homogeneous coordinates

$$[Y_0 : Y_1 : Y_2 : Y_3 : Y_4] = [1 : y_1 : y_2 : y_3 : y_4], \tag{5.8}$$

we define the canonical projective completion $\overline{\mathcal{Y}} \subseteq \mathbb{P}^4$ of \mathcal{Y} , by the homogeneous equations,

$$\begin{aligned} Y_2 Y_3 - Y_4 Y_0 &= 0, \\ Y_1 Y_4 + \epsilon_1 Y_1^2 + \epsilon_2 Y_2^2 + \epsilon_3 Y_3^2 + (\omega_1 Y_1 + \omega_2 Y_2 + \omega_3 Y_3) Y_0 + \omega_4 Y_0^2 &= 0. \end{aligned} \tag{5.9}$$

The curve at infinity, $\overline{\mathcal{Y}} \setminus \mathcal{Y}$, is described by

$$Y_1 Y_4 + \epsilon_1 Y_1^2 + \epsilon_2 Y_2^2 + \epsilon_3 Y_3^2 = 0, \quad Y_2 Y_3 = 0, \quad Y_0 = 0.$$

This quartic curve factorises into two quadratic curves,

$$C_2^\infty : \quad Y_1 Y_4 + \epsilon_1 Y_1^2 + \epsilon_3 Y_3^2 = 0, \quad Y_2 = 0, \quad Y_0 = 0,$$

and

$$C_3^\infty : \quad Y_1 Y_4 + \epsilon_1 Y_1^2 + \epsilon_2 Y_2^2 = 0, \quad Y_3 = 0, \quad Y_0 = 0,$$

which meet in two points,

$$[0 : 1 : 0 : 0 : -\epsilon_1], \quad q_1^\infty := [0 : 0 : 0 : 0 : 1].$$

The mapping π extends to a regular bi-rational mapping

$$\pi : \overline{\mathcal{X}} \rightarrow \overline{\mathcal{Y}}, \tag{5.10}$$

which is described on the three lines at infinity by

$$\begin{aligned} L_1^\infty : \quad & \pi([0 : 0 : X_2 : X_3]) = [0 : 0 : 0 : 0 : 1], \\ L_2^\infty : \quad & \pi([0 : X_1 : 0 : X_3]) = [0 : X_1^2 : 0 : X_1 X_3 : -(\epsilon_1 X_1^2 + \epsilon_3 X_3^2)], \\ L_3^\infty : \quad & \pi([0 : X_1 : X_2 : 0]) = [0 : X_1^2 : X_1 X_2 : 0 : -(\epsilon_1 X_1^2 + \epsilon_2 X_2^2)], \end{aligned}$$

see equations (5.4). In particular, $\pi(L_1^\infty) = \{q_1^\infty\}$ and $\pi(L_k^\infty) \subseteq C_k^\infty$ for $k = 2, 3$. Let us introduce some notation for the corner points of the triangle at infinity of the cubic,

$$\begin{aligned} p_{12}^\infty &= [0 : 0 : 0 : 1], \\ p_{13}^\infty &= [0 : 0 : 1 : 0], \\ p_{23}^\infty &= [0 : 1 : 0 : 0], \end{aligned}$$

so that p_{jk}^∞ is the intersection point of L_j^∞ and L_k^∞ for appropriate indices j, k , see Figure 5.1. Then π maps these corner points respectively to the following points on $\bar{\mathcal{Y}}$,

$$\begin{aligned} q_{12}^\infty &= [0 : 0 : 0 : 1 - \epsilon_3 : \epsilon_3], \\ q_{13}^\infty &= [0 : 0 : 1 - \epsilon_2 : 0 : \epsilon_2], \\ q_{23}^\infty &= [0 : 1 : 0 : 0 : -\epsilon_1]. \end{aligned}$$




FIGURE 5.1. Notation for lines and intersection points in triangle at infinity of the embedded affine cubic \mathcal{X} .

If $\epsilon_2 = \epsilon_3 = 1$, then both C_2^∞ and C_3^∞ are irreducible conics, $\pi(L_k^\infty) = C_k^\infty$ for $k = 2, 3$, and π is the blow-up of the Segre surface $\bar{\mathcal{Y}}$ at q_1^∞ , with exceptional divisor L_1^∞ .

If $\epsilon_2 = 0$, then C_3^∞ is the product of two lines

$$\begin{aligned} C_{3,a}^\infty : & \quad Y_4 + \epsilon_1 Y_1 = 0, & \quad Y_3 = 0, & \quad Y_0 = 0, \\ C_{3,b}^\infty : & \quad Y_1 = 0, & \quad Y_3 = 0, & \quad Y_0 = 0, \end{aligned}$$

which intersect at $q_{13}^\infty = [0 : 0 : 1 : 0 : 0]$, and $\pi(L_3^\infty) = C_{3,a}^\infty$.

If $\epsilon_3 = 0$, then C_2^∞ is the product of two lines

$$\begin{aligned} C_{2,a}^\infty : & \quad Y_4 + \epsilon_1 Y_1 = 0, & \quad Y_2 = 0, & \quad Y_0 = 0, \\ C_{2,b}^\infty : & \quad Y_1 = 0, & \quad Y_2 = 0, & \quad Y_0 = 0, \end{aligned}$$

which intersect at $q_{12}^\infty = [0 : 0 : 0 : 1 : 0]$, and $\pi(L_2^\infty) = C_{2,a}^\infty$.

We conclude that the quartic curve at infinity, is either the product of two irreducible conics, or the product of an irreducible conic and two lines, or the product of four lines. In Figure 5.2, the different cases are displayed.

Next, we consider singularities on $\bar{\mathcal{Y}}$. By assumption, \mathcal{X} is smooth, so \mathcal{Y} is smooth and thus singularities can only exist on the curve at infinity. The Jacobian of (5.9) with respect to Y , at $Y_0 = 0$, is given by

$$J_Y|_{Y_0=0} = \begin{bmatrix} Y_4 & 0 & Y_3 & Y_2 & 0 \\ \omega_1 Y_1 + \omega_2 Y_2 + \omega_3 Y_3 & Y_4 + 2\epsilon_1 Y_1 & 2\epsilon_2 Y_2 & 2\epsilon_3 Y_3 & Y_1 \end{bmatrix}.$$

A point Y at infinity is a singularity of $\overline{\mathcal{Y}}$ if and only if this Jacobian has rank less than two. It follows from this, that $\overline{\mathcal{Y}}$ can only have singularities at the intersection points of irreducible components of the curve at infinity. Note, furthermore, that q_1^∞ is always a regular point, and $q_{23}^\infty \in \overline{\mathcal{Y}}$ is a singular point if and only if $p_{23}^\infty \in \overline{\mathcal{X}}$ is a singular point in which case their types are the same.

Note that the Jacobian has rank less than two at q_{12}^∞ if and only if $\epsilon_3 = \omega_3 = 0$. In such case, we can determine the singularity type of q_{12}^∞ analogous to how we arrived at equation (5.5). We use local affine variables u defined through

$$[Y_0 : Y_1 : Y_2 : Y_3 : Y_4] = [u_1 : u_2 : u_1 u_3 : 1 : u_3],$$

so that q_{12}^∞ corresponds to $u = (0, 0, 0)$. The first equation in (5.9) is now trivially satisfied, and the second equation becomes

$$f(u) := u_2 u_3 + \epsilon_1 u_2^2 + \epsilon_2 u_1^2 u_3^2 + \omega_1 u_1 u_2 + \omega_2 u_1^2 u_3 + \omega_4 u_1^2 = 0.$$

The remainder of the procedure is the same as how we obtained equation (5.5). We apply a weighted scaling

$$u_k = r^{j_k} v_k \quad (1 \leq k \leq 3),$$

where r is a free scalar, and we look for triples $(j_1, j_2, j_3) \in \mathbb{Q}_{>0}^3$, such that a balance of overall weight 1 occurs.

Putting $(j_1, j_2, j_3) = (\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$, we have

$$f(u) = r(v_2 v_3 + \epsilon_1 v_2^2 + \omega_1 v_1 v_2 + \omega_4 v_1^2) + o(r) \quad (r \rightarrow 0).$$

As long as $\omega_4 \neq 0$, the leading order term has an isolated singularity at $v = (0, 0, 0)$, and q_{12}^∞ is a singularity of type A_1 .

Next, suppose that also $\omega_4 = 0$. Then we apply the locally invertible polynomial mapping $u_3 \mapsto u_3 - \omega_1 u_1$ and take weights $(j_1, j_2, j_3) = (\frac{1}{3}, \frac{1}{2}, \frac{1}{2})$, to obtain

$$f(u_1, u_2, u_3 - \omega_1 u_1) = r(v_2 v_3 + \epsilon_1 v_2^2 - \omega_1 \omega_2 v_1^3) + o(r) \quad (r \rightarrow 0).$$

As long as $\omega_1 \omega_2 \neq 0$, the leading order term has an isolated singularity at $v = (0, 0, 0)$, and q_{12}^∞ is a singularity of type A_2 .

Next, suppose that also $\omega_1 \omega_2 = 0$. Without loss of generality, we consider the case $\omega_1 = 0$. Taking $(j_1, j_2, j_3) = (\frac{1}{4}, \frac{1}{2}, \frac{1}{2})$, gives the balance

$$f(u) = r(v_2 v_3 + \epsilon_1 v_2^2 + \omega_2 v_1^2 v_3) + o(r) \quad (r \rightarrow 0).$$

Now, necessarily $\epsilon_1 = 1$ since else the Segre surface is reducible. Therefore, as long as $\omega_2 \neq 0$, the leading order term has an isolated singularity at $v = (0, 0, 0)$, and q_{12}^∞ is a singularity of type A_3 . Similarly, if $\omega_2 = 0$ but $\omega_1 \neq 0$, then q_{12}^∞ is a singularity of type A_3 .

All in all,

$$\text{type}(q_{12}^\infty) = \begin{cases} - & \text{if } \epsilon_3 = 1, \\ - & \text{if } \epsilon_3 = 0, \omega_3 \neq 0, \\ A_1 & \text{if } \epsilon_3 = 0, \omega_3 = 0, \omega_4 \neq 0, \\ A_2 & \text{if } \epsilon_3 = 0, \omega_3 = 0, \omega_4 = 0, \omega_1 \omega_2 \neq 0, \\ A_3 & \text{if } \epsilon_3 = 0, \omega_3 = 0, \omega_4 = 0, \omega_1 \omega_2 = 0, \omega_1 + \omega_2 \neq 0, \end{cases}$$

and in the final case when $\omega_k = 0$, $1 \leq k \leq 4$, the singularity is non-isolated and our running assumption that \mathcal{X} is smooth is violated. Comparing with (5.5), we see that $q_{12}^\infty \in \overline{\mathcal{Y}}$ is an A_{j-1} singularity when $p_{12}^\infty \in \overline{\mathcal{X}}$ is an A_j singularity, for $1 \leq j \leq 4$, where $A_0 := -$. Analogously, $q_{13}^\infty \in \overline{\mathcal{Y}}$ is an A_{i-1} singularity when $p_{13}^\infty \in \overline{\mathcal{X}}$ is an A_i singularity, $1 \leq i \leq 4$. The correspondence between singularities on the triangle at infinity of the cubic and singularities on the quartic curve at infinity of the Segre surface are summarised in Figure 5.2.

Finally, we discuss the relationship between affine lines on the cubic and affine lines on the Segre surface, in the following lemma.

Lemma 5.7. *Let L be any line on the cubic surface $\overline{\mathcal{X}}$, not at infinity. If L intersects only with the line L_1^∞ at infinity, then π maps L to a conic in $\overline{\mathcal{Y}}$. Otherwise, π maps L to a line in $\overline{\mathcal{Y}}$. All the lines in the Segre surface $\overline{\mathcal{Y}}$, not at infinity, arise in this way.*

Proof. Take any line L in the cubic surface $\overline{\mathcal{X}}$, not at infinity. Suppose that this line intersects only L_1^∞ at infinity. Then its affine part admits a parametrisation

$$x = ta + b \quad (t \in \mathbb{C}), \quad (5.11)$$

for some $a, b \in \mathbb{C}^3$, with $a_1 = 0$ and $a_2, a_3 \neq 0$. Therefore

$$\pi(L) = \{[t_0^2 : b_1 t_0^2 : a_2 t_1 t_0 + b_2 t_0^2 : a_3 t_1 t_0 + b_3 t_0^2 : (a_2 t_1 + b_2 t_0)(a_3 t_1 + b_3 t_0)] : [t_0 : t_1] \in \mathbb{P}^1\},$$

is a conic.

Otherwise, the affine part of the line admits a parametrisation (5.11), with either $a_2 = 0$ or $a_3 = 0$, and the image under π is thus a line, since $y_4 = y_2 y_3$ will be affine linear in t .


Conversely, take any line L in the Segre surface $\overline{\mathcal{Y}}$, not at infinity. The affine part of the line admits a parametrisation of the form

$$y = at + b,$$

for some $a, b \in \mathbb{C}^4$, with at least one $a_k \neq 0$, $1 \leq k \leq 3$, since $y_4 = y_2 y_3$. It follows that

$$\pi^{-1}(L) = \{[a_1 t_1 + b_1 t_0 : a_2 t_1 + b_2 t_0 : a_3 t_1 + b_3 t_0] : [t_0 : t_1] \in \mathbb{P}^1\},$$

is a line. In particular L is the image under π of a line in $\overline{\mathcal{X}}$ and the lemma follows. \square






FIGURE 5.2. The configurations of lines and singularities on the curves at infinity of the embedded affine cubic surface \mathcal{X} and embedded affine Segre surface \mathcal{Y} , for different choices of $\epsilon_2, \epsilon_3 \in \{0, 1\}$. Here the singularity type at each intersection point is depicted in purple, with corresponding indices satisfying $0 \leq k \leq 4$, $1 \leq i, j \leq 4$, and $A_0 := -$. The admissible values of (i, j, k) can be read off Table 5.1.

Remark 5.8. The classical Cayley–Salmon theorem [6, 44] states that any smooth cubic surface in \mathbb{P}^3 contains precisely 27 lines. Furthermore, it is well-known that any line in a smooth cubic surface intersects with precisely 10 others. Therefore, setting $\epsilon_{1,2,3} = 1$, it follows from the above lemma that the smooth Segre surface $\overline{\mathcal{Y}}$ contains precisely $27 - 10 - 1 = 16$ lines, as expected. This is also consistent with the line counting on the blow-up model in Remark 2.7.

Starting with any embedded affine cubic \mathcal{X} given by the normal form (5.1), we can carry out the above construction with respect to any of the three lines at infinity, leading, in general, to three inequivalent affine Segre surfaces. The results of this for the cubic surfaces corresponding to Painlevé equations are given in Table 5.2.

In the following sections, we will discuss some examples of this. For this purpose, and for general reference, we have provided Table 5.3 which details the possible singularity configurations and number of lines on Segre surfaces. We remark that the singularity configuration does not always determine the number of lines, and vice versa. For example, as follows from the table, there exist Segre surfaces with one singularity, of type A_3 , both with 5 lines and with 4 lines on them.

\mathcal{Y} singularities	P-eqn	L_1^∞	L_2^∞	L_3^∞
$-, -, -$	P_{VI}	$-(16, 0)$	$-(16, 0)$	$-(16, 0)$
$-, -, A_1$	P_V	$-(14, 2)$	$-(14, 2)$	$A_1(12, 0)$
$-, -, A_2$	$P_{III}^{D_6}, P_V^{\text{deg}}$	$A_1(10, 2)$	$A_1(10, 2)$	$A_2(8, 0)$
$-, -, A_3$	$P_{III}^{D_7}$	$A_2(6, 2)$	$A_2(6, 2)$	$A_3(5, 0)$
$-, -, A_4$	$P_{III}^{D_8}$	$A_3(3, 2)$	$A_3(2, 2)$	$A_4(3, 0)$
$-, A_1, A_1$	P_{IV}	$-(12, 4)$	$A_1(10, 2)$	$A_1(10, 2)$
A_1, A_1, A_1	P_{II}^{JM}	$A_1(8, 4)$	$A_1(8, 4)$	$A_1(8, 4)$
$-, A_1, A_2$	P_{II}^{FN}	$A_1(8, 4)$	$2A_1(7, 2)$	$A_2(6, 2)$
A_1, A_1, A_2	P_I	$2A_1(5, 4)$	$2A_1(5, 4)$	$A_2(4, 4)$

TABLE 5.2. Table summarising geometric data of embedded affine Segre surfaces obtained by applying the construction in Section 5.2 with respect to any of the three lines at infinity, for each of the Painlevé cubic surfaces, for generic parameter values. The rows represent the different Painlevé cubic surfaces and, in the three columns on the right, the line at infinity used in the construction is specified. Each entry takes the form ‘ $S, (m, n)$ ’, with S listing the singularities at infinity on the Segre surface, m the number of affine lines and n the number of lines at infinity.

$-$	A_1	A_2	A_3	A_3	A_4
16	12	8	5	4	3
$2A_1$	$2A_1$	$A_1 + A_2$	$A_1 + A_3$	$3A_1$	$2A_1 + A_2$
9	8	6	3	6	4
$2A_1 + A_3$	$4A_1$	D_4	D_5		
2	4	2	1		

TABLE 5.3. Possible singularity configurations and number of lines on Segre surfaces in $\mathbb{C}\mathbb{P}^4$, taken from Dolgachev [13, Table 8.6]. In each of the three blocks, the first row gives the singularities and the second row the number lines.

5.3. Affine cubic and Segre surfaces associated with P_{II} . In this section, we study the two affine cubic surfaces associated with P_{II} and corresponding Segre surfaces. In particular, we are going to use the construction in the last section to derive an explicit isomorphism between the two affine cubics. In particular, this shows that the \mathcal{Z} -Segre for P_{II}^{JM} is also a natural \mathcal{Z} -Segre for P_{II}^{FN} .

Consider the general P_{II}^{JM} cubic

$$x_1x_2x_3 - x_1 + \omega_2x_2 - x_3 - \omega_2 + 1 = 0. \quad (5.12)$$

It is smooth unless $\omega_2 = -1$ and reduces to the character variety of P_I as $\omega_2 \rightarrow 0$.

For generic ω_2 , the embedded cubic contains 9 affine lines, explicitly given by

$$\begin{aligned} L_1^{JM} &= \{x_1 + \omega_2 = 0, x_3 = 1\} & L_6^{JM} &= \{x_1 - 1 = 0, x_2 = 1\}, \\ L_2^{JM} &= \{x_3 + \omega_2 = 0, x_1 = 1\}, & L_7^{JM} &= \{x_3 - 1 = 0, x_2 = 1\}, \\ L_3^{JM} &= \{x_1 + \omega_2 = 0, x_2 = -\omega_2^{-1}\}, & L_8^{JM} &= \{x_1 = 0, x_3 - \omega_2x_2 = 1 - \omega_2\}, \\ L_4^{JM} &= \{x_3 + \omega_2 = 0, x_2 = -\omega_2^{-1}\}, & L_9^{JM} &= \{x_3 = 0, x_1 - \omega_2x_2 = 1 - \omega_2\}, \end{aligned}$$

$$L_5^{\text{JM}} = \{x_2 = 0, x_1 + x_3 = 1 - \omega_2\}.$$

We follow the construction in the previous section with respect to the line L_3^∞ at infinity and introduce variables

$$y_1 = x_1, \quad y_2 = x_2, \quad y_3 = x_3, \quad y_4 = x_1x_2,$$

to obtain the corresponding affine Segre surface $\mathcal{Y}^{\text{JM}} \subseteq \mathbb{C}^4$, defined by

$$y_1y_2 - y_4 = 0, \tag{5.13a}$$

$$y_3y_4 - y_1 + \omega_2y_2 - y_3 - \omega_2 + 1 = 0. \tag{5.13b}$$

We obtain the canonical projective completion $\bar{\mathcal{Y}}^{\text{JM}} \subseteq \mathbb{P}^4$ through homogeneous coordinates (5.8), and the curve at infinity is given by

$$Y_0 = 0, \quad Y_1Y_2 = 0, \quad Y_3Y_4 = 0.$$

This curve is the product of four lines, intersecting in four points, forming a rectangle, as in Figure 5.2d. The four intersection points are given by

$$p_k^{\text{JM}} : \quad Y_k = 1, \quad Y_j = 0 \quad (0 \leq j \leq 4, j \neq k),$$

where $1 \leq k \leq 4$. The point p_3^{JM} is an A_1 singularity and the other three are regular points in $\bar{\mathcal{Y}}^{\text{JM}}$. Note, furthermore, that $x \rightarrow y$ maps L_k^{JM} , $1 \leq k \leq 8$ to affine lines on \mathcal{Y}^{JM} , but L_9^{JM} is mapped to a conic. Thus $\bar{\mathcal{Y}}^{\text{JM}}$ is a Segre surface with one A_1 singularity and 12 lines, four of which lie at infinity, as indicated in the corresponding entry of Table 5.2.

Now, recall the affine Segre surface (4.6) of $\mathbb{P}_{\text{II}}^{\text{JM}}$ obtained by confluence in Section 4.2.4. It is affinely equivalent to \mathcal{Y}^{JM} under

$$\begin{aligned} z_1 &= \frac{\omega_2(y_4 - 1)}{1 + \omega_2}, & z_4 &= 1, \\ z_2 &= -1 - \frac{y_3}{\omega_2}, & z_5 &= \frac{y_1 - 1}{1 + \omega_2}, \\ z_3 &= \frac{1 - y_1 + \omega_2y_2 - \omega_2y_4}{1 + \omega_2}, & z_6 &= -\frac{1 + \omega_2y_2}{1 + \omega_2}, \end{aligned}$$

with parameter values in (4.6) given by $\rho_4 = \lambda_1 = 1$ and $\lambda_2 = 1 + \omega_2$.

Next, we turn our attention to the embedded affine cubic surface of $\mathbb{P}_{\text{II}}^{\text{FN}}$, defined by

$$\tilde{x}_1\tilde{x}_2\tilde{x}_3 + \tilde{x}_1^2 + \omega_1\tilde{x}_1 - \tilde{x}_2 + 1 = 0. \tag{5.14}$$

It is smooth unless $\omega_1^2 = 4$. It will be helpful to write $\omega_1 = -(v_0 + v_0^{-1})$, so that the 8 affine lines on the cubic admit a simple description,

$$\begin{aligned} L_1^{\text{FN}} &= \{\tilde{x}_1 = v_0, \tilde{x}_3 = v_0^{-1}\}, & L_5^{\text{FN}} &= \{x_2 = 1, \tilde{x}_1 + \tilde{x}_3 = v_0 + v_0^{-1}\}, \\ L_2^{\text{FN}} &= \{\tilde{x}_3 = v_0, \tilde{x}_1 = v_0^{-1}\}, & L_6^{\text{FN}} &= \{\tilde{x}_1 = v_0^{-1}, \tilde{x}_2 = 0\}, \\ L_3^{\text{FN}} &= \{\tilde{x}_1 = v_0, \tilde{x}_2 = 0\}, & L_7^{\text{FN}} &= \{v_0\tilde{x}_1 + \tilde{x}_2 = 1, \tilde{x}_3 = v_0^{-1}\}, \\ L_4^{\text{FN}} &= \{\tilde{x}_3 = v_0, \tilde{x}_1 + v_0\tilde{x}_2 = v_0\}, & L_8^{\text{FN}} &= \{\tilde{x}_1 = 0, \tilde{x}_2 = 1\}. \end{aligned}$$

Next, we consider the construction Section 3.2.2 with respect to the line L_1^∞ at infinity, so that the resulting affine Segre surface has only an A_1 singularity in its canonical projective completion, like $\bar{\mathcal{Y}}^{\text{JM}}$. So, we introduce the variables

$$\tilde{y}_1 = \tilde{x}_1, \quad \tilde{y}_2 = \tilde{x}_2, \quad \tilde{y}_3 = \tilde{x}_3, \quad \tilde{y}_4 = \tilde{x}_2\tilde{x}_3,$$

to obtain the affine Segre surface $\mathcal{Y}^{\text{FN}} \subseteq \mathbb{C}^4$ defined by

$$\tilde{y}_2\tilde{y}_3 - \tilde{y}_4 = 0,$$

$$\tilde{y}_1\tilde{y}_4 + \tilde{y}_1^2 + \omega_1\tilde{y}_1 - \tilde{y}_2 + 1 = 0.$$

Its canonical projective completion $\overline{\mathcal{Y}}^{\text{FN}} \subseteq \mathbb{P}^4$ through homogeneous coordinates \tilde{Y} , see equation (5.8), and the curve at infinity is given by

$$\tilde{Y}_0 = 0, \quad \tilde{Y}_2\tilde{Y}_3 = 0, \quad \tilde{Y}_1(\tilde{Y}_1 + \tilde{Y}_4) = 0.$$

This curve is also the product of four lines, intersecting in four points, forming a rectangle, as in Figure 5.2d. The four intersection points are given by

$$\begin{aligned} p_1^{\text{FN}} &= [0 : 1 : 0 : 0 : -1], \\ p_2^{\text{FN}} &= [0 : 0 : 0 : 0 : 1], \\ p_3^{\text{FN}} &= [0 : 0 : 0 : 1 : 0], \\ p_4^{\text{FN}} &= [0 : 0 : 1 : 0 : 0], \end{aligned}$$

where $1 \leq k \leq 4$. The point p_3^{FN} is an A_1 singularity and the other three are regular points in $\overline{\mathcal{Y}}^{\text{FN}}$. Note, furthermore, that $x \rightarrow y$ maps each of the affine lines L_k^{FN} , $1 \leq k \leq 8$, to an affine line on \mathcal{Y}^{FN} .

Since \mathcal{Y}^{JM} and \mathcal{Y}^{FN} are both smooth affine Segre surfaces with a ‘rectangle of lines’ at infinity in their canonical projective completions, with one A_1 singularity, it is natural to ask whether they might be affinely equivalent. We thus look for an $A \in GL_5(\mathbb{C})$, such that the mapping

$$\mathbb{P}^4 \rightarrow \mathbb{P}^4, Y \mapsto \tilde{Y} = AY,$$

maps \mathcal{Y}^{JM} onto \mathcal{Y}^{FN} . We require that the curves at infinity are mapped to one another, and to this end we impose that the intersection point p_k^{JM} is mapped to p_k^{FN} , $1 \leq k \leq 4$. As a result, A takes the form

$$A = \begin{bmatrix} a_{11} & 0 & 0 & 0 & 0 \\ a_{21} & a_{22} & 0 & 0 & 0 \\ a_{31} & 0 & 0 & 0 & a_{35} \\ a_{41} & 0 & 0 & a_{44} & 0 \\ a_{51} & -a_{22} & a_{53} & 0 & 0 \end{bmatrix}.$$

where we may scale A such that $a_{11} = 1$. By direct substitution of

$$\tilde{y} = A \cdot (1, y_1, y_2, y_3, y_4)^T, \quad (5.15)$$

into the equations defining \mathcal{Y}^{FN} , and simplification modulo the equations defining \mathcal{Y}^{JM} , we obtain an overdetermined linear system, from which the remaining coefficients of A are determined,

$$A = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & \frac{1-\omega_2}{\omega_1\omega_2} & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & \frac{1-\omega_2}{\omega_1\omega_2} & 0 \\ -\omega_1 & \frac{\omega_2-1}{\omega_1\omega_2} & \frac{1-\omega_2}{\omega_1} & 0 & 0 \end{bmatrix},$$

as well as a remaining necessary and sufficient condition relating ω_1 and ω_2 , namely

$$\omega_1^2 + 2 = \omega_2 + \omega_2^{-1}. \quad (5.16)$$

In other words, when (5.16) holds, then (5.15) defines an affine transformation that induces an isomorphism from \mathcal{Y}^{JM} to \mathcal{Y}^{FN} . As a consequence, we obtain the following result.

Proposition 5.9. *The affine cubic surface of P_{II}^{JM} , defined by equation (5.12), and the affine cubic surface of P_{II}^{FN} , defined by equation (5.14), are isomorphic when their parameters are related by (5.16). Using the rational parametrisation*

$$\omega_1 = -(v_0 + v_0^{-1}), \quad \omega_2 = -v_0^2, \quad v_0 \in \mathbb{C}^*,$$

an explicit isomorphism is given by

$$\tilde{x}_1 = v_0^{-1}x_1, \quad \tilde{x}_2 = 1 - x_1x_2, \quad \tilde{x}_3 = v_0^{-1}x_3,$$

with inverse

$$x_1 = v_0\tilde{x}_1, \quad x_2 = 1 + v_0^{-2} - v_0^{-1}(\tilde{x}_1 + \tilde{x}_2\tilde{x}_3), \quad x_3 = v_0\tilde{x}_3.$$

Proof. From equation (5.15), we obtain the affine transformation

$$\tilde{y}_1 = v_0^{-1}y_1, \quad \tilde{y}_2 = 1 - y_4, \quad \tilde{y}_3 = v_0^{-1}y_3, \quad \tilde{y}_4 = v_0 + v_0^{-1} - v_0^{-1}y_1 - v_0y_2,$$

which induces an isomorphism from \mathcal{Y}^{JM} to \mathcal{Y}^{FN} . Now, note that the mapping $x \mapsto y$ from the affine cubic of \mathcal{Y}^{JM} to \mathcal{Y}^{JM} is an isomorphism of affine varieties, and similarly $\tilde{x} \mapsto \tilde{y}$ is from the affine cubic of \mathcal{Y}^{FN} to \mathcal{Y}^{FN} is an isomorphism. Composing the three isomorphisms, $x \mapsto y$, $y \mapsto \tilde{y}$ and $\tilde{y} \mapsto \tilde{x}$, yields the isomorphism between the affine cubic surfaces of \mathcal{Y}^{JM} and \mathcal{Y}^{FN} in the proposition. \square

Remark 5.10. The isomorphism in Proposition 5.9 maps the line L_k^{FN} to L_k^{JM} , $1 \leq k \leq 8$. Furthermore, the inverse image of the remaining affine line, L_9^{JM} , is a conic in \mathcal{Y}^{FN} .

Remark 5.11. We note that a bi-rational mapping between the cubic surfaces of P_{II}^{FN} and P_{II}^{JM} was given in [9, Remark 2.1]. That mapping is singular along $x_2x_3 = 0$ and does not provide a global isomorphism as in Proposition 5.9.

We further obtain an affine equivalence between the \mathcal{Z} -Segre listed in Table 1.1, with parameter value $\lambda_2 = 1 - v_0^2$, and the Segre surface \mathcal{Y}^{FN} with $\omega_1 = -(v_0 + v_0^{-1})$, given by

$$\begin{aligned} z_1 &= \frac{v_0^2 \tilde{y}_2}{1 - v_0^2}, & z_4 &= 1, \\ z_2 &= \frac{\tilde{y}_3}{v_0} - 1, & z_5 &= \frac{v_0 \tilde{y}_1 - 1}{1 - v_0^2}, \\ z_3 &= \frac{v_0(\tilde{y}_4 - v_0 \tilde{y}_2)}{1 - v_0^2}, & z_6 &= \frac{v_0(v_0 - \tilde{y}_1 - \tilde{y}_4)}{1 - v_0^2}. \end{aligned}$$

5.4. Cubic and Segre surfaces associated with P_I . Consider the general embedded affine cubic for P_I given by

$$x_1x_2x_3 - x_1 - x_2 + 1 = 0.$$

It is smooth and contains 5 affine lines, given by

$$\begin{aligned} L_1^I &= \{x_1 = 0, x_2 = 1\} & L_4^I &= \{x_2 = 1, x_3 = 1\}, \\ L_2^I &= \{x_1 = 1, x_2 = 0\}, & L_5^I &= \{x_3 = 0, x_1 + x_2 = 1\}, \\ L_3^I &= \{x_1 = 1, x_3 = 1\}. \end{aligned}$$

We follow the construction in the Section 5.2 with respect to the line L_3^∞ at infinity and introduce variables

$$y_1 = x_1, \quad y_2 = x_2, \quad y_3 = x_3, \quad y_4 = x_1x_2,$$

to obtain the corresponding affine Segre surface $\mathcal{Y}^I \subseteq \mathbb{C}^4$, defined by

$$\begin{aligned} y_1y_2 - y_4 &= 0, \\ y_3y_4 - y_1 - y_2 + 1 &= 0. \end{aligned}$$

We obtain the canonical projective completion $\overline{\mathcal{Y}}^I \subseteq \mathbb{P}^4$ through homogeneous coordinates (5.8), and the curve at infinity is given by

$$Y_0 = 0, \quad Y_1 Y_2 = 0, \quad Y_3 Y_4 = 0.$$

This curve is the product of four lines, intersecting in four points, forming a rectangle, as in Figure 5.2d. The four intersection points are given by

$$p_k^I: \quad Y_k = 1, \quad Y_j = 0 \quad (0 \leq j \leq 4, j \neq k),$$

where $1 \leq k \leq 4$. The point p_3^I is an A_2 singularity and the other three are regular points in $\overline{\mathcal{Y}}^I$. Note, furthermore, that $x \rightarrow y$ maps L_k^I , $1 \leq k \leq 4$ to affine lines on \mathcal{Y}^I , but L_5^I is mapped to a conic. Thus $\overline{\mathcal{Y}}^I$ is a Segre surface with one A_2 singularity and 8 lines, four of which lie at infinity, as indicated in the corresponding entry of Table 5.2. Setting

$$\begin{aligned} z_1 &= -y_4, & z_4 &= 1, \\ z_2 &= y_3 - 1, & z_5 &= y_1 - 1, \\ z_3 &= 1 - y_1 - y_2 + y_4, & z_6 &= y_2 - 1, \end{aligned}$$

and vice-versa

$$y_1 = z_5 + 1, \quad y_2 = z_6 + 1, \quad y_3 = z_2 + 1, \quad y_4 = -z_1,$$

we obtain that \mathcal{Y}^I is isomorphic to the following \mathcal{Z} -Segre:

$$z_1 + z_3 + z_4 + z_5 + z_6 = 0 \tag{5.17a}$$

$$z_4 - 1 = 0 \tag{5.17b}$$

$$z_3 z_4 - z_1 z_2 = 0 \tag{5.17c}$$

$$z_3 z_4 - z_5 z_6 = 0. \tag{5.17d}$$

5.5. Cubic and Segre surfaces associated with $P_V^{(deg)}$. Looking at Table 4.1, we see that the $P_{III}^{D_6}$ and the $P_V^{(deg)}$ coincide, even though the parameters v_i and ν_j in the two cases don't. Denoting the $P_{III}^{D_6}$ variables and parameters with the index III and the ones of $P_V^{(deg)}$ with an index V_a is easy to see that setting

$$\begin{aligned} x_1^{V_a} &= \frac{x_1^{III}}{\sqrt{v_0^{III} v_\infty^{III}}}, & x_2^{V_a} &= \frac{x_2^{III}}{\sqrt{v_0^{III} v_\infty^{III}}}, & x_3^{V_a} &= x_3^{III}, \\ v_0^{V_a} &= \sqrt{v_0^{III} v_\infty^{III}}, & v_t^{V_a} &= \sqrt{\frac{v_\infty^{III}}{v_0^{III}}} \end{aligned}$$

we map the $P_V^{(deg)}$ cubic to the $P_{III}^{D_6}$ one. This shows that the two \mathcal{Z} -Segre surfaces coincide.

We can also produce the P_V^{deg} by blowing down the P_V^{deg} cubic

$$x_1 x_2 x_3 + x_1^2 + x_2^2 + \omega_1 x_1 + \omega_2 x_2,$$

at the line $X_0 = X_3 = 0$. By an linear affine transformation (see mathematica file), this leads to

$$z_1 + z_3 + z_4 + z_5 + z_6 = 0 \tag{5.18a}$$

$$\rho_3 z_3 + z_4 + \rho_5 z_5 + \rho_6 z_6 = 0 \tag{5.18b}$$

$$z_3 z_4 - z_1 z_2 = 0 \tag{5.18c}$$

$$z_3 z_4 - z_5 z_6 = 0, \tag{5.18d}$$

where $\rho_6 = \frac{\rho_3}{\rho_5}$. The family (5.18) only depends on two independent parameters.

5.6. **Cubic and Segre surfaces associated with $\mathbf{P}_{\text{III}}^{D_7}$.** The affine cubic surface associated with $\mathbf{P}_{\text{III}}^{D_7}$ is defined by

$$x_1x_2x_3 + x_1^2 + x_2^2 + \omega_1x_1 - x_2 = 0,$$

with $\omega_1 \in \mathbb{C}^*$. It is smooth for any value of ω_1 and reduces to the cubic surface for $\mathbf{P}_{\text{III}}^{D_8}$ as $\omega_1 \rightarrow 0$.

The embedded surface contains 7 affine lines, given by

$$\begin{aligned} L_1 &= \{x_1 = 0, x_2 = 0\}, & L_5 &= \{x_3 + \omega_1 + \omega_1^{-1} = 0, \omega_1x_1 - x_2 = 0\}, \\ L_2 &= \{x_1 = 0, x_2 = 1\}, & L_6 &= \{x_3 + \omega_1 + \omega_1^{-1} = 0, x_1 - \omega_1x_2 + \omega_1 = 0\}, \\ L_3 &= \{x_1 + \omega_1 = 0, x_2 = 0\}, & L_7 &= \{x_1 + \omega_1 = 0, 1 - x_2 + \omega_1x_3 = 0\}, \\ L_4 &= \{x_2 = 1, x_1 + x_3 + \omega_1 = 0\}, \end{aligned}$$

and three lines at infinity.

We follow the construction in Section 5.2 with respect to the line L_1^∞ at infinity and introduce variables

$$y_1 = x_1, \quad y_2 = x_2, \quad y_3 = x_3, \quad y_4 = x_2x_3,$$

to obtain the corresponding affine Segre surface $\mathcal{Y} \subseteq \mathbb{C}^4$, defined by

$$\begin{aligned} y_2y_3 - y_4 &= 0, \\ y_1y_4 + y_1^2 + y_2^2 + \omega_1y_1 - y_2 &= 0. \end{aligned}$$

We obtain the canonical projective completion $\overline{\mathcal{Y}} \subseteq \mathbb{P}^4$ through homogeneous coordinates (5.8), and the curve at infinity is given by

$$Y_0 = 0, \quad Y_2Y_3 = 0, \quad Y_1(Y_1 + Y_4) = 0.$$

This curve is the product of two lines and a conic, intersecting in three points, see Figure 5.2c. The point where the two lines at infinity intersect,

$$[Y_0 : Y_1 : Y_2 : Y_3 : Y_4] = [0 : 0 : 0 : 1 : 0],$$

is an A_2 singularity, and $\overline{\mathcal{Y}}$ is smooth elsewhere.

Note, furthermore, that $x \rightarrow y$ maps the lines L_k , $1 \leq k \leq 6$ to affine lines on \mathcal{Y} , but L_7 is mapped to a conic. Thus $\overline{\mathcal{Y}}$ is a Segre surface with one A_2 singularity and 8 lines, two of which lie at infinity, as indicated in the corresponding entry of Table 5.2. Setting

$$\begin{aligned} z_1 &= \omega_1^{-1}y_1 + y_2 + \omega_1^{-1}y_4, & z_4 &= 1 - y_2, \\ z_2 &= -1 - \omega_1^{-1}y_1 + y_2, & z_5 &= \omega_1^{-2}y_2, \\ z_3 &= -((1 + \omega_1^{-2})y_2 + \omega_1^{-1}y_4), & z_6 &= -1 + (1 + \omega_1^2)(y_2 - 1) - \omega_1y_3 + \omega_1y_4, \end{aligned}$$

we obtain the corresponding \mathcal{Z} -Segre surface

$$\begin{aligned} z_1 + z_2 + z_3 + z_4 + z_5 &= 0, \\ z_4 + \omega_1^2z_5 - 1 &= 0, \\ z_3z_4 - z_1z_2 &= 0, \\ z_3z_4 - z_5z_6 &= 0. \end{aligned}$$

This is the entry in Table 1.1 for $\mathbf{P}_{\text{III}}^{D_7}$ with $\rho_5 = \omega_1^2$.

6. POISSON STRUCTURE

Any algebraic variety defined as zero set of $n - 2$ polynomials in n variables is endowed by a natural Poisson bracket that was introduced by Nambu [33].

In this section, we study the Poisson brackets on the family of Segre surfaces \mathcal{Z}_q . In the case of the Painlevé differential equations, for each equation, there are three surfaces: namely the monodromy manifold, its blow down to the \mathcal{Y} -Segres and the isomorphic \mathcal{Z} -Segre. Here, we show that this blow down and isomorphism are in fact Poisson maps.

6.1. qPVI. For the Segre surface \mathcal{Z}_q the Nambu Poisson bracket is defined by the following formula

$$\{f, g\}_{\mathcal{Z}} := \frac{df \wedge dg \wedge dh_1 \wedge dh_2 \wedge dh_3 \wedge dh_4}{dz_1 \wedge dz_2 \wedge dz_3 \wedge dz_4 \wedge dz_5 \wedge dz_6}. \quad (6.1)$$

Applying this formula to h_1, h_3, h_4 defined by (2.2), and h_2'' defined in (2.3) in (6.1), we obtain the following:

$$\begin{aligned} \{z_1, z_2\}_{\mathcal{Z}} &= \lambda_2(z_3(z_5(-\rho_3 + \rho_5) + z_6(\rho_3 - \rho_6)) + z_4(z_5 - z_5\rho_5 + z_6(-1 + \rho_6))), \\ \{z_1, z_3\}_{\mathcal{Z}} &= z_3\lambda_2(z_5(\rho_2 - \rho_5) - z_6(\rho_2 - \rho_6)) + z_1\lambda_1(z_5\lambda_2(1 - \rho_5) - z_6\lambda_2(1 - \rho_6) + \\ &\quad + z_3(\rho_6 - \rho_5)), \\ \{z_1, z_4\}_{\mathcal{Z}} &= z_1\lambda_1\lambda_2(z_5(\rho_5 - \rho_3) - z_6(\rho_6 - \rho_3)) + z_4(z_5\lambda_2(\rho_5 - \rho_2) + z_6\lambda_2(\rho_2 - \rho_6) + \\ &\quad + z_1\lambda_1(\rho_5 - \rho_6)), \\ \{z_1, z_5\}_{\mathcal{Z}} &= z_5\lambda_2(z_4(\rho_2 - 1) + z_3(\rho_3 - \rho_2)) + z_1\lambda_1(z_5\lambda_2(\rho_3 - 1) + z_3(\rho_3 - \rho_6) + \\ &\quad z_4(\rho_6 - 1)), \\ \{z_1, z_6\}_{\mathcal{Z}} &= z_6\lambda_2(z_4 - z_4\rho_2 + z_3(\rho_2 - \rho_3)) + z_1\lambda_1(z_4 + z_6\lambda_2 - z_3\rho_3 - z_6\lambda_2\rho_3 + \\ &\quad z_3\rho_5 - z_4\rho_5), \\ \{z_2, z_3\}_{\mathcal{Z}} &= z_3\lambda_2(z_5\rho_5 - z_6\rho_6) + z_2\lambda_1(z_6\lambda_2 + z_5\lambda_2(\rho_5 - 1) + z_3\rho_5 - z_3\rho_6 - z_6\lambda_2\rho_6), \\ \{z_2, z_4\}_{\mathcal{Z}} &= z_4\lambda_2(z_6\rho_6 - z_5\rho_5) + z_2\lambda_1(z_5\lambda_2(\rho_3 - \rho_5) + z_6\lambda_2(\rho_6 - \rho_3) + z_4(\rho_6 - \rho_5)), \\ \{z_2, z_5\}_{\mathcal{Z}} &= z_5\lambda_2(z_4 - z_3\rho_3) + z_2\lambda_1(z_4 + z_5\lambda_2 - z_3\rho_3 - z_5\lambda_2\rho_3 + z_3\rho_6 - z_4\rho_6), \\ \{z_2, z_6\}_{\mathcal{Z}} &= z_6\lambda_2(z_3\rho_3 - z_4) + z_2\lambda_1(z_6\lambda_2(\rho_3 - 1) + z_3(\rho_3 - \rho_5) + z_4(\rho_5 - 1)), \\ \{z_3, z_4\}_{\mathcal{Z}} &= \lambda_1\lambda_2(z_2z_5(-\rho_2 + \rho_5) + z_2z_6(\rho_2 - \rho_6) + z_1(z_6\rho_6 - z_5\rho_5)), \\ \{z_3, z_5\}_{\mathcal{Z}} &= z_3z_5\lambda_2\rho_2 + z_2\lambda_1(z_5\lambda_2(\rho_2 - 1) + z_3(\rho_2 - \rho_6)) + z_1\lambda_1(z_5\lambda_2 + z_3\rho_6), \\ \{z_3, z_6\}_{\mathcal{Z}} &= -z_3z_6\lambda_2\rho_2 - z_1\lambda_1(z_6\lambda_2 + z_3\rho_5) + z_2\lambda_1(z_6\lambda_2 - z_3\rho_2 - z_6\lambda_2\rho_2 + z_3\rho_5), \\ \{z_4, z_5\}_{\mathcal{Z}} &= z_4z_6\lambda_2\rho_2 + z_1z_6\lambda_1\lambda_2\rho_3 + z_2\lambda_1(z_6\lambda_2(\rho_2 - \rho_3) + z_4(\rho_2 - \rho_5)) + z_1z_4\lambda_1\rho_5, \\ \{z_5, z_6\}_{\mathcal{Z}} &= \lambda_1(z_2(z_4 - z_4\rho_2 + z_3(\rho_2 - \rho_3)) + z_1(-z_4 + z_3\rho_3)), \end{aligned}$$

By construction, this bracket defines a Poisson bracket (namely it satisfies the Jacobi identity [46]) on $\mathbb{C}[z_1, \dots, z_6]$ and the functions h_1, h_2'', h_3, h_4 are central elements. This means that it can be restricted to the Segre surface

$$\mathcal{Z}_q = \text{Spec}(C[z_1, \dots, z_6]/\langle h_1, h_2'', h_3, h_4 \rangle).$$

Lemma 6.1. *The central elements of the Poisson bracket (6.2) are algebraically dependent on h_1, \dots, h_4 .*

Proof. The proof of this statement can be extracted from Section 3 of [37]. In our case the proof can also be done directly by observing that thanks to the proof of Proposition 2.4, the Jacobian matrix of h_1, \dots, h_4 has maximal rank 4. Then

because any f is central element must satisfy $\{f, z_i\}_{\mathcal{Z}} = 0$ for all $i = 1, \dots, 6$, it is easy to prove that

$$\frac{\partial f}{\partial z_j} = \sum_{i=1}^4 \alpha_i \frac{\partial h_i}{\partial z_j}, \quad j = 1, 2, \dots, 6,$$

for some polynomials $\alpha_1, \dots, \alpha_4$ that are constant on the Segre surface. We conclude that f must be algebraically dependent on h_1, \dots, h_4 . \square

6.2. Poisson structure on the monodromy manifolds of the differential equations and their blow down. In this subsection we prove the following

Lemma 6.2. *For each Painlevé differential equation, the blow down of its monodromy manifold $\mathcal{X}^{(d)}$, defined as zero set of the polynomial $\phi^{(d)}$ defined in (4.2), is a Poisson map.*

Proof. Observe that $\mathcal{X}^{(d)}$ admits the natural Poisson bracket defined by:

$$\{x_1, x_2\}_x = \frac{\partial \phi^{(d)}}{\partial x_3}, \quad \{x_2, x_3\}_x = \frac{\partial \phi^{(d)}}{\partial x_1}, \quad \{x_3, x_1\}_x = \frac{\partial \phi^{(d)}}{\partial x_2}. \quad (6.3)$$

Let us choose the following blow down:

$$\mathcal{Y}^{(d)} := \{(y_1, y_2, y_3, y_4) \in \mathbb{C}^4 \mid y_4 - y_2 y_3, y_1 y_4 + \sum_{i=1}^3 (\epsilon_i^{(d)} i^2 + \omega_i^{(d)} y_i) + \omega_4^{(d)} = 0\}. \quad (6.4)$$

Then, the Nambu Poisson bracket on the surface $\mathcal{Y}^{(d)}$ is defined as follows

$$\{f, g\}_y = \frac{df \wedge dg \wedge d\Phi_1^{(d)} \wedge d\Psi_1}{dy_1 \wedge dy_2 \wedge dy_3 \wedge dy_4}.$$

Therefore, for $i < j$,

$$\{y_i, y_j\}_y = \begin{bmatrix} 0 & -y_3 & -y_2 & 1 \\ y_4 + 2\epsilon_1^{(d)} y_1 + \omega_1^{(d)} & 2\epsilon_2^{(d)} y_2 + \omega_2^{(d)} & 2\epsilon_3^{(d)} y_3 + \omega_3^{(d)} & y_1 \end{bmatrix}_{ij}, \quad (6.5)$$

where $[\]_{ij}$ denotes the determinant of the matrix obtained erasing the i -th and the j -th columns. Using this formula, it is a straightforward computation to prove that

$$\{y_i, y_j\}_y = -\{y_i(x_1, x_2, x_3), y_j(x_1, x_2, x_3)\}_{\mathcal{X}},$$

hence proving that the blow down map is Poisson. \square

6.2.1. Poisson bracket on the \mathcal{Z} -Segre. In this subsection we prove the following

Proposition 6.3. *For each Painlevé differential equation, the linear affine transformation*

$$\Phi^{(d)} : \mathcal{Y}^{(d)} \rightarrow \mathcal{Z}^{(d)},$$

where d is an index that runs through the list of the Painlevé equations, is Poisson with Poisson inverse.

Proof. We already know that for each Painlevé equation $\Phi^{(d)}$ is invertible. We only need to prove that Poisson relations are sent to Poisson relations. We start by observing that the Poisson structure on the Segre \mathcal{Z}_1 is defined in the same way as for the \mathcal{Z}_q , namely by (6.1). The properties of this bracket are the same for all

members of Table 1.1, so that we obtain a unified formula that by specializing the parameters $\epsilon_i^{(d)}$ and $\rho_i^{(d)}, \lambda_i^{(d)}$ gives the Poisson bracket on $\mathcal{Z}^{(d)}$:

$$\begin{aligned}
\{z_1, z_2\}_{\mathcal{Z}} &= \lambda_2^{(d)} z_3 \left(z_5 (\rho_5^{(d)} - \rho_3^{(d)}) + z_6 (\epsilon_6^{(d)} \rho_3^{(d)} - \epsilon_4^{(d)} \rho_6^{(d)}) \right) \\
&\quad + \lambda_2^{(d)} z_4 \left(z_5 (1 - \epsilon_4^{(d)} \rho_5^{(d)}) + z_6 (\rho_6^{(d)} - \epsilon_6^{(d)}) \right), \\
\{z_1, z_3\}_{\mathcal{Z}} &= z_1 \lambda_1^{(d)} \left(z_5 \lambda_2^{(d)} (1 - \epsilon_4^{(d)} \rho_5^{(d)}) + z_3 (\rho_6^{(d)} - \epsilon_6^{(d)} \rho_5^{(d)}) + \right. \\
&\quad \left. + z_6 \lambda_2^{(d)} (\epsilon_4^{(d)} \rho_6^{(d)} - \epsilon_6^{(d)}) \right) + z_3 \epsilon_2^{(d)} \lambda_2^{(d)} (\rho_6^{(d)} z_6 - \rho_5^{(d)} z_5), \\
\{z_1, z_4\}_{\mathcal{Z}} &= \lambda_1^{(d)} z_1 \left(z_5 \lambda_2^{(d)} (\rho_5^{(d)} - \rho_3^{(d)}) + z_4 (\epsilon_6^{(d)} \rho_5^{(d)} - \epsilon_4^{(d)} \rho_6^{(d)}) \right. \\
&\quad \left. + z_6 \lambda_2^{(d)} (\epsilon_6^{(d)} \rho_3^{(d)} - \rho_6^{(d)}) \right) + z_4 \lambda_2^{(d)} \epsilon_2^{(d)} (z_5 \rho_5^{(d)} - z_6 \rho_6^{(d)}), \\
\{z_1, z_5\}_{\mathcal{Z}} &= \lambda_1^{(d)} z_1 \left(z_5 \lambda_2^{(d)} (\epsilon_4^{(d)} \rho_3^{(d)} - 1) + z_3 (\epsilon_6^{(d)} \rho_3^{(d)} - \rho_6^{(d)}) + z_4 (\epsilon_4^{(d)} \rho_6^{(d)} - \epsilon_6^{(d)}) \right) \\
&\quad + z_5 \epsilon_2^{(d)} \lambda_2^{(d)} (z_3 \rho_3^{(d)} - z_4), \\
\{z_1, z_6\}_{\mathcal{Z}} &= \lambda_1^{(d)} z_1 \left(z_6 \lambda_2^{(d)} (1 - \epsilon_4^{(d)} \rho_3^{(d)}) + z_3 (\rho_5^{(d)} - \rho_3^{(d)}) + z_4 (1 - \epsilon_4^{(d)} \rho_5^{(d)}) \right) \\
&\quad + z_6 \epsilon_2^{(d)} \lambda_2^{(d)} (z_4 - z_3 \rho_3^{(d)}), \\
\{z_2, z_3\}_{\mathcal{Z}} &= \lambda_1^{(d)} z_2 \left(z_5 \lambda_2^{(d)} (\epsilon_4^{(d)} \rho_5^{(d)} - 1) + z_3 (\epsilon_6^{(d)} \rho_5^{(d)} - \epsilon_4^{(d)} \rho_6^{(d)}) + \right. \\
&\quad \left. + z_6 \lambda_2^{(d)} (\epsilon_6^{(d)} - \rho_6^{(d)}) \right) + z_3 \epsilon_1^{(d)} \lambda_2^{(d)} (z_5 \rho_5^{(d)} - z_6 \rho_6^{(d)}), \\
\{z_2, z_4\}_{\mathcal{Z}} &= \lambda_1^{(d)} z_2 \left(z_5 \lambda_2^{(d)} (\rho_3^{(d)} - \rho_5^{(d)}) + z_4 (\rho_6^{(d)} - \epsilon_6^{(d)} \rho_5^{(d)}) + z_6 \lambda_2^{(d)} (\rho_6^{(d)} - \epsilon_6^{(d)} \rho_3^{(d)}) \right) \\
&\quad + z_4 \epsilon_1^{(d)} \lambda_2^{(d)} (z_6 \rho_6^{(d)} - z_5 \rho_5^{(d)}), \\
\{z_2, z_5\}_{\mathcal{Z}} &= \lambda_1^{(d)} z_2 \left(z_5 \lambda_2^{(d)} (1 - \epsilon_4^{(d)} \rho_3^{(d)}) + z_4 (\epsilon_6^{(d)} - \epsilon_4^{(d)} \rho_6^{(d)}) + z_3 (\rho_6^{(d)} - \epsilon_6^{(d)} \rho_3^{(d)}) \right) \\
&\quad + z_5 \epsilon_1^{(d)} \lambda_2^{(d)} (z_4 - z_3 \rho_3^{(d)}), \\
\{z_2, z_6\}_{\mathcal{Z}} &= \lambda_1^{(d)} z_2 \left(z_6 \lambda_2^{(d)} (\epsilon_4^{(d)} \rho_3^{(d)} - 1) + z_3 (\rho_3^{(d)} - \rho_5^{(d)}) + z_4 (\epsilon_4^{(d)} \rho_5^{(d)} - 1) \right) \\
&\quad + z_6 \epsilon_1^{(d)} \lambda_2^{(d)} (z_3 \rho_3 - z_4), \\
\{z_3, z_6\}_{\mathcal{Z}} &= \lambda_1^{(d)} (z_2 \epsilon_2^{(d)} - z_1 \epsilon_1^{(d)}) (z_6 \lambda_2 + z_3 \rho_5^{(d)}), \\
\{z_4, z_5\}_{\mathcal{Z}} &= \lambda_1 (z_2 \epsilon_2^{(d)} - z_1 \epsilon_1^{(d)}) (z_5 \lambda_2 \rho_3^{(d)} + z_4 \rho_6^{(d)}), \\
\{z_4, z_6\}_{\mathcal{Z}} &= -\lambda_1 (z_2 \epsilon_2^{(d)} - z_1 \epsilon_1^{(d)}) (z_6 \lambda_2 \rho_3^{(d)} + z_4 \rho_5^{(d)}), \\
\{z_5, z_6\}_{\mathcal{Z}} &= \lambda_1^{(d)} (z_2 \epsilon_2^{(d)} - z_1 \epsilon_1^{(d)}) (z_4 - z_3 \rho_3^{(d)}).
\end{aligned} \tag{6.6}$$

Using these formulae, by direct comparison with (6.5), we obtain

$$\{z_i, z_j\}_{\mathcal{Z}} = \frac{v_1 v_t (v_0 - 1)^2 (v_\infty^2 - 1)}{v_0 v_\infty (v_1 - 1)^2 (v_t^2 - 1)} \{z_i(y_1, \dots, y_4), z_j(y_1, \dots, y_4)\}_{\mathcal{Y}},$$

hence proving that $\Phi^{(d)}$ is Poisson. A similar computation can be done for the inverse. \square

7. CONCLUSION

In this paper, we showed that the monodromy manifold of qP_{VI} gives rise to a number of new results related to monodromy manifolds and symplectic geometry of the differential Painlevé equations. First, the continuum limit gives rise to a

Segre surface which is isomorphic to the Jimbo-Fricke cubic surface well known to be the monodromy manifold of P_{VI} . Second, we use confluence limits to obtain Segre surfaces, called \mathcal{Z} -Segre surfaces, that are isomorphic to the conventional monodromy manifolds of each Painlevé equation. Blow-downs of the latter give rise to other surfaces, which we call \mathcal{Y} -Segre surfaces. We show that the \mathcal{Z} -Segre surfaces and \mathcal{Y} -Segre surfaces are affinely equivalent and the linear affine transformation between them is Poisson. It is interesting to note that the blow-down maps from each Painlevé equation's cubic monodromy manifold to the \mathcal{Z} - and \mathcal{Y} -Segre surfaces are Poisson maps. An interesting open question is to ask whether Segre surfaces exist as monodromy manifolds for all remaining equations in Sakai's diagram.

REFERENCES

- [1] Arnol'd, V.I., Normal forms for functions near degenerate critical points, the Weyl groups of A_k , D_k , E_k and Lagrangian singularities. *Funct Anal Its Appl* **6** (1972), 254–272.
- [2] Bertola, M., Cafasso, M., Rubtsov, V., Noncommutative Painlevé equations and systems of Calogero type, *Comm. Math. Phys.* **363** (2018), 503–530.
- [3] Birkhoff, G. D., General theory of linear difference equations, *Trans. Amer. Math. Soc.*, **12** (1911), 243–284.
- [4] Borodin, A., Isomonodromy transformations of linear systems of difference equation, *Ann. of Math. (2)*, **160** (2004), no. 3:1141–1182.
- [5] Bruce, J.W., Wall, C.T.C. , “On the Classification of Cubic Surfaces”, *Journal of the London Mathematical Society* **2** (1979) 245–256.
- [6] Cayley, A., “On the triple tangent planes of surfaces of the third order”, *Cambridge and Dublin Math. J.* **4** (1849) 118–138.
- [7] Carmichael, R. D., On the Theory of Linear Difference Equations, *Amer. J. Math.*, **35** (1913), no. 2:163–182.
- [8] Cassels, J.W.S., Lectures on elliptic curves, in *Cambridge University Press*, Cambridge, **24** (1991).
- [9] Chekhov L., Mazzocco M. and Rubtsov V., Painlevé monodromy manifolds, decorated character varieties and cluster algebras, *Int. Mat. Res. Not.*
- [10] Chekhov L., Mazzocco M. and Rubtsov V., Quantised Painlevé monodromy manifolds, Sklyanin and Calabi-Yau algebras, *Advances*
- [11] Coxeter H. S. M., *Regular Polytopes*. London, Methuen & Co. Ltd (1948).
- [12] Deift P., Universality for mathematical and physical systems, in *International Congress of Mathematicians*, Vol. I, Eur. Math. Society, Zurich, 2007, [https://10.4171/022-1/7](https://doi.org/10.4171/022-1/7), 125–152.
- [13] Dolgachev I. V., *Classical algebraic geometry: A modern view*, Cambridge University Press, Cambridge (2012).
- [14] Di Francesco P., Ginsparg P., and Zinn-Justin J., 2D gravity and random matrices, *Phys. Rep.* **254** (1995), no. 1-2, 133.
- [15] Fokas, A. S.; Its, A. R.; Kapaev, A. A.; Novokshenov, V. Yu. Painlevé transcendents. The Riemann–Hilbert approach, *Mathematical Surveys and Monographs*, **128**. American Mathematical Society, Providence.
- [16] Fricke R., Klein F., Vorlesungen über die Theorie der automorphen Funktionen. I, Druck und Verlag von B.G.Teubner, Leipzig, 1897.
- [17] Gasper, G., Rahman, M., *Basic hypergeometric series*, Cambridge University Press, Cambridge, First Edition (1990).
- [18] Gross M., Hacking P., Keel S., .
- [19] Grünbaum B., *Convex Polytopes*, Springer, NY (1967).
- [20] Guzzetti, D., Tabulation of Painlevé 6 transcendents, *Nonlinearity*, **25** (2012) 3235–3276.
- [21] Hartshorne, R., *Algebraic geometry*, Springer-Verlag, New York-Heidelberg, 1977
- [22] Humphreys J. E., *Reflection Groups and Coxeter Groups*, Cambridge University Press, Cambridge UK (1992).
- [23] Iwasaki, K., A modular group action on cubic surfaces and the monodromy of the Painlevé VI equation, *Proc. Japan Acad. Ser. A Math. Sci.*, **78** (2002) 131–135.
- [24] Jimbo M., Monodromy problem and the boundary condition for some Painlevé equations, *Publ. RIMS, Kyoto Univ.*, **18** (1982) 1137–1161.
- [25] Jimbo, M. and Sakai, H., A q -analogue of the sixth Painlevé equation, *Lett. Math. Phys.*, **38** (1996) 145–154.

- [26] N. Joshi and P. Roffelsen, On the monodromy manifold of q -Painlevé VI and its Riemann-Hilbert problem, *Commun. Math. Physics*, **404** (2023) 97–149.
- [27] N. Joshi and P. Roffelsen, On the Riemann-Hilbert Problem for a q -Difference Painlevé Equation, *Commun. Math. Physics*, **384** (2021) 549–585.
- [28] Kapaev, A.A., Kitaev, A.V. Connection formulae for the first Painlevé transcendent in the complex domain. *Lett Math Phys*, **27** (19893) 243–252.
- [29] Klimeš, M., Wild monodromy of the Fifth Painlevé equation and its action on wild character variety: an approach of confluence, *Annales de l'Institut Fourier*, (2023)
- [30] Krichever, I. M., Analytic theory of difference equations with rational and elliptic coefficients and the Riemann–Hilbert problem. *Uspekhi Mat. Nauk*, **59** (2004), no. 6(360), 111–150; translation in Russian Math. Surveys 59 (2004), no. 6, 1117–1154.
- [31] Mano, T. “Asymptotic behaviour around a boundary point of the q -Painlevé VI equation and its connection problem.” *Nonlinearity*. **23** (2010): 1585–1608.
- [32] Mazzocco M., Confluences of the Painlevé equations, Cherednik algebras and q -Askey scheme, *Nonlinearity* **29** (2016), no.9:2565–2608.
- [33] Nambu, Y., Generalized Hamiltonian Dynamics, *Phys. Rev. D, Ser. 3*, **7**, 2405–2412 (1973).
- [34] Mabuchi T., Mukai S., Stability and Einstein-Kähler metric of a quartic del Pezzo surface, *Lecture Notes in Pure and Appl. Math.*, **145**, Dekker, New York, 133–160 (1993).
- [35] Noumi M., Painlevé Equations through Symmetry, American Mathematical Society, (2004).
- [36] Oblomkov A., Double Affine Hecke Algebras of Rank 1 and Affine Cubic Surfaces, *IMRN*, **2004** no.18:877–912.
- [37] Odesskii A., Rubtsov V., Polynomial Poisson algebras with regular structure of symplectic leaves, *Theoretical and Mathematical Physics*, **133**, 1321–1337 (2002)
- [38] Ohyama Y., Ramis J.P. and Sauloy J., The space of monodromy data for the Jimbo–Sakai family of q -difference equations, *Annales de la faculté des sciences de Toulouse XXIX*, 1119–1250 (2020)
- [39] Okamoto K., Sur les feuilletages associés aux équations du second ordre à points critiques fixes de P. Painlevé Japan. J. Math. (N.S.) **5**(1) (1979) 1–79.
- [40] Ramis, J.P. and Sauloy, J., Geometry of the space of monodromy data, arXiv:2301.04083 [math.DS].
- [41] Ramis, J. P., Sauloy, J. and Zhang, C., Développement asymptotique et sommabilité des solutions des équations linéaires aux q -différences, *C. R. Math. Acad. Sci. Paris*, **342** (2006), no. 7:515–518.
- [42] P. Roffelsen, On q -Painlevé VI and the geometry of Segre surfaces, arXiv:2305.17912 [nlin.SI].
- [43] Sakai H., Rational Surfaces Associated with Affine Root Systems and Geometry of the Painlevé Equations *Commun. Math. Phys.* **220**, 165–229 (2001).
- [44] Salmon, G., “On the triple tangent planes to a surface of the third order”, Cambridge and Dublin Math. J. **4** (1849) 252-260.
- [45] Sauloy, J., Systèmes aux q -différences singuliers réguliers: classification, matrice de connexion et monodromie, *Ann. Inst. Fourier (Grenoble)*, **50** (2000), no. 4:1021–1071.
- [46] Takhtajan, L., On foundation of the generalized Nambu mechanics, *Comm. Math. Phys.*, **160**, 295–315 (1994).
- [47] Tracy, C. A. and Widom, H., Level-spacing distributions and the Airy kernel, *Communications in Mathematical Physics*, **159** (1), 1994, 151–174.
- [48] Segre, C., “Etude des différentes surfaces du 4e ordre à conique double ou cuspidale (générale ou décomposée) considérées comme des projections de l’intersection de deux variétés quadratiques de l’espace à quatre dimensions”, *Mathematische Annalen* **24** (1884) 313-444.
- [49] Saito M., van der Put M., Moduli spaces for linear differential equations and the Painlevé equations, *arXiv:0902.1/02v5* (2009).

SCHOOL OF MATHEMATICS AND STATISTICS F07, THE UNIVERSITY OF SYDNEY, NEW SOUTH WALES 2006, AUSTRALIA.

Email address: nalini.joshi@sydney.edu.au

SCHOOL OF MATHEMATICS, UNIVERSITY OF BIRMINGHAM, UK.

Email address: m.mazzocco@bham.ac.uk

SCHOOL OF MATHEMATICS AND STATISTICS F07, THE UNIVERSITY OF SYDNEY, NEW SOUTH WALES 2006, AUSTRALIA.

Email address: pieter.roffelsen@sydney.edu.au